Follow-up of infants with cow's milk allergy undergoing early oral immunotherapy

Laura Badina¹, Valentina Carrato², Laura Levantino², Giulia Peruch², Egidio Barbi³, Irene Berti¹, and Giorgio Longo⁴

¹Institute for Maternal and Child Health IRCCS Burlo Garofolo ²University of Trieste Clinical Department of Medical Surgical and Health Sciences ³Institute for Maternal and Child Health, IRCCS Burlo Garofolo ⁴Institute for Maternal and Child Health IRCCS Burlo Garofolo

July 22, 2020

Hosted file

Letter to the Editor.pdf available at https://authorea.com/users/345306/articles/471556-follow-up-of-infants-with-cow-s-milk-allergy-undergoing-early-oral-immunotherapy

1 Table 1. Patients' characteristics at baseline and current follow-up studies

2

4

3	OIT-Oral Immunotheren	v: IOR=Interquartile Ranges	CM=Cour's Milk

	Baseline study cohort [4]		Follow-up cohort outcome (median duration, months: 39, IQR 32-47)
Number of patients	73		62
Male patients, number (%)	50/73 (68)		42/62 (68)
Median age, years (IQR)	0.6 (0.2-0.9)		4 (2.8-4.7)
OIT, number (%) Successfully concluded Suspended Not started for ineligibility	66/73 (90) 2/73 (3) 5/73 (7)		56/62 (90) 1/62 (2) 5/62 (8)
Type of diet, number (%) Unrestricted	66/73 (90)	Type of diet, number (%) Unrestricted	59/62 (95)
of which concluded OIT suspended OIT not eligible for OIT	66 0 0	of which concluded previous OIT suspended previous OIT previously not eligible for OIT	56 1 2
CM-free, number (%)	7/73 (10)	CM-free, number (%)	3/62 (5)
of which concluded OIT	0	of which concluded previous OIT	0
suspended OIT not eligible for OIT	2 5	suspended previous OIT previously not eligible for OIT	0 3

1