

Enhanced airway hyperresponsiveness in asthmatic children and mice with A(H1N1)pdm09 infection

Taira Ariyoshi¹, Junichiro Tezuka², Hiroki Yasudo¹, Tamaki Nakamura¹, Yasufumi Sakata¹, Takeshi Matsushige¹, Hideki Hasegawa³, Noriko Nakajima³, Akira Aina³, Atsunori Oga¹, Hiroshi Itoh¹, Komei Shirabe⁴, Shoichi Toda⁴, Ryo Atsuta⁵, Shouichi Ohga¹, and Shunji Hasegawa¹

¹Yamaguchi University Graduate School of Medicine

²Fukuoka Children's Hospital

³National Institute of Infectious Diseases

⁴Yamaguchi Prefectural Institute of Public Health and Environment

⁵Akihabara Atsuta Clinic

September 16, 2020

Abstract

Background: Severe asthma exacerbation is an important comorbidity of the 2009 H1N1 pandemic [A(H1N1)pdm09] in asthmatic patients. However, the mechanisms underlying severe asthma exacerbation remain unknown. In this study, airway hyperresponsiveness (AHR) was measured in paediatric asthma patients infected with A(H1N1)pdm09. We also evaluated AHR in asthmatic mice with A(H1N1)pdm09 infection and those with seasonal influenza for comparison. **Methods:** AHRs in asthmatic children were defined as the provocative acetylcholine concentration causing a 20% reduction in FEV_{1.0} (PC₂₀). To investigate the pathophysiology using animal models, BALB/c mice aged 6-8 weeks were sensitized and challenged with ovalbumin. Either mouse-adapted A(H1N1)pdm09, seasonal H1N1 virus (1×10⁵ pfu/20 µL), or mock treatment as a control was administered intranasally. At 3, 7, and 10 days after infection, each group of mice was evaluated for AHR by methacholine challenge using an animal ventilator, flexiVent®. Lung samples were resected and observed using light microscopy to assess the degree of airway inflammation. **Results:** AHRs in the children with bronchial asthma were temporarily increased, and alleviated by 3 months after discharge. AHR was significantly enhanced in A(H1N1)pdm09-infected asthmatic mice compared to that in seasonal H1N1-infected mice (p<0.001), peaking at 7 days post-infection and then becoming similar to control levels by 10 days post-infection. Histopathological examination of lung tissues showed more intense infiltration of inflammatory cells and severe tissue destruction in A(H1N1)pdm09-infected mice at 7 days post-infection than at 10 days post-infection. **Conclusions:** Our results suggest that enhanced AHR could contribute to severe exacerbation in human asthmatic patients with A(H1N1)pdm09 infection.

Introduction

Asthma exacerbation is a major cause of disease morbidity that increases health care costs, and in some patients, progressive loss of lung function¹. Exposure to aeroallergens and environmental factors, such as smoking, PM2.5, or diesel exhaust particles, triggers asthma exacerbation^{2,3}. Respiratory viral infection is also associated with the pathophysiology of asthma exacerbation, particularly in childhood⁴. The most prominent pathogens involved in asthma exacerbation include human rhinovirus, respiratory syncytial virus, enterovirus, influenza virus, and human metapneumovirus⁵⁻⁷.

Many severe and fatal cases of the 2009 H1N1 pandemic [A(H1N1)pdm09] infection have been reported, both in patients with underlying diseases as well as in healthy children and young adults⁸⁻¹⁰. Asthma is

among the most common underlying conditions in patients hospitalized with A(H1N1)pdm09 infection and asthmatic children show greater susceptibility to A(H1N1)pdm09 viral infection¹¹, suggesting that severe asthma exacerbation is an important comorbidity of influenza infection in patients with asthma⁸⁻¹⁰.

We previously reported that A(H1N1)pdm09 infection, but not seasonal H1N1 infection, induces severe pulmonary inflammation with elevated cytokine levels in a mouse model of asthma^{12,13}. Moreover, asthmatic model mice with A(H1N1)pdm09 infection are prone to an earlier onset of severe pulmonary inflammation compared to those with seasonal H1N1 infection¹⁴, suggesting that a hyper-cytokine condition is involved in severe pneumonia and atelectasis. Another report showed that A(H1N1)pdm09 induces AHR in a non-asthmatic mouse model¹⁵. However, no reports have evaluated the effect of A(H1N1)pdm09 on airway constriction in patients with asthma and, to date, the mechanisms of severe asthma exacerbation due to A(H1N1)pdm09 infection remain unclear.

In this study, we showed that pediatric patients with A(H1N1)pdm09 infection temporarily increased airway hyperresponsiveness (AHR) as well as asthmatic mice. Furthermore, we showed that the changes in AHR are greater in asthmatic mice with A(H1N1)pdm09 infection than those with seasonal H1N1 influenza. These findings indicate that enhanced AHR contributes to the severe asthma exacerbation phenotype triggered by A(H1N1)pdm09 infection.

Methods

Measurement of AHR in children with A(H1N1)pdm09 infection

Studies on pediatric patients were carried out in accord with the Declaration of Helsinki. This study enrolled pediatric asthmatic patients with A(H1N1)pdm09 infection and hypoxia (SpO_2 [?]90%) within the first day of fever admitted to Fukuokahigashi Medical Center between September 2009 and December 2010. Bronchial asthma was diagnosed according to the Japanese Pediatric Guidelines for the Treatment and Management of Bronchial Asthma 2008 (JPGL 2008)¹⁵. The participants were diagnosed with A(H1N1)pdm09 infection by polymerase chain reaction (PCR) at Fukuoka Institute of Health and Environmental Sciences. AHRs were measured at 1 and 3 months after discharge. Anti-inflammatory drugs and bronchodilators were prohibited 1 month after discharge. Increasing concentrations of acetylcholine (39, 78, 156, 312, 625, 1,250, 2,500, 5,000, 10,000, and 20,000 $\mu\text{g/mL}$) were inhaled by a nebulizer until the forced expiratory volume in 1 second ($\text{FEV}_{1.0}$) was reduced by 20% from a post-nebulized saline value. $\text{FEV}_{1.0}$ was measured using a spirometer (HI-801, CHEST M.I., Inc., Tokyo, Japan). AHR was defined as the provocative concentration causing a 20% fall in $\text{FEV}_{1.0}$ (PC_{20}). This study was approved by the institutional review board of Fukuokahigashi Medical Center (2020-rin-8). Parents were explained this study and signed informed consent.

Sensitization and allergen challenge in a mouse model of bronchial asthma

BALB/c mice aged 6–8 weeks were obtained from Chiyoda Kaihatsu Co. (Tokyo, Japan) and sensitized and challenged with grade II ovalbumin (Sigma–Aldrich., St. Louis, MO, USA) as previously described¹²⁻¹⁴. All animal experimentation procedures were approved by the Institutional Animal Care and Use Committee of Yamaguchi University (No. 29-S01), and all methods were conducted in accordance with approved guidelines. Additionally, animal experiments conformed to the revised Institute of Laboratory Animal Resources, Commission on Life Sciences, National Research Council “Guide for the Care and Use of Laboratory Animals” published by the National Academy Press, Washington, D.C. 1996.

Virus infection

Mouse-adapted A(H1N1)pdm09 (strain: A/Narita/1/09) or seasonal H1N1 (strain: A/Puerto Rico) viruses were provided by the National Institute of Infectious Diseases (Tokyo, Japan). On day 31, influenza virus (concentration: 1×10^5 pfu/20 μL) or vehicle (mock-infection) was administered intranasally to the mice. The number of mice in each group ranged from 4 to 7.

Measurement of AHR

Mice were anesthetized by intraperitoneal injection of xylazine (12 mg/kg) and pentobarbital (70 mg/kg) at 3, 7, and 10 days post-infection. After anesthetization, mice were cannulated and connected to an animal ventilator (flexiVent®, SCIREQ, Montreal, QC, Canada). The mice inhaled aerosolized phosphate-buffered saline or 3, 6, 12, 24, or 48 mg/mL methacholine in phosphate-buffered saline. We measured the following parameters: resistance of the respiratory system (R_{rs}) and tissue damping (G); R_{rs} and G reflect the resistance of the central and peripheral airways, respectively^{17, 18}. These parameters were measured at frequent intervals of 10–15 seconds 12 times following activation of the nebulizer.

Histological examination of the lungs

After the methacholine challenge, the lung was resected from mice that were euthanized with high dose pentobarbital (200 mg/kg). Lung tissues were fixed in 10% buffered formalin for 24 h at room temperature and then embedded in paraffin. Serial sections (3- μ m-thick) were cut and stained with haematoxylin and eosin (Muto Pure Chemicals Co., Tokyo, Japan).

Statistical analysis

For human data, differences between groups were analysed using the Wilcoxon signed rank test. For mouse data, differences between two groups and between three groups were analysed using the Mann–Whitney U test and Steel–Dwass test, respectively. When p-values less than 0.05, differences between means were considered to be statistically significant. All analyses and calculations were performed using JMP® Pro version 13.0.0 software (SAS Institute, Inc., Cary, NC, USA).

Results

Measurement of AHR in children with A(H1N1)pdm09 infection

To elucidate whether AHR was enhanced in patients with A(H1N1)pdm09 infection, we measured AHR of 12 pediatric participants infected with A(H1N1)pdm09 at 1- and 3-months after discharge (**Table 1**). PC_{20} of all patients were lower than 8,000 μ g/mL at 1month after discharge. PC_{20} significantly increased at 3 months after discharge compared to that at 1 month after discharge, suggesting that AHR was enhanced in the acute phase (1 month post discharge) of these patients (**Fig. 1** ; 1 month after discharge vs. 3 months after discharge; 1,036 vs. 1,597 μ g/mL, $p = 0.009$).

Enhanced AHR in asthmatic mice with A(H1N1)pdm09 infection

The central airway resistance revealed by the R_{rs} value was significantly increased in the A(H1N1)pdm09 mouse group compared to that in the seasonal H1N1 mouse or control group (**Fig. 2**); this was particularly prominent after treatment with 48 mg/mL methacholine at 3 and 7 days after infection [3 days post-infection A(H1N1)pdm09 vs. seasonal; 4.40 vs. 3.29 $\text{cmH}_2\text{O.s/mL}$, $p < 0.001$, vs. control; vs. 3.11 $\text{cmH}_2\text{O.s/mL}$, $p < 0.001$; 7 days post-infection A(H1N1)pdm09 vs. seasonal; 8.60 vs. 3.24 $\text{cmH}_2\text{O.s/mL}$, $p < 0.001$, vs. control; vs. 3.00 $\text{cmH}_2\text{O.s/mL}$, $p < 0.001$]. However, there was no significant difference among the three groups at 10 days post-infection. In contrast, there were no significant differences in the R_{rs} between the seasonal H1N1 and mock groups at 3, 7, or 10 days post-infection.

Next, peripheral airway resistance as reflected by G was compared among the three groups (**Fig. 3**). The peripheral airway resistance was also significantly increased in the A(H1N1)pdm09 group compared to that in the seasonal H1N1 and control groups; this difference was particularly prominent after treatment with 48 mg/mL methacholine at 3 and 7 days post-infection [3 days post-infection A(H1N1)pdm09 vs. seasonal; 25.2 vs. 16.5 $\text{cmH}_2\text{O/mL}$, $p < 0.001$, vs. control; vs. 15.5 $\text{cmH}_2\text{O/mL}$, $p < 0.001$; 7 days post-infection A(H1N1)pdm09 vs. seasonal; 50.0 vs. 16.7 $\text{cmH}_2\text{O/mL}$, $p < 0.001$, vs. control; vs. 15.6 $\text{cmH}_2\text{O/mL}$, $p < 0.001$]. However, these differences in airway resistance were not significantly different among the three groups at 10 days post-infection.

We investigated the changes in AHR of non-asthmatic mice with A(H1N1)pdm09 infection (**Figs. 2, 3**). Although AHR was enhanced with A(H1N1)pdm09 infection in non-asthmatic mice, the changes in AHR

were slight compared to the changes observed in asthmatic mice, suggesting that A(H1N1)pdm09 infection more robustly enhances AHR in asthmatic animals.

We further evaluated the alternations of AHR in asthmatic mice during the post-infection period of A(H1N1)pdm09 infection (**Fig. 4**). Airway resistance was significantly enhanced at 7 days post-infection compared to at 3- or 10-days post-infection ($p < 0.001$), whereas there were no differences in airway resistance between 3 and 10 days post-infection. When the body weight of mice was compared among the three groups to evaluate the systemic damage caused by A(H1N1)pdm09 infection at 3, 7, or 10 days post-infection, no significant differences between days were detected (data not shown).

Histopathological findings in the lungs

Haematoxylin and eosin staining of lung tissues from mice at 3, 7, and 10 days post-infection is shown in **Fig. 5**. Inflammatory cell infiltration was prominently observed in A(H1N1)pdm09-infected mice, whereas this was rarely observed in mice infected with seasonal H1N1 at 3 days post-infection (**Fig. 5A**). The lung inflammation was clearly observed in A(H1N1)pdm09- and seasonal H1N1-infected mice at 7 days post-infection, but was more severe in A(H1N1)pdm09-infected mice (**Fig. 5B**). In contrast, the magnitude of inflammation was attenuated by 10 days post-infection (**Fig. 5C**).

Discussion

In this study, we measured temporal changes of PC₂₀ in pediatric asthma patients with A(H1N1)pdm09 infection. PC₂₀ was lowest at 1 month after discharge and significantly increased at 3 months after discharge. Furthermore, we comparatively evaluated AHR in a mouse model of asthma with either A(H1N1)pdm09 infection or seasonal H1N1 infection to investigate the pathophysiology of A(H1N1)pdm09-infected asthmatic children. Enhanced AHR was observed in asthmatic mice with A(H1N1)pdm09 infection, which peaked at 7 days post-infection and subsequently diminished at 10 days post-infection. These results were similar to those of asthmatic children. Histopathological analysis showed that the onset of lung inflammation in asthmatic mice with A(H1N1)pdm09 infection occurred earlier and was more prominent compared to that in mice with seasonal H1N1 infection; these effects peaked 7 days post-infection and diminished by 10 days post-infection, which was consistent with the observed changes in AHR. These data suggest that A(H1N1)pdm09 induces enhanced AHR complication as a severe phenotype of pneumonia in mice with asthma. The desaturation and enhanced AHR observed in asthmatic patients must be induced by the pulmonary inflammation during A(H1N1)pdm09 infection, but not seasonal H1N1.

The severity of AHR reflects the inflammatory state of the airways¹⁹. Several studies have reported that AHR can be enhanced by inflammatory and Th2 cytokines, such as tumour necrosis factor (TNF)- α , interleukin (IL)-6, and IL-13²⁰⁻²². TNF- α secreted from airway macrophages or airway epithelial cells after respiratory virus infection increases levels of adhesion molecules, such as intercellular adhesion molecule-1 on epithelial cells, thereby inducing the recruitment of eosinophils and contributing to epithelial damage and AHR²³⁻²⁷. TNF- α is associated with wheezing in human infants²⁰. IL-6 is secreted from epithelial cells in respiratory virus infection and induces airway inflammation and bronchospasms in patients with asthma and upper respiratory tract infections^{21, 24}. Our previous study showed that IL-6 and TNF- α levels in the bronchoalveolar lavage fluid of A(H1N1)pdm09-infected mice were significantly higher than those in seasonal H1N1-infected mice within 3 days after infection¹⁴. Therefore, A(H1N1)pdm09 infection may enhance AHR by inducing the production of high levels of inflammatory cytokines during lung inflammation in asthmatic mice, which may also occur in human cases.

As explained above, AHR in A(H1N1)pdm09-infected children was alleviated by 3 months after discharge compared to findings at 1 month after discharge. This finding is supported by our data in A(H1N1)pdm09-infected mice showing that the enhanced AHR at 3 and 7 days post-infection decreased to the same level as in control mice at 10 days post-infection. Bozanich et al²⁸ reported that increased AHR observed in seasonal H3N1 influenza-infected non-asthmatic wild-type mice at 4 days post-infection returned to control levels at 20 days post-infection, which is consistent with our findings. Together, these data indicate that it is pivotal to treat patients with severe asthma exacerbation in the acute phase of post-A(H1N1)pdm09 infection. Estab-

lished treatments for rescuing acute severe asthma exacerbation complicated with severe pneumonia resulting from A(H1N1)pdm09 infection have not yet been developed. We are currently investigating approaches for treating acute severe asthma exacerbation occurring with A(H1N1)pdm09 infection.

There were some limitations to this study. First, we did not evaluate AHR, inflammatory or Th2 cytokines, or virus titres in the bronchoalveolar lavage fluid at the same time. Lung tissues were collected and used for pathological analysis after AHR evaluation, as we previously reported the cytokine profiles in the bronchoalveolar lavage fluid of A(H1N1)pdm09 mice¹³. Second, we could not evaluate AHRs of the pediatric participants before or during A(H1N1)pdm09 infection for ethical reasons to verify whether or not enhanced AHR during the infection were alleviated in the post-infection phase. Instead, we measured AHR at 1 and 3 months after discharge.

In conclusion, AHR was significantly enhanced in both pediatric asthma patients and asthmatic mice in acute phase of A(H1N1)pdm09 infection. Furthermore, A(H1N1)pdm09-infected asthma model mice showed more severe pulmonary inflammation in the acute phase post-infection, compared to that occurring in asthmatic mice with seasonal influenza infection. Enhanced AHR subsequently returned to normal levels with the amelioration of lung inflammation, suggesting that appropriate treatment during the acute phase after A(H1N1)pdm09 infection is essential for avoiding severe respiratory conditions.

Acknowledgments

This work was supported by Grant-in-Aid for Translational Research of Yamaguchi University Hospital 2013. We thank MT Kenzo Ikemoto (Department of Molecular Pathology, Yamaguchi University Graduate School of Medicine) for providing technical support.

Key Message

Pediatric asthma patients showed enhanced signs of airway hyperresponsiveness at 1 versus 3 months post hospital discharge. This study also used mouse models of asthma-disposed mice to test whether pandemic or seasonal strains of the influenza could enhance airway hyperresponsiveness. The pandemic A(H1N1)pdm09 strain significantly worsened airway inflammation and function compared to the seasonal flu, although these effects were limited to the acute phase at 7 days post infection. By 10 days post infection, the resolution phase was reached and differences between groups of mice infected with different influenza virus strains were negligible. In summary, our study indicates that acute influenza infection can exacerbate asthma phenotypes, particularly pandemic strains that cause severe immunopathology.

References

1. Chipps BE, Szefer SJ, Simons FE, Haselkorn T, Mink DR, Deniz Y, Lee JH; TENOR Study Group. Demographic and clinical characteristics of children and adolescents with severe or difficult-to-treat asthma. *J Allergy Clin Immunol* 2007;119:1156-1163.
2. Sheehan WJ, Phipatanakul W. Indoor allergen exposure and asthma outcomes. *Curr Opin Pediatr* 2016;28:772-777.
3. Guarnieri M, Balmes JR. Outdoor air pollution and asthma. *Lancet* 2014;383:1581-1592.
4. Jartti T, Gern JE. Role of viral infections in the development and exacerbation of asthma in children. *J Allergy Clin Immunol* 2017;140:895-906.
5. Abe N, Yasudo H, Fukano R, Nakamura T, Okada S, Wakiguchi H, Okazaki F, Shirabe K, Toda S, Okamoto R, Ouchi K, Ohga S, Hasegawa S. Multi-season analyses of causative pathogens in children hospitalized with asthma exacerbation. *Pediatr Allergy Immunol.* 2019;30:724-731.
6. Zheng XY, Xu YJ, Guan WJ, Lin LF. Regional, age and respiratory-secretion-specific prevalence of respiratory viruses associated with asthma exacerbation: a literature review. *Arch Virol* 2018;163:845-853.
7. Wark PAB, Ramsahai JM, Pathinayake P, Malik B, Bartlett NW. Respiratory viruses and asthma. *Semin Respir Crit Care Med* 2018;39:45-55.

8. Hasegawa S, Hirano R, Hashimoto K, Haneda Y, Shirabe K, Ichiyama T. Characteristics of atopic children with pandemic H1N1 influenza viral infection: pandemic H1N1 influenza reveals 'occult' asthma of childhood. *Pediatr Allergy Immunol* 2011;22:e119-e123.
9. Athanasiou M, Lytras T, Spala G, Triantafyllou E, Gkolfinopoulou K, Theocharopoulos G, Patrinos S, Danis K, Detsis M, Tsiodras S, Bonovas S, Panagiotopolous T. Fatal cases associated with pandemic influenza A (H1N1) reported in Greece. *PLoS Curr* 2010;2:RRN1194.
10. Reichert T, Chowell G, Nishiura H, Christensen RA, McCullers JA. Does glycosylation as a modifier of original antigenic sin explain the case age distribution and unusual toxicity in pandemic novel H1N1 influenza? *BMC Infect Dis* 2010;10:5.
11. Jain S, Kamimoto L, Bramley AM, Schmitz AM, Benoit SR, Louie J, Sugerman DE, Druckenmiller JK, Ritger KA, Chugh R, Jasuja S, Deutscher M, Chen S, Walker JD, Duchin JS, Lett S, Soliva S, Wells EV, Swardlow D, Uyeki TM, Fiore AE, Olsen SJ, Fry AM, Bridges CB, Finelli L. 2009 Pandemic Influenza A (H1N1) Virus Hospitalization Investigation Team. Hospitalized patients with 2009 H1N1 influenza in the United States, April-June 2009. *N Engl J Med* 2009;61:1935-1944.
12. Okada S, Hasegawa S, Hasegawa H, Aina A, Atsuta R, Ikemoto K, Sasaki K, Toda S, Shirabe K, Takahara M, Harada S, Morishima T, Ichiyama T. Analysis of bronchoalveolar lavage fluid in a mouse model of bronchial asthma and H1N1 2009 infection. *Cytokine* 2013;63:194-200.
13. Hasegawa S, Wakiguchi H, Okada S, Gui Kang Y, Fujii N, Hasegawa M, Hasegawa H, Aina A, Atsuta R, Shirabe K, Toda S, Wakabayashi-Takahara M, Morishima T, Ichiyama T. Cytokine profile of bronchoalveolar lavage fluid from a mouse model of bronchial asthma during seasonal H1N1 infection. *Cytokine* 2014;69:206-210.
14. Fujimoto Y, Hasegawa S, Matsushige T, Wakiguchi H, Nakamura T, Hasegawa H, Nakajima N, Aina A, Oga A, Itoh H, Shirabe K, Toda S, Atsuta R, Morishima T, Ohga S. Pulmonary inflammation and cytokine dynamics of bronchoalveolar lavage fluid from a mouse model of bronchial asthma during A(H1N1)pdm09 influenza infection. *Sci Rep* 2017;7:9128.
15. Shim DH, Park YA, Kim MJ, Hong JY, Baek JY, Kim KW, Byun YH, Seong BL, Ryu S, Song MK, Hong KJ, Na W, Song D, Park JH, Sohn MH, Lee JM. Pandemic influenza virus, pH1N1, induces asthmatic symptoms via activation of innate lymphoid cells. *Pediatr Allergy Immunol* 2015;26:780-788.
16. Kondo N, Nishimuta T, Nishima S, Morikawa A, Aihara Y, Akasaka T, Akasawa A, Adachi Y, Arakawa H, Ikarashi T, Ikebe T, Inoue T, Iwata T, Urisu A, Ebisawa M, Ohya Y, Okada K, Odajima H, Katsunuma T, Kameda M, Kurihara K, Kohno Y, Sakamoto T, Shimojo N, Suehiro Y, Tokuyama K, Nambu M, Hamasaki Y, Fujisawa T, Matsui T, Matsubara T, Mayumi M, Mukoyama T, Mochizuki H, Yamaguchi K, Yoshihara S. Japanese pediatric guidelines for the treatment and management of bronchial asthma 2008. *Pediatr Int.* 2010;52:319-26.
17. McGovern TK, Robichaud A, Fereydoonza L, Schuessler TF, Martin JG. Evaluation of respiratory system mechanics in mice using the forced oscillation technique. *J Vis Exp* 2013;75:e50172.
18. Siddiqui S, Tsuchiya K, Risse PA, Bullimore SR, Benedetti A, Martin JG. Site of allergic airway narrowing and the influence of exogenous surfactant in the Brown Norway rat. *PLoS One* 2012;7:e29381.
19. Postma DS, Kerstjens HA. Characteristics of airway hyperresponsiveness in asthma and chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1998;158:S187-S192.
20. Balfour-Lynn IM, Valman HB, Wellings R, Webster AD, Taylor GW, Silverman M. Tumour necrosis factor-alpha and leukotriene E4 production in wheezy infants. *Clin Exp Allergy* 1994;24:121-126.
21. Yasuda H, Suzuki T, Zayasu K, Ishizuka S, Kubo H, Sasaki T, Nishimura H, Sekizawa K, Yamaya M. Inflammatory and bronchospastic factors in asthma exacerbations caused by upper respiratory tract infections. *Tohoku J Exp Med* 2005;207:109-118.
22. Risse PA, Jo T, Suarez F, Hirota N, Tolloczko B, Ferraro P, Grutter P, Martin JG. Interleukin-13 inhibits proliferation and enhances contractility of human airway smooth muscle cells without change in contractile phenotype. *Am J Physiol Lung Cell Mol Physiol* 2011;300:L958-L966.
23. Gern JE, Dick EC, Lee WM, Murray S, Meyer K, Handzel ZT, Busse WW. Rhinovirus enters but does not replicate inside monocytes and airway macrophages. *J Immunol* 1996;156:621-627.
24. Yamaya M. Virus infection-induced bronchial asthma exacerbation. *Pulm Med* 2012;2012:834826.

25. Rothlein R, Czajkowski M, O'Neill MM, Marlin SD, Mainolfi E, Merluzzi VJ. Induction of intercellular adhesion molecule 1 on primary and continuous cell lines by pro-inflammatory cytokines. Regulation by pharmacologic agents and neutralizing antibodies. J Immunol 1988;141:1665-1669.
26. Wegner CD, Gundel RH, Reilly P, Haynes N, Letts LG, Rothlein R. Intercellular adhesion molecule-1 (ICAM-1) in the pathogenesis of asthma. Science 1990;247:456-459.
27. Malaviya R, Laskin JD, Laskin DL. Anti-TNF α therapy in inflammatory lung diseases. Pharmacol Ther. 2017;180:90-98.
28. Bozanich E, Gualano RC, Zosky GR, Larcombe AN, Turner DJ, Hantos Z, Sly PD. Acute Influenza A infection induces bronchial hyper-responsiveness in mice. Respir Physiol Neurobiol 2008;162:190-196.

Table 1. Clinical characteristics of patients infected with A(H1N1)pdm09 at 1 month after discharge. FVC, forced vital capacity; FEV_{1.0}, forced expiratory volume in 1 s; V₅₀, maximal flow at 50% vital capacity.

n=12	
Age (years)	7.7 (4.9–11.8)
Sex (M/F)	9/3
IgE (IU/mL)	932 (124–2,680)
%FVC	90.6 (70.0–119)
%FEV _{1.0}	96.4 (75.8–111)
%V ₅₀	97.2 (55.5–160)
Previous asthma diagnosis	7

Figure legends

Figure 1. The provocative concentration causing a 20% reduction in the forced expiratory volume in 1 second (PC₂₀) at 1 and 3 months after discharge. *: children with previous asthma diagnosis; *children without previous asthma diagnosis*.


Figure 2. Resistance of respiratory system (R_{rs}) to methacholine challenge in asthma model mice infected with vehicle (mock), A(H1N1)pdm09, or seasonal H1N1 (A/Puerto Rico) at 3 (A), 7 (B), and 10 (C) days post-infection. Data are presented as means \pm standard error of the mean (SEM); *: asthma – mock; : asthma – A(H1N1)pdm09; *asthma – seasonalH1N1*; *non – asthma – A(H1N1)pdm09*. Differences between means are presented as follows : *asthma – mock vs. asthma – seasonalH1N1*, * $p < 0.05$; *asthma – seasonalH1N1 vs. asthma – A(H1N1)pdm09*, + $p < 0.05$; *asthma – A(H1N1)pdm09 vs. asthma – mock*, + + $p < 0.05$; and *asthma – A(H1N1)pdm09 vs. non – asthma – A(H1N1)pdm09*, SS $p < 0.05$.

Figure 3. Tissue damping (G) of methacholine challenge in asthma model mice infected with vehicle (mock), A(H1N1)pdm09, or seasonal H1N1 (A/Puerto Rico) at 3 (A), 7 (B), and 10 (C) days post-infection. Data are presented as means \pm standard error of the mean (SEM); *: asthma – mock; : asthma – A(H1N1)pdm09; *asthma – seasonalH1N1*; *non – asthma – A(H1N1)pdm09*. Differences between means are presented as follows : *asthma – mock vs. asthma – seasonalH1N1*, * $p < 0.05$; *asthma – seasonalH1N1 vs. asthma – A(H1N1)pdm09*, + $p < 0.05$; *asthma – A(H1N1)pdm09 vs. asthma – mock*, + + $p < 0.05$; and *asthma – A(H1N1)pdm09 vs. non – asthma – A(H1N1)pdm09*, SS $p < 0.05$.

Figure 4. Changes in airway hyperresponsiveness of A(H1N1)pdm09-infected mice at 3, 7, and 10 days post-infection. (A) Resistance of respiratory system (R_{rs}) to methacholine challenge. (B) Tissue damping (G) of methacholine challenge. Data are presented as means \pm standard error of the mean (SEM); *: day 3, : day 7, *day10*. Differences between means are presented as follows : *day3 vs. day7*, * $p < 0.05$; *day3 vs. day10*, + $p < 0.05$; and *day7 vs. day10*, + + $p < 0.05$.

Figure 5. Histopathological findings after influenza infection. Photomicrographs of haematoxylin and eosin-stained lung tissue at 3 (A), 7 (B), and 10 (C) days post-infection with mock, seasonal H1N1 (A/Puerto

Rico), or A(H1N1)pdm09 influenza virus. Representative data are shown for tissues from one of four to six independent mice per group.



Hosted file

Figure1-5.pdf available at <https://authorea.com/users/349602/articles/481470-enhanced-airway-hyperresponsiveness-in-asthmatic-children-and-mice-with-a-h1n1-pdm09-infection>