

Small gaps are preferred for the scatter hoarding of acorns by rodents

Fei Yu¹, Shanshan Wei¹, Yang Wang¹, Yingchen Li¹, Lixia Zhang¹, Qingwei Lin¹, Jianmin Ma¹, and Xianfeng Yi²

¹Henan Normal University

²Qufu Normal University

October 20, 2020

Abstract

In general, it is accepted that gap formation significantly affects the placement of scatter-hoarded seeds by small rodents, but the effects of different forest gap sizes on the seed-eating and scatter-hoarding behaviors of small rodents remain unclear. Thus, we examined the effects of a closed canopy forest, forest edge, and gaps with different sizes on the spatial dispersal of *Quercus variabilis* acorns and cache placement by small rodents using coded plastic tags in the Taihang Mountains, China. The seeds were removed rapidly and there were significant differences in the seed-eating and caching strategies between the stand types. We found that *Q. variabilis* acorns were usually eaten after being removed from the closed canopy forest and forest edges. By contrast, the *Q. variabilis* acorns in the forest gap stands were more likely to be scatter hoarded. The dispersal distances of *Q. variabilis* acorns were significantly longer in the forest gap plots compared with the closed canopy and forest edge plots. However, the proportions of scatter-hoarded seeds did not increase significantly as the gap size increased. In small-scale oak reforestation projects or research, creating small gaps to promote rodent-mediated seed dispersal may effectively accelerate forest recovery and successional processes.

Hosted file

manuscript.pdf available at <https://authorea.com/users/368810/articles/487832-small-gaps-are-preferred-for-the-scatter-hoarding-of-acorns-by-rodents>