

Evidencias competencia 5

Omar Escamilla García.¹

¹Instituto Tecnológico Superior Zacatecas Occidente

18 de junio de 2020

Problemas.

Problema 1.

A person jumps from a fourth-story window 15.0 m above a firefighter's safety net. The survivor stretches the net 1.0 m before coming to rest, Fig. 46. (a) What was the average deceleration experienced by the survivor when she was slowed to rest by the net? (b) What would you do to make it "safer" (that is, to generate a smaller deceleration): would you stiffen or loosen the net? Explain.




Figura 1: This is a caption

A

$$Y = y_o + V_o t - \frac{1}{2} g t^2$$

$$V^2 = V_o^2 - 2g(y - y_o)$$

$$V = v_o - gt$$

Solución:

Primero se calcula la velocidad con la que toca la red.

$$V^2 = (0)^2 - 2(9.81)(0 - 15m)$$

$$2\left(9.81 \frac{m}{s}\right)(15m)$$

$$V = \sqrt{2(9.81) \frac{m^2}{s}}(15m)$$

$$V = 17.15 \frac{m}{s}$$

Aceleración

$$V^2 = V_o^2 - 2a(y - y_o)$$

$$2a(y - y_o) = v^2 - v_o^2$$

$$2a = \frac{v^2 - v_o^2}{y - y_o}$$

Variables:

$$V_o = 17.5 \frac{m}{s} \quad V = 0 \frac{m}{s} \quad Y_o = 1m \quad Y = 0m$$

$$a = \frac{\left(0 \frac{m}{s}\right)^2 - \left(17.5 \frac{m}{s}\right)^2}{2(0 - 1m)} = \frac{\left(17.5 \frac{m}{s}\right)^2}{-2}$$

$$a = 147 \frac{m}{s^2} \text{ Se acelera}$$

¿Cómo le harías para que la desaceleración fuera menor?

R= Permitiría que la red se elongue más.

Problema 2.

A person driving her car at 45 km/h approaches an intersection just as the traffic light turns yellow. She knows that the yellow light lasts only 2.0 s before turning to red, and she is 28 m away from the near side of the intersection (Fig. 51). Should she try to stop, or should she speed up to cross the intersection before the light turns red? The intersection is 15 m wide. Her car's maximum deceleration is -5.8 m/s^2 , whereas it can accelerate from 45 km/h to 65 km/h in 6.0 s. Ignore the length of her car and her reaction time.




Figura 2: This is a captiion

Caso 1

$$V_o = 12,5 \frac{\text{m}}{\text{s}} \quad x = 28\text{m} \quad t = 2\text{s}$$

$$a = -5,8 \frac{\text{m}^2}{\text{s}} \quad V_f = 0 \frac{\text{m}}{\text{s}}$$

$$V^2 = V_o^2 + 2a (x - x_o)$$

$$2a (X - X_o) = V^2 - V_o^2$$

$$X - X_o = \frac{V - V_o}{2a} = \frac{-\left(12,5 \frac{\text{m}}{\text{s}}\right)}{-2\left(5,8 \frac{\text{m}^2}{\text{s}}\right)} = 13,46 \text{ m}$$

Caso 2

Aceleración del auto a ver si alcanza la intersección.

$$a = \frac{V - V_o}{t}$$

$$\frac{(18.05 \frac{m}{s} - 12.5 \frac{m}{s})}{6s} 0.925 \frac{m}{s}$$

$$X = X_o + V_o t + \frac{1}{2} a t^2$$

$$(12.5 \frac{m}{s})(2s) + \frac{1}{2}(0.925)(2s^2) = 26.85 m$$

Le conviene detenerse, dado que sólo podrá aumentar 26.85m y el trayecto es de 4.

Problema 3.

(II) Extreme-sports enthusiasts have been known to jump off the top of El Capitan, a sheer granite cliff of height 910 m in Yosemite National Park. Assume a jumper runs horizontally off the top of El Capitan with speed 5.0 m/s and enjoys a freefall until she is 150 m above the valley floor, at which time she opens her parachute (Fig. 41). (a) How long is the jumper in freefall? Ignore air resistance. (b) It is important to be as far away from the cliff as possible before opening the parachute. How far from the cliff is this jumper when she opens her chute?




Figura 3: This is a caption

En dirección x

$$V_{xo} = \frac{x}{t}$$

$$x = V_{xo} t$$

En dirección y

$$Y = y_o + V_{oy} t - \frac{1}{2} g t^2$$

$$150 \text{ m} = 910 \text{ m} - \frac{1}{2} g t^2$$

$$\frac{1}{2} g t^2 = 910 \text{ m} - 150 \text{ m} = 760 \text{ m}$$

$$t^2 = \frac{2(760 \text{ m})}{9.81 \frac{\text{m}}{\text{s}^2}}$$

$$t = \sqrt{\frac{2(760 \text{ m})}{9.81 \frac{\text{m}}{\text{s}^2}}} = 12.44 \text{ s tiempo para la caída libre}$$

$$X = V_{xo} t = \left(5 \frac{\text{m}}{\text{s}}\right) (12.44 \text{ s}) = 62.2 \text{ m se encontraba lejos}$$

Link del video

<https://youtu.be/M9DKgmNF0vY>