

Amelioration of cadmium chloride and arsenic trioxide induced genotoxicity in *Oreochromis mossambicus* using ocimum and ascorbic acid.

Dr. Papri saha¹ and Post doctoral article²

¹Paprisaha76@gmail.com

²Affiliation not available

September 3, 2019

Abstract

The genotoxic effects of cadmium chloride and arsenic trioxide was assessed singly and conjointly in *Oreochromis mossambicus* (tilapia) with end points such as chromosome aberration, red blood cell anomaly , anomalous nuclei determination and sperm head anomaly detection.

The primary objective were to examine If cadmium chloride and arsenic trioxide as well as ascorbic acid (AA) and ocimum, had, any genotoxic effect of their own on non target aquatic organisms and secondly if ascorbic acid and ocimum had any ameliorating effect on cadmium chloride and arsenic trioxide induced genotoxicity in the fish, tilapia.

key words

Chromosome aberration(CA), genotoxicity, mutation, cell anomaly, ocimum, AA, cadmium chloride, arsenic trioxide.

Introduction (summary)

Heavy metals are known to cause abundance of effects on all living organisms including fish (olsson,1996, Sorenson,1991).

The overall objective of study is to ascertain the level and quantum of genotoxic effects of cadmium chloride and arsenic trioxide as compared to DW treated controls and also to find out if certain antioxidants like AA and ocimum can nullify the oxidative metabolism of these chemical by producing electrophilic metabolites.

Methodology

Somatic chromosomes were prepared as the conventional colchicine citrate flame drying technique(khud-abukhsh ,1979) for both control and treated series.

For micronuclei testing , the procedure of Schmidt(1976) was followed.

For sperm head anomaly, procedure of Manna and Biswas was followed, further followed by microscopic analysis.

Experimental set up was provided by taking

18 to 22gms fish, collected from local pond and then acclimatized in concrete vats maintained under similar conditions.

Conclusion

Maximum amelioration have been found at 48hr interval for cadmiumchloride and arsenic trioxide induced genotoxicity, when treated conjointly with either ocimum or with ascorbic acid, in tilapia.

However both AA and ocimum showed certain amount of genotoxicity when treated individually in the fish specimens.

The dose at which maximum amelioration could be found was noted, which was 0.02 percent for AA and 0.5 percent for ocimum.

The antioxidative property is mainly due to phenolic acid, flavonoids and anthocyanins present in vitamin c and basil (ocimum).

Data format

Table

Hosted file 1566749313445503611934.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Hosted file 1566749424738-1994465419.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Hosted file 1566749494967-1298195396.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Hosted file 1566822712854-1994465419.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Hosted file 1566822591764503611934.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Hosted file 1566822359978-492617179.jpg available at <https://authorea.com/users/272838/articles/387371-amelioration-of-cadmium-chloride-and-arsenic-trioxide-induced-genotoxicity-in-oreochromis-mossambicus-using-ocimum-and-ascorbic-acid>

Abbreviations

CA..chromosome aberration

MN.micronuclei

AA..ascorbic acid

Dw..distilled water

Reference

Chandra, P. and Khuda_BUKhsh, A.R. (2004). Genotoxic effects of cadmium chloride and azadirachtin treated singly and in combination in fish. *Ecotox Environ Safety*. 58: 194_201.

Davison, k., Mann k.k and Miller, W.H.JR. (2002). Arsenic trioxide mechanism of action. *Seminar Hematol*. 39: 3_7.

Al.Sabti and Metcalfe C.D. (1995). Fish micronuclei for assessing genotoxicity in water. *Mutat Red* . 343: 121_135.

De Flora, S . and Ramel, C. (1988). Mechanism of inhibitor of mutagenesis and carcinogenesis classification and overview. *Mutat Red*. 202: 285_306.

Chandra, P. and khudabukhsh, A. R. (2001). An assay of genotoxicity produced by cadmium chloride in fish (*Oreochromis mossambicus*) and efficacy of vitamin c in its alteration. *Process xth All India Cong cytol Genet, kalyani, West Bengal oct29_31 , 1999*, In: *pers cytol Genet*, (Eds G.k Manna and S.C Roy). Springer. 10: 583_590.

Authors contribution and information

I, Dr. Papri Saha, am a former ICAR fellow at University of Kalyani and at present, Lecturer in BRSN college, and also engaged in post doctoral work.

Acknowledgement

I would like to thank the library staffs and my fellow mates, and all my well wishers for helping me with this article .

I would also like to thank brsnc, for cooperating with me to carry out my post doctoral articles, apart from my regular teaching schedule in the college.