

Table 2: Univariate analysis of prognostic variables affecting survival in pB-LBL

Variables (n)	5yr-EFS (95%CI)	P-value	5yr-OS (95%CI)	P-value
Age				
<10 years (14)	76.6±12.1%	0.731	92.9±6.9%	0.982
≥10 years (7)	85.7±13.2%		85.7±13.2%	
Sex				
Male (15)	71.1±12.4%	0.184	86.7±8.8%	0.229
Female (6)	100%		100%	
Symptom duration				
<3months (15)	93.3±6.4%	0.030	93.3±6.4%	0.124
≥3months (6)	44.4±22.2%		83.3±15.2%	
Primary tumor site				
Osteolytic lesion (4)	37.5±28.6%	0.161	75±21.7%	0.08
Soft tissue mass (11)	81.8±11.6%		90.9±8.7%	
Other sites [¥] (6)	100%		100%	
Baseline LDH				
Normal (11)	90.9±8.7%	0.132	100%	0.037
Elevated (8)	60±18.2%		75±15.3%	
Stage				
Limited stage (9)	77.8±13.9%	0.763	88.9±10.5%	0.735
Advanced Stage (12)	80.2±12.8%		91.7±8.0%	
Bone marrow involvement				
No (12)	74.1±12.1%	0.378	83.3±15.2%	0.693
Yes [€] (9)	83.3±10.8%		85.7±13.2%	
Interim response				
Complete response (15)	79.4±10.6%	0.823	86.7±8.8%	0.893
Partial response (6)	75.0±21.7%		75.0±21.7%	
[¥] Other sites included: 2-Testicular mass,1-Right atrial mass, 1-Mediastinal mass, 1 Subdural mass, 1-Lymph node enlargement. [€] Bone marrow involvement: 1-Morphologic, 4-MDD±PET, 4-PET only				

