

The normalized Laplacians spectrum and characteristic parameters of a class of irregular networks

Jia-Bao Liu¹, Qian Zheng^{1,*}, Jinde Cao²

¹School of Mathematics and Physics, Anhui Jianzhu University, Hefei 230601, PR China

²School of Mathematics, Southeast University, Nanjing 210096, China

Abstract. The normalized Laplacian plays an indispensable role in exploring the structural properties of irregular graphs. Let $L_n^{8,4}$ represents a linear octagonal-quadrilateral network. Then, by identifying the opposite lateral edges of $L_n^{8,4}$, we get the corresponding Möbius graph $MQ_n(8,4)$. In this paper, starting from the decomposition theorem of polynomials, we infer that the normalized Laplacian spectrum of $MQ_n(8,4)$ can be determined by the eigenvalues of two symmetric quasi-triangular matrices \mathcal{L}_A and \mathcal{L}_S of order $4n$. Next, owing to the relationship between the two matrix roots and the coefficients mentioned above, we derive the explicit expressions of the degree-Kirchhoff indices and the complexity of $MQ_n(8,4)$.

Keywords: Irregular network; Möbius graphs; Normalized Laplacian; Degree-Kirchhoff index; Complexity

1. Introduction

It is well established that networks can be represented by graphs. The graphs we consider in this paper are simple, undirected and connected. Let's first recall some definitions commonly used in graph theory. Suppose G represents a simple undirected graph with $|V_G| = n$ and $|E_G| = m$. For more notation, one can be referred to [1].

Note that $D(G) = \text{diag}\{d_1, d_2, \dots, d_n\}$ represents a degree matrix, where d_p is the degree of v_p . $A(G)$ is the adjacency matrix of G . The Laplacian matrix of G is $L(G) = D(G) - A(G)$. The (p,q) -entry of the normalized Laplacian matrix is given by

$$(\mathcal{L}(G))_{pq} = \begin{cases} 1, & p = q; \\ -\frac{1}{\sqrt{d_p d_q}}, & p \neq q \text{ and } v_p \sim v_q; \\ 0, & \text{otherwise.} \end{cases} \quad (1.1)$$

As a matter of fact, there are many parameters that can be used to describe the structure and properties of molecular graphs in graph networks. One of the parameters based on resistance distance is defined as Wiener index [2, 3], which is

$$W(G) = \sum_{i < j} d_{ij},$$

where $d_{ij} = d_G(v_i, v_j)$ represents the length of the shortest path between two vertices v_i and v_j in G . Wiener index is widely used in chemical and mathematical research. For details, see [4–7].

The parameter of resistance distance was first proposed by Klein and Randić [8] in 1993. It means that if every edge of a graph G is regarded as a unit resistance, then the distance between any two vertices i and j in G is called resistance distance, which is denoted as r_{ij} . Similar to Wiener index, we give the

E-mail address: liujiabaoad@163.com, zhengqian19960202@163.com, jdcao@seu.edu.cn.

* Corresponding author.

expression of the Kirchhoff index [9, 10] according to the resistance distance, namely

$$Kf(G) = \sum_{i < j} r_{ij} = n \sum_{i=2}^n \frac{1}{\mu_i}.$$

In 2007, Chen and Zhang [11] proposed that the eigenvalues and eigenvectors of normalized Laplacian spectrum can be used to describe the resistance distance, and an observation that prompted a new characteristic parameter, called the degree-Kirchhoff index, which is a kind of structural descriptor [11]. However, it is very difficult to calculate the degree-Kirchhoff index from the complexity division of graphs, so it is very important to find the explicit expression of degree-Kirchhoff index. In recent years, many scholars have devoted themselves to the study of degree-Kirchhoff index of various graphs. Huang and Li et al. [12, 13] proved the degree-Kirchhoff index of linear hexagonal chains and linear polyomino chains, successively. H. Bian et al. [14] determined the normalized Laplacians and degree-Kirchhoff index of cylinder phenylene chain. Zhao and Liu et al. [15] described the normalized Laplacian and degree-Kirchhoff index of linear octagonal-quadrilateral networks. For more excellent results, please refer to [16–21]. After learning the excellent works of scholars, in this paper, we use the correlation properties of normalized Laplace matrix to calculate the degree-Kirchhoff index and the complexity of Möbius graph of linear octagonal-quadrilateral networks.

The investigation of complex graph and irregular network has gone through a spectacular development in the past decades. Especially in organic chemistry, more and more attention has been paid to the application of polyomino in polycyclic aromatic compounds. Many scholars are interested in the study of linear octagonal networks and related molecular graphs. As we all know, linear octagonal network is an octagonal system without branch compression. It is constructed by regularly inserting some new points on the straight line of the linear polyomino network. The research on the structure and properties of this kind of natural graph network lays a solid foundation for the advancement of theoretical chemistry, as well as for the development of applied mathematics.

Let $L_n^{8,4}$ be the linear octagonal-quadrilateral networks and octagons and quadrilaterals are connected by a common edge, which depicted in Figure 1. Then the corresponding Möbius graph $MQ_3(8, 4)$ of octagonal-quadrilateral networks is obtained by the reverse identification of the opposite edge by $L_n^{8,4}$, see Figure 2. Obviously, we can obtained that $|V_{MQ_n}(8, 4)| = 8n$, $|E_{MQ_n}(8, 4)| = 10n$.




Figure 1: Linear octagonal-quadrilateral networks.

The rest of the paper will be divided into the following several sections: In Section 2, we put forward some basic notation and related lemmas. In Section 3, we determine the normalized Laplacian spectrum of $MQ_n(8, 4)$. In Section 4, we committed to give the Kemeny's constant, the degree-Kirchhoff index and the complexity of $MQ_n(8, 4)$.




Figure 2: Graph $MQ_3(8,4)$.

2. Preliminary

In this section, we introduce some common symbols and related calculation methods [1], which are applied to the rest of the article.

The characteristic polynomial of matrix R of order n is defined as $P_R(x) = \det(xI - R)$. It's not difficult to find that π is an automorphism of G , we can write the product of disjoint 1-cycles and transposition, namely

$$\pi = (\bar{1})(\bar{2}) \cdots (\bar{m})(1, 1')(2, 2') \cdots (k, k').$$

Then one has $|V(G)| = m + 2k$, let $v_0 = \{\bar{1}, \bar{2}, \cdots \bar{m}\}$, $v_1 = \{1, 2 \cdots k\}$, $v_2 = \{1', 2' \cdots k'\}$. Thus the Laplacians matrix can be expressed in the form of block matrix, that is

$$\mathcal{L}(G) = \begin{pmatrix} \mathcal{L}_{V_0 V_0} & \mathcal{L}_{V_0 V_1} & \mathcal{L}_{V_0 V_2} \\ \mathcal{L}_{V_1 V_0} & \mathcal{L}_{V_1 V_1} & \mathcal{L}_{V_1 V_2} \\ \mathcal{L}_{V_2 V_0} & \mathcal{L}_{V_2 V_1} & \mathcal{L}_{V_2 V_2} \end{pmatrix},$$

where

$$\mathcal{L}_{V_0 V_1} = \mathcal{L}_{V_0 V_2}, \quad \mathcal{L}_{V_1 V_2} = \mathcal{L}_{V_2 V_1}, \quad \text{and} \quad \mathcal{L}_{V_1 V_1} = \mathcal{L}_{V_2 V_2}.$$

Let

$$P = \begin{pmatrix} I_m & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}}I_k & \frac{1}{\sqrt{2}}I_k \\ 0 & \frac{1}{\sqrt{2}}I_k & -\frac{1}{\sqrt{2}}I_k \end{pmatrix},$$

then

$$P' \mathcal{L}(G) P = \begin{pmatrix} \mathcal{L}_A(G) & 0 \\ 0 & \mathcal{L}_S(G) \end{pmatrix},$$

noted that P' is the transposition of P , where

$$\mathcal{L}_A = \begin{pmatrix} \mathcal{L}_{V_0 V_0} & \sqrt{2}\mathcal{L}_{V_0 V_1} \\ \sqrt{2}\mathcal{L}_{V_1 V_0} & \mathcal{L}_{V_1 V_1} + \mathcal{L}_{V_1 V_2} \end{pmatrix}, \quad \mathcal{L}_S = \mathcal{L}_{V_1 V_1} - \mathcal{L}_{V_1 V_2}.$$

Lemma 2.1. [12] Let $\mathcal{L}(L_n)(G)$, $\mathcal{L}_A(G)$, $\mathcal{L}_S(G)$ are determined as above, then

$$P_{\mathcal{L}(L_n)}(G) = P_{\mathcal{L}_A}(G) P_{\mathcal{L}_S}(G).$$

Lemma 2.2. Let G is a graph with $|V_G| = n$ and $|E_G| = m$, and $0 = \mu_1 < \mu_2 \leq \cdots \leq \mu_n (n \geq 2)$ are the eigenvalues of $\mathcal{L}(G)$. Then we can quickly confirm that the following formulas holds.

(a) [22] The Kemeny's constant of G can be denoted

$$K_c(G) = \sum_{i=2}^n \frac{1}{\mu_i}.$$

(b) [11] The degree-Kirchhoff index of G is defined as

$$Kf^*(G) = 2m \sum_{k=2}^n \frac{1}{\mu_k}.$$

(c) [1] *The number of spanning trees of G can also be called the complexity of G . Then the complexity of G is*

$$\prod_{i=1}^n d_i \sum_{k=2}^n \lambda_k = 2m\tau(G).$$

3. The normalized Laplacian spectrum of $MQ_n(8, 4)$

In this section, we focus on obtain the normalized Laplacian spectrum of $MQ_n(8, 4)$ by lemma 2.1.

Given an $n \times n$ matrix T , and put deleting the p_1th , p_2th , \dots p_kth rows and columns of T are expressed as $T[\{p_1, p_2, \dots, p_k\}]$. With a suitable labeling, the vertices of $MQ_n(8, 4)$ show in Figure 2. Apparently, $\pi = (1, 1')(2, 2') \dots (4n, (4n)')$ is an automorphism of $MQ_n(8, 4)$. Then $v_0 = \emptyset$, $v_1 = \{1, 2, 3, \dots, 4n\}$ and $v_2 = \{1', 2', 3', \dots, (4n)'\}$. Besides, we express $\mathcal{L}_A(MQ_n(8, 4))$ and $\mathcal{L}_S(MQ_n(8, 4))$ as \mathcal{L}_A and \mathcal{L}_S . Then one can get

$$\mathcal{L}_A = \mathcal{L}_{V_1 V_1} + \mathcal{L}_{V_1 V_2}, \quad \mathcal{L}_S = L_{V_1 V_1} - \mathcal{L}_{V_1 V_2}.$$

In views of Equation (1.1), we have

and

$$\mathcal{L}_{V_1 V_2} = \begin{pmatrix} \frac{-1}{3} & & & & & & & \frac{-1}{3} \\ & 0 & & & & & & \\ & & 0 & & & & & \\ & & & \frac{-1}{3} & & & & \\ & & & & 0 & & & \\ & & & & & 0 & & \\ & & & & & & \ddots & \\ & & & & & & & \frac{-1}{3} \\ & & & & & & & 0 \\ & & & & & & & & 0 \\ \frac{-1}{3} & & & & & & & & & \frac{-1}{3} \end{pmatrix}_{(4n) \times (4n)}.$$

Hence,

$$\mathcal{L}_A = \begin{pmatrix} \frac{2}{3} & \frac{-1}{\sqrt{6}} & & & & & & & \frac{-1}{3} \\ \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & & & & & \\ & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} & & & & & \\ & & \frac{-1}{\sqrt{6}} & \frac{2}{3} & \frac{-1}{3} & & & & \\ & & & \frac{2}{3} & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & & & \\ & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & & \\ & & & & \ddots & \ddots & \ddots & & \\ & & & & \frac{-1}{3} & \frac{2}{3} & \frac{-1}{\sqrt{6}} & & \\ & & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & \\ & & & & & \ddots & \ddots & \ddots & \\ & & & & & \frac{2}{3} & \frac{-1}{\sqrt{6}} & \frac{-1}{2} & \\ & & & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & \\ & & & & & & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} \\ & & & & & & & \frac{-1}{\sqrt{6}} & \frac{2}{3} \\ & & & & & & & & \frac{-1}{3} \end{pmatrix}_{(4n) \times (4n)},$$

and

$$\mathcal{L}_S = \begin{pmatrix} \frac{4}{3} & \frac{-1}{\sqrt{6}} & & & & & & & \frac{-1}{3} \\ \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & & & & & \\ & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} & & & & & \\ & & \frac{-1}{\sqrt{6}} & \frac{4}{3} & \frac{-1}{3} & & & & \\ & & & \frac{4}{3} & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & & & \\ & & & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & \\ & & & & \ddots & \ddots & \ddots & & \\ & & & & \frac{-1}{3} & \frac{4}{3} & \frac{-1}{\sqrt{6}} & & \\ & & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & & \\ & & & & & \ddots & \ddots & \ddots & \\ & & & & & \frac{4}{3} & \frac{-1}{\sqrt{6}} & \frac{-1}{2} & \\ & & & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & \\ & & & & & & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} \\ & & & & & & & \frac{-1}{\sqrt{6}} & \frac{4}{3} \\ & & & & & & & & \frac{-1}{3} \end{pmatrix}_{(4n) \times (4n)}.$$

Assuming that $0 = \eta_1 < \eta_2 \leq \eta_3 \leq \dots \leq \eta_{4n}$ are the roots of $P_{\mathcal{L}_A}(x) = 0$, and $0 < \varphi_1 \leq \varphi_2 \leq \varphi_3 \leq \dots \leq \varphi_{4n}$ are the roots of $P_{\mathcal{L}_S}(x) = 0$, respectively. Then according to lemma 2.1 we can get the spectrum of $MQ_n(8, 4)$ is just $\eta_1, \eta_2, \dots, \eta_{4n}, \varphi_1, \varphi_2, \dots, \varphi_{4n}$, and it is directly to check that $\eta_1 = 0, \eta_p > 0 (p = 2, 3, \dots, 4n)$, and $\varphi_q > 0 (q = 1, 2, \dots, 4n)$.

Nextly, we will committed to calculate some main results of $MQ_n(8, 4)$ related to the normalized Laplacian spectrum.

4. The degree-Kirchhoff index and the complexity of $MQ_n(8, 4)$

In this section, we first introduce some theorems, which are obtained by describing the eigenvalues and eigenvectors of normalized Laplacian matrix. Then obtained the Kemeny's constant, the degree-Kirchhoff index and the complexity of $MQ_n(8, 4)$ based on these theorems.

Theorem 4.1.

$$\sum_{p=2}^{4n} \frac{1}{\eta_p} = \frac{200n^2 - 11}{60}.$$

Proof. Let

$$\begin{aligned} P_{\mathcal{L}_S}(x) &= \det(xI - \mathcal{L}_A) = x^{4n} + a_1 x^{4n-1} + \dots + a_{4n-1} x + a_{4n} \\ &= x(x^{4n-1} + a_1 x^{4n-2} + \dots + a_{4n-2} x + a_{4n-1}), \quad a_{4n-1} \neq 0. \end{aligned}$$

$$\mathcal{L}_A^2 = \begin{pmatrix} 1 & & & & & & & & & & \\ \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{6}} & & & & & & & & & \\ & \frac{-1}{3} & \frac{-1}{3} & & & & & & & & \\ & & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{6}} & & & & & & & \\ & & & 1 & \frac{-1}{2} & & & & & & \\ & & & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} & & & & & \\ & & & & \frac{-1}{\sqrt{6}} & \frac{-1}{3} & \frac{-1}{3} & & & & \\ & & & & & \frac{-1}{2} & \ddots & & & & \\ & & & & & & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} & & \\ & & & & & & & \frac{-1}{\sqrt{6}} & \frac{-1}{3} & \frac{-1}{3} & \\ & & & & & & & & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & \frac{-1}{3} \\ & & & & & & & & & \frac{-1}{\sqrt{6}} & 1 \end{pmatrix}_{(4n) \times (4n)},$$

and

$$\mathcal{L}_A^3 = \begin{pmatrix} \frac{2}{3} & & & & & & & & & & \\ \frac{-1}{3} & \frac{-1}{3} & & & & & & & & & \\ & \frac{-1}{\sqrt{6}} & \frac{-1}{\sqrt{6}} & & & & & & & & \\ & & 1 & \frac{-1}{2} & & & & & & & \\ & & \frac{-1}{2} & 1 & \frac{-1}{\sqrt{6}} & & & & & & \\ & & & \frac{-1}{\sqrt{6}} & \frac{-1}{3} & \frac{-1}{3} & & & & & \\ & & & & \frac{-1}{3} & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & & & & \\ & & & & & \frac{-1}{\sqrt{6}} & \ddots & & & & \\ & & & & & & \frac{-1}{\sqrt{6}} & \frac{2}{3} & \frac{-1}{3} & & \\ & & & & & & & \frac{-1}{3} & \frac{-1}{3} & \frac{-1}{\sqrt{6}} & \\ & & & & & & & & \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} \\ & & & & & & & & & \frac{-1}{2} & 1 \end{pmatrix}_{(4n) \times (4n)}.$$

In this way, we can get four facts.

Fact 1. For $1 \leq p \leq 4n$,

$$r_p^0 = \begin{cases} (p+1) \left(\frac{1}{36} \right)^{\frac{p}{4}}, & \text{if } p \equiv 0(\text{mod}4); \\ \frac{1}{3}(p+1) \left(\frac{1}{36} \right)^{\frac{p-1}{4}}, & \text{if } p \equiv 1(\text{mod}4); \\ \frac{1}{6}(p+1) \left(\frac{1}{36} \right)^{\frac{p-2}{4}}, & \text{if } p \equiv 2(\text{mod}4); \\ \frac{1}{12}(p+1) \left(\frac{1}{36} \right)^{\frac{p-3}{4}}, & \text{if } p \equiv 3(\text{mod}4). \end{cases}$$

Fact 2. For $1 \leq p \leq 4n$,

$$r_p^1 = \begin{cases} (p+1) \left(\frac{1}{36} \right)^{\frac{p}{4}}, & \text{if } p \equiv 0(\text{mod}4); \\ \frac{1}{2}(p+1) \left(\frac{1}{36} \right)^{\frac{p-1}{4}}, & \text{if } p \equiv 1(\text{mod}4); \\ \frac{1}{4}(p+1) \left(\frac{1}{36} \right)^{\frac{p-2}{4}}, & \text{if } p \equiv 2(\text{mod}4); \\ \frac{1}{12}(p+1) \left(\frac{1}{36} \right)^{\frac{p-3}{4}}, & \text{if } p \equiv 3(\text{mod}4). \end{cases}$$

Fact 3. For $1 \leq p \leq 4n$,

$$r_p^2 = \frac{1}{2}r_{p-2}^0 - \frac{1}{9}r_{p-3}^1.$$

Fact 4. For $1 \leq p \leq 4n$,

$$r_p^3 = \frac{2}{3}r_{p-1}^0 - \frac{1}{9}r_{p-2}^1.$$

Proof of Fact 1. Take $r_p^0 = \det R_p^0$, $r_p^1 = \det R_p^1$, $r_p^2 = \det R_p^2$ and $r_p^3 = \det R_p^3$. By a straightforward calculation, one can get the following values, see Table 1.

Table 1: Initial value.

r_p^0	Value	r_p^0	Value	r_p^0	Value	r_p^0	Value
r_1^0	$\frac{2}{3}$	r_2^0	$\frac{1}{2}$	r_3^0	$\frac{1}{3}$	r_4^0	$\frac{5}{36}$
r_5^0	$\frac{3}{54}$	r_6^0	$\frac{7}{216}$	r_7^0	$\frac{1}{54}$	r_8^0	$\frac{1}{144}$

For $4 \leq p \leq 4n - 1$, we can get expending $\det R_p^0$ with respect to its last row yields

$$r_p^0 = \begin{cases} \frac{2}{3}r_{p-1}^0 - \frac{1}{6}r_{p-2}^0, & \text{if } p \equiv 0(\text{mod}4); \\ \frac{2}{3}r_{p-1}^0 - \frac{1}{9}r_{p-2}^0, & \text{if } p \equiv 1(\text{mod}4); \\ r_{p-1}^0 - \frac{1}{6}r_{p-2}^0, & \text{if } p \equiv 2(\text{mod}4); \\ r_{p-1}^0 - \frac{1}{4}r_{p-2}^0, & \text{if } p \equiv 3(\text{mod}4). \end{cases}$$

For $1 \leq p \leq n - 1$, let $a_p = r_{4p}^0$; $0 \leq p \leq n - 1$, $b_p = r_{4p+1}^0$, $c_p = r_{4p+2}^0$, $d_p = r_{4p+3}^0$. Then we can get $a_1 = \frac{5}{36}$, $b_0 = \frac{2}{3}$, $c_0 = \frac{1}{2}$, $d_0 = \frac{5}{36}$, $b_1 = \frac{3}{54}$, $c_1 = \frac{7}{216}$, $d_1 = \frac{1}{54}$, and for $p \geq 2$, we have

$$\begin{cases} a_p = \frac{2}{3}d_{p-1} - \frac{1}{6}c_{p-1}; \\ b_p = \frac{2}{3}a_p - \frac{1}{9}d_{p-1}; \\ c_p = b_p - \frac{1}{6}a_p; \\ d_p = c_p - \frac{1}{4}b_p. \end{cases}$$

Then it's no difficult to obtain that

$$\begin{cases} a_p = 18c_p - 24d_p; \\ b_p = 4c_p - 4d_p; \\ c_p = \frac{1}{18}c_{p-1} - \frac{1}{1296}c_{p-2}; \\ d_p = \frac{1}{18}d_{p-1} - \frac{1}{1296}d_{p-2}. \end{cases} \quad (4.3)$$

According to the equation of d_p in (4.3) is $x^2 - \frac{1}{18}x + \frac{1}{1296} = 0$, and its two roots are $\frac{1}{36}$ and $\frac{1}{36}$. Therefore, $d_p = (x_p + y)(\frac{1}{36})^p$ is the general solution. Then we can get $x = \frac{1}{3}$ and $y = \frac{1}{3}$.

Thus, we can obtained $d_p = \frac{1}{3}(p+1)(\frac{1}{36})^p$ ($p \geq 1$). Similarly, we have $c_p = (\frac{2p}{3} + \frac{1}{2})(\frac{1}{36})^p$ ($p \geq 1$); $a_p = (4p+1)(\frac{1}{36})^p$ ($p \geq 1$) and $b_p = \frac{2}{3}(2p+1)(\frac{1}{36})^p$ ($p \geq 1$).

The result as desired. ■

By the similar consideration, Facts 2 is available. Then based on the conclusion of Facts 1 and 2, we quickly get Facts 3 and 4.

Now, we will further calculate $(-1)^{4n-1}a_{4n-1}$ and $(-1)^{4n-2}a_{4n-2}$ in equation (4.2). For the sake of discussion, it is assumed that $r_0 = 1$.

Claim 1. $(-1)^{4n-1}a_{4n-1} = 40n^2\left(\frac{1}{36}\right)^n$.

Proof of Claim 1. Since the $(-1)^{4n-1}a_{4n-1}$ is the total of all the principal minors of order $4n-1$ of \mathcal{L}_A , we have

$$\begin{aligned} (-1)^{4n-1}a_{4n-1} &= \sum_{p=1}^{4n} \det \mathcal{L}_A[p] = \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p] + \sum_{p=1, p \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p] \\ &\quad + \sum_{p=2, p \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p] + \sum_{p=3, p \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p]. \end{aligned}$$

where

$$\begin{aligned} \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p] &= \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{4n-p}^0 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^0); \\ \sum_{p=1, p \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p] &= \sum_{p=1, p \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{4n-p}^1 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^1); \\ \sum_{p=2, p \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p] &= \sum_{p=2, p \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{4n-p}^2 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^2); \\ \sum_{p=3, p \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p] &= \sum_{p=3, p \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{4n-p}^3 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^3). \end{aligned}$$

By Fact 1-2, we have

$$\begin{aligned} \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{4n-p}^0 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^0) &= \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} \left[\frac{p}{12} \left(\frac{1}{36}\right)^{\frac{p-4}{4}} (4n-p+1) \frac{1}{12} \left(\frac{1}{36}\right)^{\frac{4n-p}{4}} \right. \\ &\quad \left. - \frac{1}{9} (p-1) \frac{1}{4} \left(\frac{1}{36}\right)^{\frac{p-4}{4}} \frac{1}{12} (4n-p) \left(\frac{1}{36}\right)^{\frac{4n-p-4}{4}} \right] \\ &= \sum_{p=4, p \equiv 0 \pmod{4}}^{4n} 12n \left(\frac{1}{36}\right)^n \\ &= 12n^2 \left(\frac{1}{36}\right)^n. \end{aligned}$$

Similarly, by Fact 1-4 we can get

$$\begin{aligned} \sum_{p=1, p \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{4n-p}^1 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^1) &= 12n^2 \left(\frac{1}{36}\right)^n; \\ \sum_{p=2, p \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{4n-p}^2 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^2) &= 8n^2 \left(\frac{1}{36}\right)^n; \\ \sum_{p=3, p \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{4n-p}^3 - \frac{1}{9} r_{p-2}^1 r_{4n-p-1}^3) &= 8n^2 \left(\frac{1}{36}\right)^n. \end{aligned}$$

Hence, according to the above results, we have

$$(-1)^{4n-1}a_{4n-1} = \sum_{p=1}^{4n} \det \mathcal{L}_A[p] = 40n^2 \left(\frac{1}{36}\right)^n.$$

The proof of Claim 1 completed. ■

Claim 2. $(-1)^{4n-2}a_{4n-2} = \frac{2}{3}(200n^4 - 11n^2)\left(\frac{1}{36}\right)^n$.

Proof of Claim 2. It's not hard to see that $(-1)^{4n-2}a_{4n-2}$ is the total of the those principal minors \mathcal{L}_A , which have $(4n-2)$ rows and columns. Thus we have

$$(-1)^{4n-2}a_{4n-2} = \sum_{1 \leq i < j \leq 4n} \det \mathcal{L}_A[p, q]. \quad (4.4)$$

By equation (4.4), it can be seen that the change of i and j values will lead to different $\det \mathcal{L}_A[p, q]$ results. Therefore, we will choose different p and q to list the following equations.

$$\begin{aligned} \sum_{1 \leq p < q \leq 4n} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 3 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\ &+ \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q]. \end{aligned}$$

where by Fact 1-4, we can compute the following results

Case 1.

$$\begin{aligned} \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{q-p-1}^0 r_{4n-q}^0 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^0 r_{4n-q-1}^0) \\ &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 0 \pmod{4}}^{4n} 9(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\ &= 12(n^4 - n^2) \left(\frac{1}{36}\right)^n. \end{aligned}$$

Case 2.

$$\begin{aligned}
\sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{q-p-1}^0 r_{4n-q}^1 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^0 r_{4n-q-1}^1) \\
&= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 1 \pmod{4}}^{4n-3} 9(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{3}{2} (8n^4 - 12n^3 + n^2 + 3n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 3.

$$\begin{aligned}
\sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{q-p-1}^0 r_{4n-q}^2 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^0 r_{4n-q-1}^2) \\
&= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 2 \pmod{4}}^{4n-2} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 + 8n^3 + 4n^2 + 4n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 4.

$$\begin{aligned}
\sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{q-p-1}^0 r_{4n-q}^3 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^0 r_{4n-q-1}^3) \\
&= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 3 \pmod{4}}^{4n-1} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 - 4n^3 + n^2 - 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 5.

$$\begin{aligned}
\sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{q-p-1}^1 r_{4n-q}^0 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^1 r_{4n-q-1}^0) \\
&= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 0 \pmod{4}}^{4n} 9(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{3}{2} (8n^4 + 12n^3 + n^2 - 3n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 6.

$$\begin{aligned}
\sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{q-p-1}^1 r_{4n-q}^1 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^1 r_{4n-q-1}^1) \\
&= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 1 \pmod{4}}^{4n-3} 9(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= 12(n^4 - n^2) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 7.

$$\begin{aligned}
\sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{q-p-1}^1 r_{4n-q}^2 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^1 r_{4n-q-1}^2) \\
&= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 2 \pmod{4}}^{4n-2} 9(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 + 4n^3 + n^2 + 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 8.

$$\begin{aligned}
\sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{q-p-1}^1 r_{4n-q}^3 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^1 r_{4n-q-1}^3) \\
&= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 3 \pmod{4}}^{4n-1} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 - 8n^3 + 4n^2 - 4n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 9.

$$\begin{aligned}
\sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{q-p-1}^2 r_{4n-q}^0 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^2 r_{4n-q-1}^0) \\
&= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 0 \pmod{4}}^{4n} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 + 8n^3 + 4n^2 + 4n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 10.

$$\begin{aligned}
\sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{q-p-1}^2 r_{4n-q}^1 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^2 r_{4n-q-1}^1) \\
&= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 - 4n^3 + n^2 - 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 11.

$$\begin{aligned}
\sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{q-p-1}^2 r_{4n-q}^2 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^2 r_{4n-q-1}^2) \\
&= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 2 \pmod{4}}^{4n-2} 4(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{16}{3} (n^4 - n^2) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 12.

$$\begin{aligned}
\sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{q-p-1}^2 r_{4n-q}^3 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^2 r_{4n-q-1}^3) \\
&= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 3 \pmod{4}}^{4n-1} 4(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{2}{3} (8n^4 + 4n^3 + n^2 + 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 13.

$$\begin{aligned}
\sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 0 \pmod{4}}^{4n} (r_{p-1}^0 r_{q-p-1}^3 r_{4n-q}^0 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^3 r_{4n-q-1}^0) \\
&= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 0 \pmod{4}}^{4n} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 + 4n^3 + n^2 + 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 14.

$$\begin{aligned}
\sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 1 \pmod{4}}^{4n-3} (r_{p-1}^0 r_{q-p-1}^3 r_{4n-q}^1 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^3 r_{4n-q-1}^1) \\
&= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 1 \pmod{4}}^{4n-3} 6(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= (8n^4 - 8n^3 + 4n^2 - 4n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 15.

$$\begin{aligned}
\sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} (r_{p-1}^0 r_{q-p-1}^3 r_{4n-q}^2 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^3 r_{4n-q-1}^2) \\
&= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} 4(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{2}{3} (8n^4 - 4n^3 + n^2 - 5n) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Case 16.

$$\begin{aligned}
\sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] &= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 3 \pmod{4}}^{4n-1} (r_{p-1}^0 r_{q-p-1}^3 r_{4n-q}^3 - \frac{1}{9} r_{p-2}^1 r_{q-p-1}^3 r_{4n-q-1}^3) \\
&= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} 4(q-p)(4n-q+p) \left(\frac{1}{36}\right)^n \\
&= \frac{16}{3} (n^4 - n^2) \left(\frac{1}{36}\right)^n.
\end{aligned}$$

Then, according to the value of p , the above sixteen cases can be divided into the following four categories.

$$\begin{aligned}
F_0 &= \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\
&+ \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 0 \pmod{4}}^{4n-4} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\
&= \frac{1}{2}(80n^4 - 28n^3 - 11n^2 + 7n) \left(\frac{1}{36} \right)^n.
\end{aligned}$$

$$\begin{aligned}
F_1 &= \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\
&+ \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 1 \pmod{4}}^{4n-3} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\
&= \frac{1}{2}(80n^4 + 28n^3 - 11n^2 - 7n) \left(\frac{1}{36} \right)^n.
\end{aligned}$$

$$\begin{aligned}
F_2 &= \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\
&+ \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 2 \pmod{4}}^{4n-2} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\
&= \frac{1}{3}(80n^4 + 20n^3 + n^2 + 7n) \left(\frac{1}{36} \right)^n.
\end{aligned}$$

$$\begin{aligned}
F_3 &= \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 0 \pmod{4}}^{4n} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 3 \pmod{4}}^{4n-2} \sum_{q \equiv 1 \pmod{4}}^{4n-3} \det \mathcal{L}_A[p, q] \\
&+ \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 2 \pmod{4}}^{4n-2} \det \mathcal{L}_A[p, q] + \sum_{p \equiv 3 \pmod{4}}^{4n-1} \sum_{q \equiv 3 \pmod{4}}^{4n-1} \det \mathcal{L}_A[p, q] \\
&= \frac{1}{3}(80n^4 - 20n^3 + 10n^2 - 7n) \left(\frac{1}{36} \right)^n.
\end{aligned}$$

Substituting F_0 , F_1 , F_2 , and F_3 to Equation(4.4), one has

$$(-1)^{4n-2} a_{4n-2} = F_0 + F_1 + F_2 + F_3 = \frac{2}{3}(200n^4 - 11n^2) \left(\frac{1}{36} \right)^n.$$

The result as desired. ■

So substituting the results of Claim 1 and 2 into Equation (4.2) yields

$$\begin{aligned}
\sum_{p=2}^{4n} \frac{1}{\eta_p} &= \frac{(-1)^{4n-2} a_{4n-2}}{(-1)^{4n-1} a_{4n-1}} \\
&= \frac{\frac{2}{3}(200n^4 - 11n^2) \left(\frac{1}{36} \right)^n}{40n^2 \left(\frac{1}{36} \right)^n} \\
&= \frac{200n^2 - 11}{60}.
\end{aligned}$$

Fact 7. For $1 \leq q \leq 4n$,

$$s_q^2 = \frac{7}{6}s_{q-2}^0 - \frac{1}{9}s_{q-3}^1.$$

Fact 8. For $1 \leq q \leq 4n$,

$$s_q^3 = \frac{4}{3}s_{q-1}^0 - \frac{1}{9}s_{q-2}^1.$$

Proof of Fact 5. Take $s_q^0 = \det S_q^0$, $s_q^1 = \det S_q^1$, $s_q^2 = \det S_q^2$ and $s_q^3 = \det S_q^3$. By direct calculation, it's no difficult to get the following values, see Table 2.

Table 2: Initial value.							
s_q^0	Value	s_q^0	Value	s_q^0	Value	s_q^0	Value
s_1^0	$\frac{4}{3}$	s_2^0	$\frac{7}{6}$	s_3^0	$\frac{5}{6}$	s_4^0	$\frac{33}{36}$
s_5^0	$\frac{61}{54}$	s_6^0	$\frac{211}{216}$	s_7^0	$\frac{25}{36}$	s_8^0	$\frac{989}{1296}$

For $4 \leq q \leq 4n$, we have the expansion-formula of the $\det S_q^0$ with respect to its last row yields

$$s_q^0 = \begin{cases} \frac{4}{3}t_{q-1}^0 - \frac{1}{6}s_{q-2}^0, & \text{if } q \equiv 0(\text{mod}4); \\ \frac{4}{3}t_{q-1}^0 - \frac{1}{9}s_{q-2}^0, & \text{if } q \equiv 1(\text{mod}4); \\ s_{q-1}^0 - \frac{1}{6}s_{q-2}^0, & \text{if } q \equiv 2(\text{mod}4); \\ s_{q-1}^0 - \frac{1}{4}s_{q-2}^0, & \text{if } q \equiv 3(\text{mod}4). \end{cases}$$

For $1 \leq q \leq n$, let $A_q = s_{4q}$; $0 \leq q \leq n-1$, $B_q = s_{4q+1}$, $C_q = s_{4q+2}$, $D_q = s_{4q+3}$. Then we may obtain that

$$\begin{cases} A_q = \frac{4}{3}D_{q-1} - \frac{1}{6}C_{q-1}; \\ B_q = \frac{4}{3}A_q - \frac{1}{9}D_{q-1}; \\ C_q = B_q - \frac{1}{6}A_q; \\ D_q = C_q - \frac{1}{4}B_q. \end{cases} \quad (4.6)$$

From the first three equations in (4.6), one can get $A_q = \frac{12}{13}C_q + \frac{1}{78}C_{q-1}$. Next, substituting A_q to the third equation, one has $B_q = \frac{15}{13}C_q + \frac{1}{468}C_{q-1}$. Then substituting B_q to the fourth equation, we have $D_q = \frac{37}{52}C_q - \frac{1}{1872}C_{q-1}$. Finally, Substituting A_q and d_q to the first equation, one has $c_q - 30c_{q-1} + c_{q-2} = 0$. Thus

$$C_q = k_1 \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^q + k_2 \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^q.$$

In views of $C_0 = \frac{7}{6}$, $C_1 = \frac{211}{216}$, we have

$$\begin{cases} k_1 + k_2 = \frac{7}{6}; \\ k_1 \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right) + k_2 \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right) = \frac{211}{216}, \end{cases}$$

and

$$\begin{cases} k_1 = \left(\frac{7}{12} + \frac{53\sqrt{14}}{336} \right); \\ k_2 = \left(\frac{7}{12} - \frac{53\sqrt{14}}{336} \right). \end{cases}$$

Thus it is routine to deduce that

$$\begin{cases} A_q = \left(\frac{4}{8} + \frac{9\sqrt{14}}{56}\right)\left(\frac{5}{12} + \frac{\sqrt{14}}{9}\right)^q + \left(\frac{4}{8} - \frac{9\sqrt{14}}{56}\right)\left(\frac{5}{12} - \frac{\sqrt{14}}{9}\right)^q, & \text{if } q \equiv 0(\text{mod } 4); \\ B_q = \left(\frac{2}{3} + \frac{31\sqrt{14}}{168}\right)\left(\frac{5}{12} + \frac{\sqrt{14}}{9}\right)^q + \left(\frac{2}{3} - \frac{31\sqrt{14}}{168}\right)\left(\frac{5}{12} - \frac{\sqrt{14}}{9}\right)^q, & \text{if } q \equiv 1(\text{mod } 4); \\ C_q = \left(\frac{7}{12} + \frac{53\sqrt{14}}{336}\right)\left(\frac{5}{12} + \frac{\sqrt{14}}{9}\right)^q + \left(\frac{7}{12} - \frac{53\sqrt{14}}{336}\right)\left(\frac{5}{12} - \frac{\sqrt{14}}{9}\right)^q, & \text{if } q \equiv 1(\text{mod } 4); \\ D_q = \left(\frac{5}{12} + \frac{25\sqrt{14}}{224}\right)\left(\frac{5}{12} + \frac{\sqrt{14}}{9}\right)^q + \left(\frac{5}{12} - \frac{25\sqrt{14}}{224}\right)\left(\frac{5}{12} - \frac{\sqrt{14}}{9}\right)^q, & \text{if } q \equiv 3(\text{mod } 4). \end{cases}$$

The result as desired. ■

In the same way, we can quickly prove the result of fact 6.

Then we expand $\det s_q^2$ and s_q^3 according to the properties of determinant, and we can get facts 7 and

8.

Now by exploiting the property of determinant, we can get

$$\begin{aligned} \det \mathcal{L}_S &= \begin{vmatrix} \frac{4}{3} & \frac{-1}{\sqrt{6}} & 0 & \cdots & 0 & 0 & \frac{1}{3} \\ \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & \cdots & 0 & 0 & 0 \\ 0 & \frac{-1}{2} & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 & \frac{-1}{\sqrt{6}} \\ \frac{1}{3} & 0 & 0 & \cdots & 0 & \frac{-1}{\sqrt{6}} & \frac{4}{3} \end{vmatrix}_{4n} \\ &= \begin{vmatrix} \frac{4}{3} & \frac{-1}{\sqrt{6}} & 0 & \cdots & 0 & 0 & 0 \\ \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & \cdots & 0 & 0 & 0 \\ 0 & \frac{-1}{2} & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 & \frac{-1}{\sqrt{6}} \\ \frac{1}{3} & 0 & 0 & \cdots & 0 & \frac{-1}{\sqrt{6}} & \frac{4}{3} \end{vmatrix}_{4n} + \begin{vmatrix} \frac{4}{3} & \frac{-1}{\sqrt{6}} & 0 & \cdots & 0 & 0 & \frac{1}{3} \\ \frac{-1}{\sqrt{6}} & 1 & \frac{-1}{2} & \cdots & 0 & 0 & 0 \\ 0 & \frac{-1}{2} & 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 & \frac{-1}{2} & 0 \\ 0 & 0 & 0 & \cdots & 0 & 1 & 0 \\ \frac{1}{3} & 0 & 0 & \cdots & 0 & \frac{-1}{\sqrt{6}} & 0 \end{vmatrix}_{4n} \\ &= s_{4n}^0 - \frac{1}{9}s_{4n-2}^1 + 2\left(\frac{1}{36}\right)^n \end{aligned}$$

Together with Facts 1 and 2, we can obtain one interesting Claim.

Claim 3. $\det \mathcal{L}_S = \left(\frac{5}{12} + \frac{\sqrt{14}}{9}\right)^n + \left(\frac{5}{12} - \frac{\sqrt{14}}{9}\right)^n + 2\left(\frac{1}{36}\right)^n$.

Then we're going to concentrate on calculating $(-1)^{4n-1}b_{4n-1}$.

Claim 4. $(-1)^{4n-1}b_{4n-1} = \frac{41n\sqrt{14}}{28} \left[\frac{\left((15+4\sqrt{14})^n - (15-4\sqrt{14})^n\right)}{\left((15+4\sqrt{14})^n + (15-4\sqrt{14})^n\right) + 2} \right]$.

Proof. Since the $(-1)^{4n-1}b_{4n-1}$ is the total of all the principal minors of order $4n-1$ of \mathcal{L}_S , we have

$$\begin{aligned} (-1)^{4n-1}b_{4n-1} &= \sum_{q=1}^{4n} \det \mathcal{L}_S[q] = \sum_{q=4, q \equiv 0(\text{mod } 4)}^{4n} \det \mathcal{L}_S[q] + \sum_{q=1, q \equiv 1(\text{mod } 4)}^{4n-3} \det \mathcal{L}_S[q] \\ &\quad + \sum_{q=2, q \equiv 2(\text{mod } 4)}^{4n-2} \det \mathcal{L}_S[q] + \sum_{q=3, q \equiv 3(\text{mod } 4)}^{4n-1} \det \mathcal{L}_S[q]. \end{aligned}$$

where

$$\det \mathcal{L}_S[q] = \begin{cases} s_{q-1}^0 s_{4n-q}^0 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^0, & \text{if } q \equiv 0(\text{mod}4); \\ s_{q-1}^0 s_{4n-q}^1 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^1, & \text{if } q \equiv 1(\text{mod}4); \\ s_{q-1}^0 s_{4n-q}^2 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^2, & \text{if } q \equiv 2(\text{mod}4); \\ s_{q-1}^0 s_{4n-q}^3 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^3, & \text{if } q \equiv 3(\text{mod}4). \end{cases} \quad (4.7)$$

For $q \equiv 0(\text{mod}4)$ and $4 \leq q \leq 4n-4$, in views of (4.7) and Fact 5-8, one gets

$$\begin{aligned} \det \mathcal{L}_S[q] &= s_{q-1}^0 s_{4n-q}^0 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^0 \\ &= \left[\left(\frac{5}{12} + \frac{25\sqrt{14}}{224} \right) \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^{\frac{j-4}{4}} + \left(\frac{5}{12} - \frac{25\sqrt{14}}{224} \right) \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^{\frac{j-4}{4}} \right] \\ &\quad \times \left[\left(\frac{4}{8} + \frac{25\sqrt{14}}{56} \right) \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^{\frac{4n-j}{4}} + \left(\frac{4}{8} - \frac{25\sqrt{14}}{56} \right) \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^{\frac{4n-j}{4}} \right] \\ &\quad - \frac{1}{9} \left[\left(\frac{3}{8} + \frac{23\sqrt{14}}{224} \right) \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^{\frac{j-4}{4}} + \left(\frac{3}{8} - \frac{23\sqrt{14}}{224} \right) \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^{\frac{j-4}{4}} \right] \\ &\quad \times \left[\left(\frac{5}{12} + \frac{25\sqrt{14}}{224} \right) \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^{\frac{4n-j-4}{4}} + \left(\frac{5}{12} - \frac{25\sqrt{14}}{224} \right) \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^{\frac{4n-j-4}{4}} \right] \\ &= \frac{15n\sqrt{14}}{56} \left[\left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n - \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n \right]. \end{aligned}$$

Similarly, for $q \equiv 1(\text{mod}4)$ and $1 \leq q \leq 4n-3$, we have

$$\begin{aligned} \sum_{q=1, q \equiv 1(\text{mod}4)}^{4n-3} \det \mathcal{L}_S[q] &= s_{q-1}^0 s_{4n-q}^1 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^1 \\ &= \frac{15n\sqrt{14}}{56} \left[\left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n - \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n \right]. \end{aligned}$$

For $q \equiv 2(\text{mod}4)$ and $2 \leq q \leq 4n-2$, we have

$$\begin{aligned} \sum_{q=2, q \equiv 2(\text{mod}4)}^{4n-2} \det \mathcal{L}_S[q] &= s_{q-1}^0 s_{4n-q}^2 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^2 \\ &= \frac{13n\sqrt{14}}{28} \left[\left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n - \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n \right]. \end{aligned}$$

For $q \equiv 3(\text{mod}4)$ and $3 \leq q \leq 4n-1$, we have

$$\begin{aligned} \sum_{q=3, q \equiv 3(\text{mod}4)}^{4n-1} \det \mathcal{L}_S[q] &= s_{q-1}^0 s_{4n-q}^3 - \frac{1}{9} s_{q-2}^1 s_{4n-q-1}^3 \\ &= \frac{13n\sqrt{14}}{28} \left[\left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n - \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n \right]. \end{aligned}$$

Thus, one has the following equation

$$(-1)^{4n-1} b_{4n-1} = \frac{41n\sqrt{14}}{28} \left[\left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n - \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n \right].$$

Therefore, substituting the results of claim 3 and 4 into (4.5) can be obtained

$$\sum_{q=1}^{4n} \frac{1}{\varphi_q} = \frac{41n\sqrt{14}}{28} \left[\frac{((15+4\sqrt{14})^n - (15-4\sqrt{14})^n)}{((15+4\sqrt{14})^n + (15-4\sqrt{14})^n) + 2} \right].$$

The result as desired. \blacksquare

Note that $|E_{MQ_n(8,4)}| = 10n$. Take the results of Theorems 4.1 and 4.2 to Lemma 2.2 (a) and (b), we can immediately get the following two Theorems.

Theorem 4.3. *Let $MQ_n(8,4)$ be a möbius graph with n octagonal and n quadrilateral. Then*

$$\begin{aligned} Kc(MQ_n(8,4)) &= \sum_{p=2}^{4n} \frac{1}{\eta_p} + \sum_{q=1}^{4n} \frac{1}{\varphi_q} \\ &= \frac{200n^2 - 11}{60} + \frac{41n\sqrt{14}}{28} \left[\frac{((15 + 4\sqrt{14})^n - (15 - 4\sqrt{14})^n)}{((15 + 4\sqrt{14})^n + (15 - 4\sqrt{14})^n) + 2} \right]. \end{aligned}$$

Theorem 4.4. *Let $MQ_n(8,4)$ be a möbius graph with n octagonal and n quadrilateral. Then*

$$\begin{aligned} Kf^*(MQ_n(8,4)) &= 20n \left(\sum_{p=2}^{4n} \frac{1}{\eta_p} + \sum_{q=1}^{4n} \frac{1}{\varphi_q} \right) \\ &= 20n \left(\frac{200n^2 - 11}{60} + \frac{41n\sqrt{14}}{28} \left[\frac{((15 + 4\sqrt{14})^n - (15 - 4\sqrt{14})^n)}{((15 + 4\sqrt{14})^n + (15 - 4\sqrt{14})^n) + 2} \right] \right) \\ &= \frac{200n^3 - 11n}{3} + 20n\varrho(n), \end{aligned}$$

$$\text{where } \varrho(n) = \frac{41n\sqrt{14}}{28} \left[\frac{((15 + 4\sqrt{14})^n - (15 - 4\sqrt{14})^n)}{((15 + 4\sqrt{14})^n + (15 - 4\sqrt{14})^n) + 2} \right].$$

The degree-Kirchhoff indices of Möbius graph of linear octagonal-quadrilateral networks, see Table 3.

Table 3: The degree-Kirchhoff indices of $MQ_1(8,4), MQ_2(8,4), \dots, MQ_{16}(8,4)$.

G	$Kf^*(G)$	G	$Kf^*(G)$	G	$Kf^*(G)$	G	$Kf^*(G)$
$MQ_1(8,4)$	165.50	$MQ_5(8,4)$	11054.43	$MQ_9(8,4)$	57442.75	$MQ_{13}(8,4)$	164937.53
$MQ_2(8,4)$	963.33	$MQ_6(8,4)$	18322.78	$MQ_{10}(8,4)$	77587.71	$MQ_{14}(8,4)$	204359.11
$MQ_3(8,4)$	2775.12	$MQ_7(8,4)$	28210.28	$MQ_{11}(8,4)$	101951.83	$MQ_{15}(8,4)$	249599.85
$MQ_4(8,4)$	6005.23	$MQ_8(8,4)$	41116.93	$MQ_{12}(8,4)$	130935.10	$MQ_{16}(8,4)$	301059.74

Finally, we will concentrate on calculate the complexity of $MQ_n(8,4)$.

Theorem 4.5. *Let $MQ_n(8,4)$ denote a Möbius graph of linear octagonal-quadrilateral networks of length $n \geq 2$. Then*

$$\tau(MQ_n(8,4)) = 4n \left((15 + 4\sqrt{14})^n + (15 - \sqrt{14})^n + 2 \right).$$

Proof.

By Claim 1, one can get

$$\prod_{p=2}^{4n} \eta_p = (-1)^{4n-1} a_{4n-1} = 40n^2 \left(\frac{1}{36} \right)^n.$$

Similarly, according to Claim 3, we have

$$\prod_{q=1}^{4n} \varphi_q = \det \mathcal{L}_S = \left(\frac{5}{12} + \frac{\sqrt{14}}{9} \right)^n + \left(\frac{5}{12} - \frac{\sqrt{14}}{9} \right)^n + 2 \left(\frac{1}{36} \right)^n.$$

Note that $\prod_{i=1}^{8n} d_i(MQ_n) = 2^{4n}3^{4n}$ and $|E_{MQ_n}(8, 4)| = 10n$. By lemma 2.4, one gets

$$\tau(Q_n(8, 4)) = \frac{1}{10n} \prod_{p=2}^{4n} \eta_p \cdot \prod_{q=1}^{4n} \varphi_q = 4n \left((15 + 4\sqrt{14})^n + (15 - \sqrt{14})^n + 2 \right).$$

This completes the proof. ■

Thus we can get the complexity of $MQ_n(8, 4)$, which are listed in Table 4.

Table 4: The complexity of Q_1, Q_2, \dots, Q_{10} .

G	$\tau(G)$	G	$\tau(G)$
Q_1	128	Q_5	17379554400
Q_2	7200	Q_7	607607778176
Q_3	322944	Q_8	20809093939328
Q_4	12902464	Q_9	701525710449792
Q_5	483303040	Q_{10}	23358178980900000

Funding

This work was supported in part by Anhui Provincial Natural Science Foundation under Grant 2008085J01, and by National Natural Science Foundation of China Grant 11601006.

References

- [1] F.R.K. Chung, Spectral graph theory, American Mathematical Society Providence, RI, 1997.
- [2] H. Wiener, Structural determination of paraffin boiling points, Journal of the American Chemical Society 69 (1947) 17-20.
- [3] A. Dobrynin, Branchings in trees and the calculation of the Wiener index of a tree, Match Communications in Mathematical and in Computer Chemistry 41 (2000) 119-134.
- [4] A. Dobrynin, R. Entringer, I. Gutman, Wiener index of trees: Theory and Applications, Acta Applicandae Mathematicae 66 (2002) 211-249.
- [5] A. Dobrynin, I. Gutman, S. Klavžar, and P. Žigert, Wiener index of hexagonal systems, Acta Applicandae Mathematicae 72 (2002) 94-247.
- [6] I. Gutman, S. C. Li, W. Wei, Cacti with n vertices and t cycles having extremal Wiener index, Discrete Applied Mathematics 232 (2017) 189-200.
- [7] I. Gutman, S. Klavžar, B. Mohar, MATCH Commun. Math. Comput. Chem 35 (1997) 1-259.
- [8] D.J. Klein, M. Randić, Resistance distances, Journal of Mathematical Chemistry 12 (1993) 81-95.
- [9] D.J. Klein, Resistance-distance sum rules, Croatica Chemica Acta 75 (2002) 633-649.
- [10] D.J. Klein, O. Ivanciuc, Graph cyclicity, excess conductance, and resistance deficit, Journal of Mathematical Chemistry 30 (2001) 271-287.
- [11] H. Y. Chen, F. J. Zhang, Resistance distance and the normalized Laplacian spectrum, Discrete Appl. Math. 155 (2007) 654-661.
- [12] J. Huang, S. C. Li, L. Q. Sun, The normalized Laplacians, degree-Kirchhoff index and the spanning trees of linear hexagonal chains, Discrete Appl. Math. 207 (2016) 67-79.
- [13] J. Huang, S. C. Li, X. C. Li, The normalized Laplacian, degree-Kirchhoff index and spanning trees of the linear polyomino chains, Appl. Math. Comput. 289 (2016) 324-334.
- [14] X.L. Ma, H. Hong, The normalized Laplacians, degree-Kirchhoff index and the spanning trees of cylinder phenylene chain, Polycyclic Aromatic Compounds. (2019)DOI: 10.1080/10406638.2019.1665553.
- [15] J.B. Liu, J. Zhao, Z. Zhu, On the number of spanning trees and normalized Laplacian of linear octagonal-quadrilateral networks, International Journal of Quantum Chemistry 119 (17) (2019) e25971.

- [16] Y. J. Peng, S. C. Li, On the Kirchhoff index and the number of spanning trees of linear phenylenes, *MATCH Commun. Math. Comput. Chem.* 77 (2017) 765-780.
- [17] X.Y. Geng, P. W, On the Kirchhoff indices and the number of spanning trees of Möbius phenylenes chain and cylinder phenylenes chain, *Polycyclic Aromatic Compounds* (2019)DOI: 10.1080/10406638.2019.1693405.
- [18] J.B. Liu, Z.Y. Shi, Y.H. Pan, Computing the Laplacian spectrum of linear octagonal-quadrilateral networks and its applications, *Polycyclic Aromatic Compounds*. (2020)DOI: 10.1080/10406638.2020.1748666.
- [19] J. Huang, S.C. Li, X. Li, The normalized Laplacian, degree-Kirchhoff index and spanning trees of the linear polyomino chains, *Applied Mathematics and Computation* 289 (2016) 324-334.
- [20] Z.X. Zhu, J.B. Liu, The normalized Laplacian, degree-Kirchhoff index and the spanning tree numbers of generalized phenylenes, *Discrete Applied Mathematics* 254 (2019) 256-267.
- [21] S. Li, W. Sun, S. Wang, Multiplicative degree-Kirchhoff index and number of spanning trees of a zigzag polyhex nanotube TUHC[2n,2], *International Journal of Quantum Chemistry* 119 (17) (2019) e25969.
- [22] S. Butler, Algebraic aspects of the normalized Laplacian, in: A. Beveridge, J. Griggs, L. Hogben, G. Musiker, P. Tetali (eds.), *Recent Trends in Combinatorics*, vol. to appear of The IMA Volumes in Mathematics and its Applications, IMA, 2016.