

Table 1: Descriptive results related with urological complaints, and percentages of these findings according to groups.

Parameter	Group 1 (%) (+)	Group 1 (%) (-)	Group 2 (%) (+)	Group 2 (%) (-)
Dysuria	8.64	91.36	40	60
Testicular Swelling	1.23	98.77	10	90
History of BCG Vaccine	8.64	91.36	60	40
History of Epididymo-orchitis	7.4	92.6	30	70
History of Urological Surgery	2.46	97.54	0	100
Comorbid Diseases	18.51	81.49	20	80

Table 2: Statistical comparison of groups in terms of inflammatory markers and disease duration.

Parameter	Group 1 (Median-IQR)	Group 2 (Median-IQR)	p value
Neutrophil Count ($10^3/\text{mm}^3$)	3.74 (2.36)	4.19 (2.86)	0.395
Lymphocyt Count ($10^3/\text{mm}^3$)	1.62 (0.84)	1.99 (1.33)	0.069
Neutrophil-Lymphocyte Ratio	2.16 (2.32)	2.21 (1.11)	0.909
C-Reactive Protein (mg/L)	4.6 (27.95)	0 (58.75)	0.478
D-Dimer ($\mu\text{g/L}$)	266 (428)	186.5 (202.25)	0.193
Duration of COVID-19 (Day)	12 (9)	10 (7.75)	0.465

Table 3: Statistical comparison of groups according to the lymphocytopenia levels.

Parameter	Lymphopenia ($<1.5 \times 10^3/\text{mm}^3$) (n=37)	Lymphopenia ($>1.5 \times 10^3/\text{mm}^3$) (n=54)	p value
Age (Years)*	40 (23.5)	29 (27.5)	0.038 ^x
Duration of COVID-19 (Day)*	11 (10.5)	12 (6.25)	0.704 ^x
Dysuri	5.4%	14.81%	0.757 ^y
Testicular Pain	10.81%	12.96%	0.159 ^y
Testicular Swelling	5.4%	1.85%	0.786 ^y
History of BCG Vaccine	51.35%	51.85%	0.795 ^y
History of Epididymo-orchitis	13.51%	7.4%	0.338 ^y
History of Urological Surgery	2.7%	1.85%	0.786 ^y
Comorbid Diseases	18.91%	18.51%	0.962 ^y

*Median (IQR), ^x: Mann-Whitney U test, ^y: Chi-square test.