

**Tab.2** Characteristic parameters of the spatial distribution of cypress and topography in each runoff plot

N O	Topographic relief (m)	Surface roughness	Surface cutting depth (cm)	Runoff path density (m/100m <sup>2</sup> )	L(r) index	Contagion index	Stand density of cypress (ind/100m <sup>2</sup> )	Surface runoff coefficient
1	1.71	1.22	9.54	12.44	-0.10	0.41	14	0.62
2	1.54	1.23	8.71	9.64	0.22	0.48	22	0.51
3	1.44	1.35	10.32	11.22	0.08	0.42	34	0.56
4	1.36	1.21	3.49	10.08	-0.05	0.53	22	0.39
5	1.71	1.31	6.94	9.62	0.22	0.45	36	0.60
6	1.39	1.61	10.76	9.92	-0.13	0.54	66	0.32
7	1.61	1.72	6.95	8.02	0.16	0.43	36	0.35
8	1.31	1.62	5.94	6.96	0.26	0.64	54	0.21
9	1.66	1.45	8.73	6.52	0.03	0.47	70	0.33
10	1.31	1.73	6.32	10.08	0.19	0.60	68	0.20
11	1.51	1.65	7.43	7.9	0.03	0.40	34	0.33
12	1.55	1.46	4.76	9.3	0.08	0.47	10	0.41