

	<b>SARS-CoV-2 only (n=249)</b>	<b>SARS-CoV-2 coinfection with another respiratory virus (n=33)</b>	<b>P- value</b>
<b>Age (Median, IQR)</b>	59 (49-73)	58 (45-69)	0.26
<b>Male gender</b>	175 (70%)	24 (73%)	0.84
<b>Fever</b>	202 (81%)	21 (64%)	0.037
<b>Dyspnea</b>	137 (55%)	16 (48%)	0.57
<b>Expectoration</b>	10 (5%)	2 (10%)	0.6
<b>Cough</b>	184 (74%)	22 (67%)	0.4
<b>Myalgia</b>	74 (30%)	10 (30%)	1
<b>Headache</b>	36 (14%)	10 (30%)	0.04
<b>Symptoms duration, median (IQR)</b>	4 [3-7]	3 [2-7]	0.26
<b>ICU admission, day 1</b>	64 (26%)	0 (0%)	<0.001
<b>ICU admission, day 14</b>	31 (15%)	3 (9%)	0.27
<b>Hospital mortality</b>	49 (20%)	5 (15%)	0.48

**Table 1. Main clinical characteristics of patients presenting a SARS-CoV-2 infection with or without any other respiratory viruses coinfection.**