

**TABLE 2 Characteristics of study participants who experienced  $\geq 10\%$  decline in LVEF and one who experience a decline in LEVF to a final value of  $< 50\%$**

	<b><math>\geq 10\%</math> decline</b>					<b>LVEF <math>&lt; 50\%</math></b>
	Patient 1	Patient 2	Patient 3	Patient 4	Patient 5	Patients 6
Age (yrs.)	15	3	9	4	2	3
Sex	Male	Male	Male	Female	Male	Male
BSA (m <sup>2</sup> )	1.2	0.53	1.1	0.59	0.49	0.56
Nutrition status	Underweight	Normal	Normal	Normal	Normal	Underweight
Diagnosis	BL	BL	BL	BL	Rhabdo-myosarcoma	Metastatic Wilms Tumour
<b>(At baseline)</b>						
BP (mmHg)	110/70	91/52	119/69	100/60	90/71	107/78
Heart Rate(b/min)	102	110	74	125	153	131
Temperature	36.5°C	36.5°C	36.8°C	36.5°C	35.8°C	36.8°C
Haemoglobin(g/dl)	8.6	10.9	12.3	9.4	5.8	6.5
<b>Cardiac Echo (LVEF-%)</b>						
• LVEF0	66	71	70	72	72	55
• LVEF1	54	61	60	61	67	46
• LVEF2	52	61	65	66	60	56
• LVEF3	56	54	-	61	-	61

BSA- Body Surface Area, LVEF 0- Left Ventricular Ejection Fraction at baseline, LVEF 1 – Left Ventricular Ejection Fraction at one month, LVEF 2- Left Ventricular Ejection Fraction at six months, LVEF 3- Left Ventricular Ejection Fraction at 12 months