

Table 4. Appropriateness of the decisions made by midwives, characteristics of pregnant women, and work shift during which the episode was assessed.

Inappropriateness (n=95)			Appropriateness (n=2,647)		p
Quantitative	Mean	SD	Mean	SD	
Age	29.3	5.6	30.7	5.6	0.019
Weight gain (kg)	11.4	5.0	12.3	5.0	0.072
BMI	29.0	4.7	29.0	4.7	0.993
Presence of pain in the last episode	3.7	2.6	3.6	2.6	0.598
Qualitative	n	%	n	%	
Clinically relevant personal history					
Absence	87	91.6	2450	96.6	0.875
Presence	8	8.4	197	96.1	
Clinically relevant surgical history					
Absence	58	61.1	1652	96.6	0.872
Presence	37	38.9	995	96.4	
Parity					
Nulliparous	36	37.6	1130	96.9	0.192
Multiparous	48	50.4	1098	95.8	
Nationality					
Spanish	14	14.7	1172	98.8	0.464
non-Spanish	6	6.3	801	99.3	
Smoker					
No	80	84.2	2144	96.4	0.307
Yes	8	8.4	333	97.7	
Funding					
Private	0	0.0	34	100.0	1
Social Security	20	21.1	1975	99.0	
High-risk at admission					
No	62	65.8	2032	97.0	0.014
Yes	33	34.2	615	94.9	
Day of the week					
Monday	18	18.9	433	96.0	0.222

Tuesday	11	2. 8	378	97.2	
Wednesday	14	3. 7	365	96.3	
Thursday	14	3. 5	382	96.5	
Friday	13	3. 4	373	96.6	
Saturday	19	5. 2	345	94.8	
Sunday	6	1. 6	371	98.4	
Shift of the widwife					
Morning	22	2. 0	1062	98.0	
Afternoon	29	4. 2	664	95.8	0.00
Night	44	4. 6	921	95.4	4
