

Table 4 In vitro susceptibility testing of the <i>Myroides odoratimimus</i> isolates		
Antibiotics	(n), MIC value (µg/mL)	Interpretation (S, I, R)
Aztreonam	(n=25/25), ≥ 64 µg/mL	100% R
Ceftazidime	(n=25/25), ≥ 64 µg/mL	100% R
Ciprofloxacin	(n=25/25), ≥ 4 µg/mL	100% R
Cefepime	(n=24/25), ≥ 32 µg/mL, (n=1/25) 16 µg/mL	96.0% R, 4.0% I
Gentamicin	(n=25/25), ≥ 16 µg/mL	100% R
Imipenem	(n=25/25), ≥ 16 µg/mL	100% R
Levofloxacin	(n=25/25), ≥ 8 µg/mL	100% R
Meropenem	(n=25/25), ≥ 16 µg/mL	100% R
Netilmicin	(n=25/25), ≥ 32 µg/mL	100% R
Piperacillin	(n=25/25), ≥ 128 µg/mL	100% R
Trimethoprim / Sulfamethoxazole	(n=25/25), ≥ 320 µg/mL	100% R
Tetracycline	(n=25/25), ≥ 16 µg/mL	100% R
Tobramycin	(n=25/25), ≥ 16 µg/mL	100% R
Piperacillin / Tazobactam	(n=24/25), ≥ 128 µg/mL, (n=1/25) ≤4 µg/mL	96.0% R, 4.0% S
Amikacin	(n=25/25), ≥ 64 µg/mL	100% R
Abbreviations: MIC, Minimum inhibitory concentration; R, resistant; I, intermediate; S, susceptible		