

<b>Diagnostic test</b>	<b>Female</b>	<b>Male</b>	<b>&lt; 55 years</b>	<b>55 years</b>	<b>rUTI</b>	<b>Total Cohort</b>
<b>Cystoscopy (700)</b>	<b>n=427</b>	<b>n=273</b>	<b>n= 241</b>	<b>n=459</b>	<b>n=331</b>	<b>n=700</b>
Normal	418 (98%)	258 (94.5%)	238 (98.8%)	438 (95.4%)	316 (95.5%)	676 (96.6%)
Abnormal	9 (2%)	15 (5.5%)	3 (1.2%)	21 (4.6%)	15 (4.5%)	24 (3.4%)
<b>Imaging (621)</b>	<b>n=380</b>	<b>n=241</b>	<b>n=212</b>	<b>n=409</b>	<b>n=298</b>	<b>n=621</b>
Normal	350 (92%)	226 (93.7%)	197 (93%)	379 (92.7%)	273 (91.6%)	576 (92.8%)
Abnormal	30 (8%)	15 (6.3%)	15 (7%)	30 (7.3%)	25 (8.4%)	45 (7.2%)
<b>Cytology (521)</b>	<b>n=310</b>	<b>n=211</b>	<b>n=168</b>	<b>n=353</b>	<b>n=243</b>	<b>n=521</b>
Normal	308 (99.4%)	210 (99.6%)	168 (100%)	350 (99%)	240 (98.8%)	518 (99.5%)
Abnormal	2 (0.6%)	1 (0.4%)	0 (0%)	3 (1%)	3 (1.2%)	3 (0.5%)

**Table 3. Investigational findings stratified by gender and age and those meeting guideline criteria for rUTI**