

Table-2 The differences between clinicopathological characteristics, perioperative factors and groups

	Group 1	Group 2	p Value
ASA 1-2	29 (80.6%)	13 (46.4%)	0.004
ASA 3	7 (19.4%)	15 (53.6%)	
Comorbidity			0.025
Present	17 (47.2%)	7 (25%)	
Absent	17 /47.2%)	21 (75%)	
Preoperative Hospitalization period, day			
Comorbidity present	3.9±2.6	5.9±2.5	0.047
Comorbidity absent	6.1±4.0	9.8±4.2	0.008
Surgery Duration (min) (mean±SD)	125.4±29.5	148.9±39.9	0.018
Hematocrit (n, %)			0.014
<30	16 (44.4%)	21 (75%)	
Atropine (n, %)			>0.999
Administered	34 (94.4%)	26 (92.9%)	
Ephedrine (n, %)			<0.001
Administered	33 (91.7%)	15 (53.6%)	
Midazolam (n, %)			0.223
Administered	19 (52.8%)	19 (67.9%)	
Crystalloid (ml) (Mean±SD)	2222±806	1821±495	0.087
Colloid Replacement (n, %)			
Administered	21 (58.3%)	18 (64.3%)	0.628
Not administered	15 (41.7%)	10 (35.7%)	
Hemorrhage (ml) (Mean±SD)	633.3±305.9	764.3±322.3	0.046
Erythrocyte Replacement (n, %)			
Administered	21 (58.3%)	19 (67.9%)	0.892