

	Total (n=239)	Pandemy (-) (n=108, 45.2%)	Pandemy (+) (n=131, 54.8%)	P Value
Age (mean, std)	58.8 (12.1)	59.8 (12.3)	58.0 (11.9)	0.26
Sex (male)	175 (72.9%)	77 (70.6%)	98 (74.8%)	0.47
Diabetes Mellitus	71 (29.6%)	34 (31.2%)	37 (28.2%)	0.61
Hypertension	104 (43.3%)	52 (47.7%)	52 (39.7%)	0.21
Renal insufficiency	13 (5.4%)	5 (4.6%)	8 (6.1%)	0.60
CAD	63 (26.3%)	32 (29.4%)	31 (23.7%)	0.31
CHF	10 (4.2%)	7 (6.4%)	3 (2.3%)	0.11
Atrial Fibrillation	9 (3.8%)	8 (7.3%)	1 (0.8%)	0.01
COPD	9 (3.8%)	3 (2.8%)	6 (4.6%)	0.45
SVE	11 (4.6%)	8 (7.3%)	3 (2.3%)	0.06
Malignancy	10 (4.2%)	7 (6.4%)	3 (2.4%)	0.12
STEMI (n, %)	91 (38.1%)	32 (29.6%)	59 (45.0%)	0.01
NSTEMI	91(38.1)	45(41.7%)	46(35.1%)	0.29
USAP	57(23.8%)	31(28.7%)	26(19.8%)	0.11
LVEF (%) <i>Median (Q1-Q3)</i>	50 (40-55)	55 (40-60)	50 (40-55)	0.06
Leading symptom				
Angina	212 (89.1%)	95 (88.8%)	117 (89.3%)	0.29
Dyspnea	8 (3.3%)	6 (5.6%)	2 (1.5%)	
Syncope	2 (0.8%)	1 (0.9%)	1 (0.8%)	
A-V block	6 (2.5%)	3 (2.8%)	3 (2.3%)	
Hypotension	2 (0.8%)	0	2 (1.5%)	
VT	2 (0.8%)	0	1 (1.5%)	
SVT	1 (0.4%)	1(0.9%)	0	
Arrest	5 (2.1%)	4 (3.7%)	1 (0.8%)	
Diagnostic angiogram only	84 (35.0%)	47 (43.1%)	37 (28.2%)	0.01
History of CABG	7 (2.9%)	3 (2.8%)	4 (3.1%)	0.95

Procedural exitus	1 (0.4%)	0	1 (0.8%)	0.96
Acute stent thrombosis	10 (4.2%)	0	10 (7.6%)	0.003
Number of pictures taken during intervention (Median, Q1-Q3)	16 (8-22)	16 (8-22)	16 (8-23)	0.96
CABG referral	35(14.6%)	27 (25.0%)	8(6.1%)	<0.001

Table 1: Baseline clinical characteristics of the study patients

CAD:coronary artery disease, CHF:congestif heart failure, COPD:chronic obstructive pulmonary disease, SVE:cerebrovascular event,STEMI:ST- segment elevation myocardial infarction, NSTEMI:non st elevation myocardial infarction, USAP:unstabil angina pectoris, LVEF:left ventricular ejection fraction, A-V block:atrioventricular block, VT:ventricular tachycardia, SVT:supraventricular tachycardia, CABG:coronary artery by pass grefting.