

**Table 1. Demographics and clinical characteristics of study patients**

	Total (n:107)	not using ACEI (n:52)	using ACEI (n:55)	p
<b>Age (years)</b>	68.49±11.95(38-97)	70.13±10.16(43-91)	66.93±13.33(38-97)	0.163
<b>Gender</b>				
Male (n,%)	54(50.5)	27(51.9)	27(49.1)	0.770
Female (n,%)	53(49.5)	25(48.1)	28(50.9)	
<b>Past medical history(n,%)</b>				
Diabetes	51(47.7)	29(55.8)	22(40)	0.103
CAD	34(31.8)	18(34.6)	16(29.1)	0.540
COPD	11(10.3)	3(5.8)	8(14.5)	0.135
CRF	15(14)	9(17.3)	6(10.9)	0.341
<b>Prognosis (n,%)</b>				
Discharged	83(77.6)	35(67.3)	48(87.3)	<b>0.013</b>
Death	24(22.4)	17(32.7)	7(12.7)	
<b>No ICU care (n,%)</b>	84(78.5)	40(76.9)	44(80)	0.161
<b>ICU care (n,%)</b>	19(17.8)	12(23.1)	7(12.7)	
<b>Radiologic findings (n,%)</b>				
Unilateral ground-glass opacity	13(12.1)	7(13.5)	6(10.9)	0.686
Bilateral ground-glass opacity	90(84.1)	45(86.5)	45(81.8)	0.504
Multiple lobe lesions	57(53.3)	37(71.2)	20(36.4)	<b>&lt;0.001</b>
<b>Symptom duration (days)</b>	4(1-30)	4(1-30)	5(1-30)	0.406
<b>Clinical symptoms (n, %)</b>				
Fever	51(47.7)	26(50)	25(45.5)	0.638
Cough and sputum	57(53.8)	23(44.2)	34(63)	0.053
Dyspnea	49(45.8)	23(44.2)	26(47.3)	0.752
Sore throat	4(3.7)	1(1.9)	3(5.5)	0.618
Fatigue	30(28)	14(26.9)	16(29.1)	0.803
Diarrhea	15(14)	9(17.3)	6(10.9)	0.341
Chest pain	4(3.7)	1(1.9)	3(5.5)	0.618
Myalgia or arthralgia	6(5.6)	3(5.8)	3(5.5)	0.999
Anosmi	2(1.9)	0(0)	2(3.6)	0.496

Data are presented as mean±SD (range), median (range) or n (%). Student's t test, Mann-Whitney U test, Pearson chi-square test, Fisher's Exact test.

Abbreviations: CAD: coronary artery disease; COPD: chronic obstructive pulmonary disease; CRF: chronic renal failure; ICU: Intensive Care Unit.