

**Table 2: Use of Cardioscopy to Remove PFE**

Author	N	Tumor Location	Technique (Optics type)	Outcome
Je, H.G. <sup>36</sup>	1	LV	R Mini Thor, TM (Aesop robotic)	NAE, CPB 57 min, XC 24 min.
Espada R. <sup>38</sup>	1	LV	TM, Stern Stryker camera 30 degree; concomitant CABG	NAE
Kaneko Y. <sup>39</sup>	1	LV	TM Stern, right chest thoracoscopic assistance	NAE; CPB 79 min, XC 34 min.
Misumi T. <sup>40</sup>	2	LV	Pt 1 – TM Stern, Olympus endoscope Pt 2 – TA Stern, Olympus endoscope	NR
Ishida K. <sup>41</sup>	1	LV	TA Stern EndoEye Olympus scope	NAE.
Walkes <sup>42</sup>	1	LV	TA Stern, 2mm thoracoscope	NAE. No recurrence 8 months.
Toeg H.D. <sup>43</sup>	1	LV	TA Stern, video mediastinoscope	NAE. No recurrence 1 year.
Domenech A. et al. <sup>44</sup>	1	LV	Transmur Stern	NAE. No recurrence 1 year.
Matsuo Y. et al. <sup>45</sup>	1	LV	TA Stern fiberoptic scope	NAE
Ariyoshi T. et al. <sup>46</sup>	1	LV	TM Stern, videoscope	NAE
Allen K.B. et al. <sup>47</sup>	1	LV	TA Stern. videothoracoscope	NAE

**TM – transmitral; Stern – Sternotomy; TA – Transaortic; Mini Thor – Mini Thoracotomy; NAE – No adverse events; NR – Not reported**