

**Table 1.** Health professionals surveyed by age and sex, and occurrence of conflicts

<b>Health workers</b>	<b>Total (%)</b>	<b>Man</b>	<b>Woman</b>	<b>Age (<math>\mu</math>)</b>	<b>Experienced conflicts</b>
<b>Undergraduate intern doctors</b>	92 (1.5%)	48.9%	51.1%	23.7	56.0%
<b>Social service intern doctors</b>	767 (12.7%)	43.4%	56.6%	25.1	79.4%
<b>Resident doctors</b>	796 (13.2%)	49.9%	50.1%	28.8	46.8%
<b>Nurse practitioners</b>	1669 (27.7%)	20.7%	79.3%	24.0	37.9%
<b>Nursing technicians</b>	989 (16.4%)	24.9%	75.1%	21.2	29.9%
<b>Technicians in radiology</b>	49 (0.8%)	53.1%	46.9%	23.0	40.0%
<b>Odontology practitioner</b>	567 (9.4%)	28.2%	71.8%	24.4	23.3%
<b>Resident odontologist</b>	12 (0.2%)	41.7%	58.3%	29.6	40.0%
<b>Chemistry, Pharmacy and Biologist practitioner, Chemist-biologist parasitologist and Pharmacy</b>	113 (1.9%)	38.1%	61.9%	23.5	22.9%
<b>Clinical Laboratory technician</b>	122 (2.0%)	25.4%	74.6%	18.1	27.1%
<b>Rehabilitation, physical, and occupational therapy practitioners</b>	247 (4.1%)	20.2%	79.8%	23.5	23.9%
<b>Physical therapy and rehabilitation technician</b>	17 (0.3%)	47.1%	52.9%	24.0	28.6%
<b>Practitioners in nutrition</b>	408 (8.8%)	15.7%	84.3%	23.8	20.3%
<b>Nutrition and dietetics technician</b>	25 (0.4%)	8.0%	92.0%	23.7	8.7%
<b>Optometry technician</b>	9 (0.1%)	22.2%	77.8%	32.2	44.4%
<b>Psychology practitioners</b>	87 (1.4%)	23.0%	77.0%	24.1	11.0%
<b>Social work practitioners</b>	29 (0.5%)	17.2%	82.8%	25.1	28.6%
<b>Social work technician</b>	13 (0.2%)	7.7%	92.3%	19.4	10.0%

<b>Other professions*</b>	9 (0.1%)	33.3%	66.7%	23.1	0.0%
<b>Total</b>	6020 (100.0%)	29.7%	70.3%	24.2	39.5%

\*Engineering, physics, and administration practitioners who answered the survey