

Table 1. Differences of the studied variables between Event-No versus Event-Yes groups

Studied variables	Event-No (n=18)	Event-Yes (n=10)	P value
Age at diagnosis (mo), median (IQR)	15 (8.7 to 32.5)	15 (7.7 to 39.5)	0.88
Age at LT (mo), median (IQR)	27.7 (17.6 to 45.2)	29.6 (26 to 48.8)	0.38
Time from diagnosis to LT (mo), median (IQR)	7.5 (5 to 11)	13.5 (7.7 to 20)	0.11
Time from diagnosis to LT ≥ 12 mo, n(%)	4 (22.2)	6 (60)	0.09
Weight at LT (kg), median (IQR)	11.6 (10.1 to 14)	12 (11.1 to 17.4)	0.42
PELD, median (IQR)	-3.5 (-8 to -2.7)	- 8 (-10 to -1.7)	0.38
PRETEXT, n(%)			0.38
III	2 (11.2)	2 (20)	
IV	15 (83.3)	6 (60)	
Rescue LT, n(%)	1 (5.5)	2 (20)	0.28
Lung Metastasis Pre Qtx, n(%)	4 (22.2)	2 (20)	1
	N=15	N=8	
Type of PreLT Qtx, n(%)			0.22
PLADO	10 (66.7)	3 (37.5)	
Other	5 (33.3)	5 (62.5)	
	N=18	N=10	
PostLT Qtx, n(%)	15 (83.3)	10 (100)	0.28
AFP levels Pre Qtx, median (IQR)	15684.5 (1099.7 to 104600.5)	3000 (836.7 to 210126.2)	0.48
AFP levels Post Qtx, median (IQR)	2636 (107.9 to 12109)	15690.5 (2751.6 to 78516.5)	0.07
%AFP reduction Post Qtx, n(%)			0.05
<70%	5 (27.8)	7 (70)	
$\geq 70\%$	13 (72.2)	3 (30)	
Liver explant findings			
Vascular Invasion, n(%)			0.002
No	14 (77.8)	3 (30)	
Microvascular	3 (16.7)	3 (30)	

Macrovascular	1 (5.5)	4 (40)
---------------	---------	--------

Mo, months; IQR, interquartile range; LT, liver transplantation; Qtx, chemotherapy.