

	Gender n (%)	Median age (range)	Presenting features n (%)	HNC subsite n (%)	T classification n (%)	Treatment intent n (%)
2020	M – 23 (79.3) F – 6 (20.7)	68 (44-93)	Airway compromise – 16 (55.2) Dysphagia & malnutrition – 10 (34.5) Bleeding into UADT – 3 (10.3)	Larynx – 16 (55.2) Hypopharynx – 7 (24.1) Oropharynx – 4 (13.8) Thyroid – 2 (6.9)	T3/T4 – 27 (93.1) Other – 2 (6.9)	Curative – 15 (51.7) Palliative – 14 (48.3)
2019	M – 9 (81.8) F – 2 (18.2)	64 (50-88)	Airway compromise – 5 (45.5) Dysphagia & malnutrition – 3 (27.3) Bleeding into UADT – 1 (9.1) Bleeding from neck mass – 2 (18.2)	Larynx – 6 (54.5) Hypopharynx – 2 (18.2) Oropharynx – 1 (9.1) Neck – 2 (18.2)	T3/T4 – 10 (90.9) Other – 1 (9.1)	Curative – 7 (63.6) Palliative – 4 (36.4)
2018	M – 6 (66.7) F – 3 (33.3)	70 (56-91)	Airway compromise – 6 (66.7) Dysphagia & malnutrition – 1 (11.1) Bleeding into UADT – 1 (11.1) Bleeding from neck mass – 1 (11.1)	Larynx – 6 (66.7) Hypopharynx – 1 (11.1) Oropharynx – 1 (11.1) Neck – 1 (11.1)	T3/T4 – 9 (100)	Curative – 6 (66.7) Palliative – 3 (33.3)
2017	M – 10 (76.9) F – 3 (23.1)	68 (58-89)	Airway compromise – 7 (53.8) Dysphagia & malnutrition – 3 (23.1) Bleeding into UADT – 3 (23.1)	Larynx – 8 (61.5) Hypopharynx – 4 (30.8) Oropharynx – 1 (7.7)	T3/T4 – 11 (84.6) Other – 2 (15.4)	Curative – 8 (61.5) Palliative – 5 (38.5)
Total	M – 48 (77.4) F – 14 (22.6)	68 (44-93)	Airway compromise – 34 (54.8) Dysphagia & malnutrition – 17 (27.4) Bleeding into UADT – 8 (12.9) Bleeding from neck mass – 3 (4.8)	Larynx – 36 (58.1) Hypopharynx – 14 (22.6) Oropharynx – 7 (11.3) Neck – 3 (4.8) Thyroid – 2 (3.2)	T3/T4 – 57 (91.9) Other – 5 (8.1)	Curative – 36 (58.1) Palliative – 26 (41.9)

Table 1: Patient demographics, presenting symptoms, diagnosis, disease stage, and treatment intent for the study cohort as a whole and by year of presentation. Each year represents the six-month period of June – November inclusive. UADT = upper aerodigestive tract.