

	Delirium (n=96)	No-Delirium (n=340)	p value
Operation time (min)	128.8±36.5	130.2±46.0	.133
Anesthesia time (min)	141.3±56.2	151.5±68.5	.216
Preoperative anxiety	32(33.33%)	75(22.06%)	.065
Dexmedetomidine use	89(92.71%)	321(94.41%)	.256
Propofol use	92(95.83%)	332(97.65%)	.278
ASA PS			
II	35(36.46%)	72(21.18%)	.041
III	28(29.17%)	92(27.06%)	.102
Estimated blood loss (ml)	834.8±87.5	521.2±76.8	< .001
Transfusion (ml)	426.9±48.2	321.5±49.0	.003
Lowest heart rate	49.3±17.5	48.4±16.1	.102
Lowest mean arterial pressure (mmHg)	53.8±12.3	61.1±18.2	< .001
SAS	4.2±0.8	7.8±1.2	< .001
SAS 0-3	62(64.58%)	40(11.76%)	< .001
SAS 4-6	27(28.13%)	93(27.35%)	< .001
SAS 7-10	7(7.29%)	207(60.88%)	< .001

Table 3. Intraoperative anaesthesia data

Data are presented as n (%) or mean±SD. ASA PS, American Society of Anesthesiologists Physical Status; SAS, surgical Apgar score;