

Table 1. The unadjusted and adjusted independent associations between the predictor variable, formal engagement with allergy-trained clinicians, and four mutually exclusive clinical comfort outcome variables.

		Unadjusted Univariate Logistic Regression			Adjusted* Multivariate Logistic Regression		
		Odds Ratio (OR)	95% C.I.	<i>p</i> -value	Odds Ratio (OR)	95% C.I.	<i>p</i> -value
Outcome Variable: Comfort Evaluating Infant/Toddler Food Allergy							
Engagement with Allergy							
Yes		7.64	1.74 to 33.58	0.007	5.13	0.85 to 30.95	0.075
No		1.00	<i>Ref</i>		1.00	<i>Ref</i>	
Outcome Variable: Comfort Managing Infant/Toddler Food Allergy							
Engagement with Allergy							
Yes		4.83	0.84 to 27.93	0.078	2.12	0.25 to 17.78	0.487
No		1.00	<i>Ref</i>		1.00	<i>Ref</i>	
Outcome Variable: Comfort Diagnosing Infant/Toddler Anaphylaxis							
Engagement with Allergy							
Yes		2.53	0.68 to 9.39	0.166	1.23	0.26 to 5.91	0.795
No		1.00	<i>Ref</i>		1.00	<i>Ref</i>	
Outcome Variable: Comfort Treating Infant/Toddler Anaphylaxis							
Engagement with Allergy							
Yes		7.41	1.46 to 37.57	0.016	8.27	1.16 to 59.01	0.035
No		1.00	<i>Ref</i>		1.00	<i>Ref</i>	

95% *C.I.*, 95% confidence interval

*Adjusted for years postgraduate training