

Turán type inequalities for the supertrigonometric functions

Lu-Lu Geng^{1,2}, Xiao-Jun Yang^{1,2,3†}, Jian-Gen Liu^{1,2}

1. School of Mathematics, China University of Mining and Technology, Xuzhou, Jiangsu, 221116, P. R. China.

2. State Key Laboratory for Geomechanics and Deep Underground Engineering, China University of Mining and Technology, Xuzhou, Jiangsu, 221116, P. R. China.

3. School of Mechanics and Civil Engineering, China University of Mining and Technology, Xuzhou, Jiangsu, 221116, P. R. China.

Abstract: This paper is devoted to the study of Turán type inequalities for some well-known special functions such as supersine and supercosine which are derived by using a new form of the Cauchy-Bunyakovsky-Schwarz inequality.

Key words: Turán type inequality; Supersine; Supercosine; Gauss hypergeometric function; Cauchy-Bunyakovsky-Schwarz inequality.

Mathematics Subject Classification: 33C05; 33C20; 33C99.

Introduction

The importance, in many fields of mathematics, of the inequalities of the type proved by Turán [1]

$$f_n(x)f_{n+2}(x) - [f_{n+1}(x)]^2 \leq 0, \quad n = 0, 1, 2, \dots, \quad (1)$$

is the well known as Turán type inequalities[2].

Many Turán type inequalities have been investigated in the literature. For example, Joshi and Bissu [3] presented some two-sided inequalities for the ratio of modified Bessel functions of the first kind in 1991. Recently, Segura [4] introduced the bounds for ratios of modified Bessel functions associated with Turán type inequalities in 2011. Baricz [5] recommended Turán type inequalities for q -hypergeometric functions in 2013.

Throughout this paper, let R and N be the sets of the real and integral numbers. The discrete version of the well-known Cauchy-Schwarz inequality [6, 7]

$$\left(\sum_{i=1}^n a_i b_i\right)^2 \leq \sum_{i=1}^n a_i^2 \sum_{i=1}^n b_i^2, \quad a_i, b_i \in R, \quad (2)$$

and its integral representation in the space of continuous real-valued functions $C([a, b], R)$, i.e. the Cauchy-Bunyakovsky-Schwarz (CBS) inequality [6, 7]

$$\left(\int_a^b [u(t)]^{\frac{1}{2}} [v(t)]^{\frac{1}{2}} dt\right)^2 \leq \left(\int_a^b u(t) dt\right) \left(\int_a^b v(t) dt\right), \quad (3)$$

plays an important role in the different branches of modern mathematics.

[†]Corresponding author E-mail: dyangxiaojun@163.com

In 2006, Laforgia and Natalini [8] used the following form of the CBS inequality:

$$\left(\int_a^b g(t)[f(t)]^{\frac{m+n}{2}} dt\right)^2 \leq \left(\int_a^b g(t)[f(t)]^m dt\right)\left(\int_a^b g(t)[f(t)]^n dt\right), \quad (4)$$

to establish some new Turán type inequalities involving the special functions [9–12] as gamma, polygamma and Riemann's zeta function. Here f and g are the non-negative functions of a real variable and $m, n \in R$, R is the set of real numbers, such that the involved integrals in (1) exist.

In 2018, Bhandari and Bissu presented a new form of the generalized CBS inequality [13]

$$\left(\int_a^b [u(t)]^\eta [v(t)]^\eta [r(t)]^\eta dt\right)^2 \leq \left(\int_a^b [u(t)]^{\eta-l} [v(t)]^{\eta-m} [r(t)]^{\eta-n} dt\right)\left(\int_a^b [u(t)]^{\eta+l} [v(t)]^{\eta+m} [r(t)]^{\eta+n} dt\right), \quad (5)$$

in which $\eta, l, m, n \in R$, u, v , and r are real integrable functions such that the involved integrals in (5) exist.

In 2021, the supersine via Gauss hypergeometric series, proposed by author [14], was defined as

$$\begin{aligned} {}_2\text{Supersin}_1(a, b; c; x) &= \sum_{n=0}^{\infty} \frac{(a)_{2n+1}(b)_{2n+1}}{(c)_{2n+1}} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \\ &= \frac{\Gamma(c)}{2i\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} [(1-xt)^{-a} - (1+xt)^{-a}] dt, \end{aligned} \quad (6)$$

where $a, b, c, x \in R$, $n \in N$, and $|x| < 1$.

Meanwhile, the supercosine via Gauss hypergeometric series, proposed by author [14], was defined by

$$\begin{aligned} {}_2\text{Supercos}_1(a, b; c; x) &= \sum_{n=0}^{\infty} \frac{(a)_{2n}(b)_{2n}}{(c)_{2n}} \frac{(-1)^n x^{2n}}{(2n)!} \\ &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} {}_1\text{Supercos}_0(a; -; xt) dt, \end{aligned} \quad (7)$$

where $a, b, c, x \in R$, $n \in N$, and $|x| < 1$.

The aim of this paper is to study Turán type inequalities for the supersine and supercosine by using a new form of the generalized CBS inequality.

The structure of the paper is as follows: In Section 2, some of main results about Turán type inequality for the supersine and supercosine are obtained and proved in detail, In Section 3, we make a summary and prospect of this paper.

Main results

In this section, we will focus on Turán type inequalities for the supersine and supercosine.

Theorem 1. *Let $b > |\alpha|$, $c - b > |\beta|$ and $|x| \leq 1$. Then Turán type inequality for the supersine is given as*

$$\begin{aligned} [{}_2\text{Supersin}_1(a, b; c; x)]^2 &\leq \frac{B(b - \alpha, c - b - \beta)B(b + \alpha, c - b + \beta)}{[B(b, c - b)]^2} \\ &\quad \times {}_2\text{Supersin}_1(a - \gamma, b - \alpha; c - (\alpha + \beta); x) \\ &\quad \times {}_2\text{Supersin}_1(a + \gamma, b + \alpha; c + (\alpha + \beta); x). \end{aligned} \quad (8)$$

Proof. In the beginning, for (6), let $u(t) = t^{b-1}$, $v(t) = (1-t)^{c-b-1}$ and $r(t) = (1-xt)^{-a} - (1+xt)^{-a}$. Then the form of the supersine is as follows

$$\begin{aligned} \int_0^1 u(t)v(t)r(t)dt &= \int_0^1 t^{b-1}(1-t)^{c-b-1}[(1-xt)^{-a} - (1+xt)^{-a}]dt \\ &= \frac{2i\Gamma(b)\Gamma(c-b)}{\Gamma(c)} {}_2\text{Supersin}_1(a, b; c; x). \end{aligned} \quad (9)$$

On the basis of the inequality (5), we can receive

$$\begin{aligned} & \left(\int_0^1 t^{\eta(b-1)}(1-t)^{\eta(c-b-1)}[(1-xt)^{-a} - (1+xt)^{-a}]^\eta dt \right)^2 \\ & \leq \int_0^1 t^{(\eta-l)(b-1)}(1-t)^{(\eta-m)(c-b-1)}[(1-xt)^{-a} - (1+xt)^{-a}]^{\eta-n} dt \\ & \quad \times \int_0^1 t^{(\eta+l)(b-1)}(1-t)^{(\eta+m)(c-b-1)}[(1-xt)^{-a} - (1+xt)^{-a}]^{\eta+n} dt. \end{aligned} \quad (10)$$

Consequently, the following results can be obtained by applying (9) and (10),

$$\begin{aligned} & [{}_2\text{Supersin}_1(\eta a, \eta(b-1)+1; \eta(c-2)+2; x)]^2 \\ & \leq \frac{[\Gamma\{\eta(c-2)+2\}]^2}{[\Gamma\{\eta(b-1)+1\}]^2[\Gamma\{\eta(c-b-1)+1\}]^2} \\ & \quad \times \frac{\Gamma[(\eta-l)(b-1)+1]\Gamma[(\eta-m)(c-b-1)+1]}{\Gamma[(\eta-l)(b-1)+(\eta-m)(c-b-1)+2]} \\ & \quad \times \frac{\Gamma[(\eta+l)(b-1)+1]\Gamma[(\eta+m)(c-b-1)+1]}{\Gamma[(\eta+l)(b-1)+(\eta+m)(c-b-1)+2]} \\ & \quad \times {}_2\text{Supersin}_1((\eta-n)a, (\eta-l)(b-1)+1; (\eta-l)(b-1)+(\eta-m)(c-b-1)+2; x) \\ & \quad \times {}_2\text{Supersin}_1((\eta+n)a, (\eta+l)(b-1)+1; (\eta+l)(b-1)+(\eta+m)(c-b-1)+2; x). \end{aligned} \quad (11)$$

If $p_1 = \eta(b-1)+1$, $p_2 = \eta(c-b-1)+1$, $p_3 = \eta a$, $q_1 = l(b-1)$, $q_2 = m(c-b-1)$, and $q_3 = na$ in inequality (11), then Turán type inequality for the supersine can be written as

$$\begin{aligned} [{}_2\text{Supersin}_1(p_3, p_1; p_1+p_2; x)]^2 & \leq \frac{[\Gamma(p_1+p_2)]^2}{[\Gamma(p_1)]^2[\Gamma(p_2)]^2} \frac{\Gamma(p_1-q_1)\Gamma(p_2-q_2)}{\Gamma[(p_1-q_1)+(p_2-q_2)]} \frac{\Gamma(p_1+q_1)\Gamma(p_2+q_2)}{\Gamma[(p_1+q_1)+(p_2+q_2)]} \\ & \quad \times {}_2\text{Supersin}_1(p_3-q_3, p_1-q_1; (p_1-q_1)+(p_2-q_2); x) \\ & \quad \times {}_2\text{Supersin}_1(p_3+q_3, p_1+q_1; (p_1+q_1)+(p_2+q_2); x). \end{aligned} \quad (12)$$

Using the relationship between beta and gamma functions

$$B(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)}, \quad m, n > 0, \quad (13)$$

Turán type inequality can also be shown in the following form

$$\begin{aligned} [{}_2\text{Supersin}_1(p_3, p_1; p_1+p_2; x)]^2 & \leq \frac{B(p_1-q_1, p_2-q_2)B(p_1+q_1, p_2+q_2)}{[B(p_1, p_2)]^2} \\ & \quad \times {}_2\text{Supersin}_1(p_3-q_3, p_1-q_1; (p_1-q_1)+(p_2-q_2); x) \\ & \quad \times {}_2\text{Supersin}_1(p_3+q_3, p_1+q_1; (p_1+q_1)+(p_2+q_2); x), \end{aligned} \quad (14)$$

where $p_1 > |q_1|$, $p_2 > |q_2|$ and $|x| \leq 1$.

When $\eta = 1$, the inequality (11) reduces to following inequality for the supersine

$$\begin{aligned}
[{}_2\text{Supersin}_1(a, b; c; x)]^2 &\leq \frac{[\Gamma(c)]^2}{[\Gamma(b)]^2[\Gamma(c-b)]^2} \\
&\times \frac{\Gamma[b-l(b-1)]\Gamma[(c-b)-m(c-b-1)]}{\Gamma[c-l(b-1)-m(c-b-1)]} \\
&\times \frac{\Gamma[b+l(b-1)]\Gamma[(c-b)+m(c-b-1)]}{\Gamma[c+l(b-1)+m(c-b-1)]} \\
&\times {}_2\text{Supersin}_1((1-n)a, b-l(b-1); c-l(b-1)-m(c-b-1); x) \\
&\times {}_2\text{Supersin}_1((1+n)a, b+l(b-1); c+l(b-1)+m(c-b-1); x).
\end{aligned} \tag{15}$$

Suppose $l(b-1) = \alpha$, $m(c-b-1) = \beta$, $na = \gamma$, $b > |\alpha|$, $c-b > |\beta|$ and $|x| \leq 1$. Then, through (13) the inequality (15) transforms to the much nicer version of Turán type inequality for the supersine as follows

$$\begin{aligned}
[{}_2\text{Supersin}_1(a, b; c; x)]^2 &\leq \frac{B(b-\alpha, c-b-\beta)B(b+\alpha, c-b+\beta)}{[B(b, c-b)]^2} \\
&\times {}_2\text{Supersin}_1(a-\gamma, b-\alpha; c-(\alpha+\beta); x) \\
&\times {}_2\text{Supersin}_1(a+\gamma, b+\alpha; c+(\alpha+\beta); x).
\end{aligned} \tag{16}$$

Theorem 2. For $b > |\lambda|$, $c-b > |\mu|$ and $|x| \leq 1$, we have Turán type inequality for the supercosine

$$\begin{aligned}
[{}_2\text{Supercos}_1(a, b; c; x)]^2 &\leq \frac{B(b-\lambda, c-b-\mu)B(b+\lambda, c-b+\mu)}{[B(b, c-b)]^2} \\
&\times {}_2\text{Supercos}_1(a-\omega, b-\lambda; c-(\lambda+\mu); x) \\
&\times {}_2\text{Supercos}_1(a+\omega, b+\lambda; c+(\lambda+\mu); x).
\end{aligned} \tag{17}$$

Proof . At first, basing on (6), if $u(t) = t^{b-1}$, $v(t) = (1-t)^{c-b-1}$, and $r(t) = {}_1\text{Supercos}_0(a; -; xt)$, we have

$$\begin{aligned}
\int_0^1 u(t)v(t)r(t)dt &= \int_0^1 t^{b-1}(1-t)^{c-b-1} {}_1\text{Supercos}_0(a; -; xt)dt \\
&= \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} {}_2\text{Supercos}_1(a, b; c; x).
\end{aligned} \tag{18}$$

By the generalized CBS inequality, the following inequality for the supercosine can be written as

$$\begin{aligned}
&(\int_0^1 t^{\eta(b-1)}(1-t)^{\eta(c-b-1)} [{}_1\text{Supercos}_0(a; -; xt)]^\eta)^2 \\
&\leq \int_0^1 t^{(\eta-l)(b-1)}(1-t)^{(\eta-m)(c-b-1)} [{}_1\text{Supercos}_0(a; -; xt)]^{\eta-n} dt \\
&\times \int_0^1 t^{(\eta+l)(b-1)}(1-t)^{(\eta+m)(c-b-1)} [{}_1\text{Supercos}_0(a; -; xt)]^{\eta+n} dt.
\end{aligned} \tag{19}$$

Thus, the following results can be obtained by applying (18) and (19),

$$\begin{aligned}
& [{}_2\text{Supercos}_1(\eta a, \eta(b-1)+1; \eta(c-2)+2; x)]^2 \\
& \leq \frac{[\Gamma\{\eta(c-2)+2\}]^2}{[\Gamma\{\eta(b-1)+1\}]^2 [\Gamma\{\eta(c-b-1)+1\}]^2} \\
& \times \frac{\Gamma[(\eta-l)(b-1)+1] \Gamma[(\eta-m)(c-b-1)+1]}{\Gamma[(\eta-l)(b-1)+(\eta-m)(c-b-1)+2]} \\
& \times \frac{\Gamma[(\eta+l)(b-1)+1] \Gamma[(\eta+m)(c-b-1)+1]}{\Gamma[(\eta+l)(b-1)+(\eta+m)(c-b-1)+2]} \\
& \times {}_2\text{Supercos}_1((\eta-n)a, (\eta-l)(b-1)+1; (\eta-l)(b-1)+(\eta-m)(c-b-1)+2; x) \\
& \times {}_2\text{Supercos}_1((\eta+n)a, (\eta+l)(b-1)+1; (\eta+l)(b-1)+(\eta+m)(c-b-1)+2; x).
\end{aligned} \tag{20}$$

Next, let $p_1 = \eta(b-1)+1$, $p_2 = \eta(c-b-1)+1$, $p_3 = \eta a$, $q_1 = l(b-1)$, $q_2 = m(c-b-1)$, and $q_3 = na$ in inequality (20). Then we can receive Turán type inequality for the supercosine as follows:

$$\begin{aligned}
[{}_2\text{Supercos}_1(p_3, p_1; p_1+p_2; x)]^2 & \leq \frac{[\Gamma(p_1+p_2)]^2}{[\Gamma(p_1)]^2 [\Gamma(p_2)]^2} \frac{\Gamma(p_1-q_1) \Gamma(p_2-q_2)}{\Gamma[(p_1-q_1)+(p_2-q_2)]} \frac{\Gamma(p_1+q_1) \Gamma(p_2+q_2)}{\Gamma[(p_1+q_1)+(p_2+q_2)]} \\
& \times {}_2\text{Supercos}_1(p_3-q_3, p_1-q_1; (p_1-q_1)+(p_2-q_2); x) \\
& \times {}_2\text{Supercos}_1(p_3+q_3, p_1+q_1; (p_1+q_1)+(p_2+q_2); x).
\end{aligned} \tag{21}$$

On the other hand, for $p_1 > |q_1|$, $p_2 > |q_2|$ and $|x| \leq 1$, Turán type inequality for the supercosine can also be expressed in the following form by applying (13),

$$\begin{aligned}
[{}_2\text{Supercos}_1(p_3, p_1; p_1+p_2; x)]^2 & \leq \frac{B(p_1-q_1, p_2-q_2) B(p_1+q_1, p_2+q_2)}{[B(p_1, p_2)]^2} \\
& \times {}_2\text{Supercos}_1(p_3-q_3, p_1-q_1; (p_1-q_1)+(p_2-q_2); x) \\
& \times {}_2\text{Supercos}_1(p_3+q_3, p_1+q_1; (p_1+q_1)+(p_2+q_2); x).
\end{aligned} \tag{22}$$

After that, suppose $\eta = 1$. Then inequality (20) can be simplified to the following inequality

$$\begin{aligned}
[{}_2\text{Supercos}_1(a, b; c; x)]^2 & \leq \frac{[\Gamma(c)]^2}{[\Gamma(b)]^2 [\Gamma(c-b)]^2} \\
& \times \frac{\Gamma[b-l(b-1)] \Gamma[(c-b)-m(c-b-1)]}{\Gamma[c-l(b-1)-m(c-b-1)]} \\
& \times \frac{\Gamma[b+l(b-1)] \Gamma[(c-b)+m(c-b-1)]}{\Gamma[c+l(b-1)+m(c-b-1)]} \\
& \times {}_2\text{Supercos}_1((1-n)a, b-l(b-1); c-l(b-1)-m(c-b-1); x) \\
& \times {}_2\text{Supercos}_1((1+n)a, b+l(b-1); c+l(b-1)+m(c-b-1); x).
\end{aligned} \tag{23}$$

Finally, for $l(b-1) = \lambda$, $m(c-b-1) = \mu$, $na = \omega$, $b > |\lambda|$, $c-b > |\mu|$ and $|x| \leq 1$, by using (13) Turán type inequality (23) for the supercosine can be further simplified to

$$\begin{aligned}
[{}_2\text{Supercos}_1(a, b; c; x)]^2 & \leq \frac{B(b-\lambda, c-b-\mu) B(b+\lambda, c-b+\mu)}{[B(b, c-b)]^2} \\
& \times {}_2\text{Supercos}_1(a-\omega, b-\lambda; c-(\lambda+\mu); x) \\
& \times {}_2\text{Supercos}_1(a+\omega, b+\lambda; c+(\lambda+\mu); x).
\end{aligned} \tag{24}$$

Conclusion

In our work we had addressed the theorems about Turán type inequalities for the supersine and supercosine and proved it by using a new form of the generalized CBS inequality. This paper will strongly promote the rapid development of the field of special functions.

Acknowledgements

This work are supported by the Yue-Qi Scholar of the China University of Mining and Technology (No.102504180004), Graduate Research and Innovation Projects of Jiangsu Province(KYCX21_2145).

References

- [1] P. Turán. On the zeros of the polynomials of Legendre. *Macromolecular Chemistry and Physics*, 1950, 75(3): 113-122.
- [2] G. Szegő. On an inequality of P. Turán concerning Legendre polynomials. *Bulletin of the American Mathematical Society*, 1948, 54(4): 401-405.
- [3] C. M. Joshi, S. K. Bissu. Some inequalities of Bessel and modified Bessel functions. *Journal of the Australian Mathematical Society*, 1991, 50(2): 333-342.
- [4] J. Segura. Bounds for ratios of modified Bessel functions and associated Turán-type inequalities. *Journal of Mathematical Analysis and Applications* 2011, 374(2): 516-528.
- [5] A. Baricz, Raghavendar, et al. Turán type inequalities for q -hypergeometric functions. *Journal of Approximation Theory*, 2013, 168: 69-79.
- [6] H. Alzer. On the Cauchy-Schwarz inequality. *Journal of Mathematical Analysis and Applications*, 1999, 234(1): 6-14.
- [7] D. S. Mitrinović, J. E. Pečarić, and A. M. Fink. *Classical and New Inequalities in Analysis*. Dordrecht: Kluwer Academic; 1993.
- [8] A. Laforgia, P. Natalini. On some Turán type inequalities. *Journal of Inequalities and Applications*, 2006.
- [9] G. E. Andrews, R. Askey, R. Roy. *Special Functions*. Nature, 2001.
- [10] W. T. Sulaiman. Turán type inequalities for some special functions. *Australian Journal of Mathematical Analysis and Applications*, 2012, 9(1).
- [11] K. Mehrez, S. M. Sitnik. Turán type inequalities for classical and generalized Mittag-Leffler functions. *Analysis Mathematica*, 2018, 44(4): 521-541.
- [12] R. W. Barnard, M. B. Gordy, K. C. Richards. A note on Turán type and mean inequalities for the Kummer

- function. *Journal of Mathematical Analysis and Applications*, 2009, 349(1): 259-263.
- [13] P. K. Bhandari, S. K. Bissu. Turán type inequalities for Gauss and confluent hypergeometric functions via Cauchy-Bunyakovsky-Schwarz inequality. *Communications of the Korean Mathematical Society*, 2018, 33(4): 1285-1301.
- [14] X. J. Yang. *An Intruduction to Hypergeometric, Supertrigonometric, and Superhyperbolic functions*. Elsevier, 2021.