

Impulsive fractional partial differential system and its correct integral solution*

Xianmin Zhang^{1†}, Zuohua Liu², Zuming Peng¹, Yali He¹, Shixian Yang¹

1. School of Mathematics and Statistics, Yangtze Normal University, Chongqing 408100, China

2. School of Chemistry and Chemical Engineering, Chongqing University, Chongqing 400044, China

Abstract: It is found that there are two piecewise functions satisfy the conditions in the impulsive fractional partial differential system (IFrPDS), which deduce that the three different integral solutions of the IFrPDS given in the cited papers are inappropriate. Next, by applying two limit properties of the IFrPDS and the properties of piecewise function, the new formula of solution of the IFrPDS is discovered that is the integral equation with an arbitrary continuously differentiable function of t on $[0, c]$ to reveal the non-uniqueness of the IFrPDE's solution. Finally, an example is provided to expound the computation of the solution of the IFrPDS.

Key Words: impulsive fractional partial differential equations; equivalent integral equation; non-uniqueness of solution

MSC2010: 26A33; 34A08; 34A37; 34A12

1 Introduction

Impulsive fractional order system (IFOS) has become a focus of research and several hundreds articles are found by searching the topic of impulsive fractional order system from Web of science. For the IFOS, its integral solution (or equivalent integral equation) is key in studying numerical solution [1, 2], existence of solution [3–11], oscillation behavior [12, 13], periodic motion [14], solvability [15], asymptotic behavior of solution [16], stability [17–19], integral solution [20–27] and general solution [28] etc.

Furthermore, the impulsive fractional partial differential order system (IFrPDOS) was firstly introduced in [29] by

$$\begin{cases} {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta, s \neq s_j \ (j = 1, 2, \dots, J), \\ \Upsilon(s_j^+, t) - \Upsilon(s_j^-, t) = U_j(\Upsilon(s_j^-, t)), & t \in [0, c], j = 1, 2, \dots, J, \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c], \end{cases} \quad (1.1)$$

where ${}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e$ represents the Caputo fractional derivative with order $e = (e_1, e_2)$ ($e_1, e_2 \in (0, 1]$), $\Theta = [0, b] \times [0, c]$, $b = s_{J+1} > s_J > \dots > s_1 > s_0 = 0$, $F : \Theta \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $U_j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ ($j = 1, 2, \dots, J$),

*
†Corresponding: z6x2m@126.com (X. Zhang)

$\eta : [0, b] \rightarrow \mathbb{R}^n$ and $\xi : [0, c] \rightarrow \mathbb{R}^n$ with $\eta(0) = \xi(0)$.

Remark 1.1. Let $\Theta_0 = [0, s_1] \times [0, c]$, $\Theta_j = (s_j, s_{j+1}] \times [0, c]$, $\bar{\Theta}_0 = \Theta_0$ and $\bar{\Theta}_j = [s_j, s_{j+1}] \times [0, c]$ ($j = 1, 2, \dots, J$), $\Lambda(s, t) = \eta(s) + \xi(t) - \eta(0)$ and $F d\tau d\epsilon = F(\epsilon, \tau, w(\epsilon, \tau)) d\tau d\epsilon$ throughout this paper.

For (1.1), it is found that there are three different integral solution in existing research. To study some properties of solution, the integral solution of (1.1) was given in [29–32] by

$$\Upsilon(s, t) = \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s_j^+, t) + \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_j, j = 1, \dots, J, \end{cases} \quad (1.2)$$

or equivalently,

$$\Upsilon(s, t) = \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s, t) + \sum_{i=1}^j \int_{s_{i-1}}^{s_i} \int_0^t \frac{(s_i-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ + \sum_{i=1}^j [U_i(\Upsilon(s_i^-, t)) - U_i(\Upsilon(s_i^-, 0))] + \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \\ (s, t) \in \Theta_j, j = 1, 2, \dots, J. \end{cases} \quad (1.3)$$

In contrast, in [33], it was thought that the integral solution of (1.1) is another integral equality

$$\Upsilon(s, t) = \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s, t) + \sum_{i=1}^j [U_i(\Upsilon(s_i^-, t)) - U_i(\Upsilon(s_i^-, 0))] \\ + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_j, j = 1, 2, \dots, J. \end{cases} \quad (1.4)$$

Moreover, by combining the above two thoughts of integral solution, the third integral solution of (1.1) was given in [34].

Remark 1.2. By Lemma 2.3, in fact, (1.2) and (1.3) are the integral solution of the system

$$\begin{cases} {}^C_{(s_i^+, 0+)} \mathcal{D}_{(s,t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta_i, \text{ here } i = 0, 1, \dots, J, \\ \Upsilon(s_j^+, t) - \Upsilon(s_j^-, t) = U_j(\Upsilon(s_j^-, t)), & t \in [0, c], j = 1, 2, \dots, J, \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c], \end{cases} \quad (1.5)$$

However, recently it is found that there are two piecewise functions satisfying the conditions in (1.1), which derive that the given integral solutions in the cited papers are inappropriate. Therefore, in what follows, we will re-search the integral solution of (1.1).

We arrange the remainder of the paper as follows. Firstly, we review some definitions and properties of the fractional derivative and the fractional integration in Section 2. Secondly, we elaborate that there are two piecewise functions satisfying the conditions in (1.1), and then we re-explore the integral solution of (1.1) by some properties of piecewise function in Section 3. Finally, an example is provided to show the computation of integral solution of the IFrPDOS's solution in Section 4.

2 Preliminaries

Let $L^1(\Theta, \mathbb{R}^n)$ denote the space of the Lebesgue-integrable function $\Upsilon : \Theta \rightarrow \mathbb{R}^n$ with

$$\|\Upsilon(\mu, \nu)\|_1 = \int_0^b \int_0^c \|\Upsilon(\mu, \nu)\| d\nu d\mu$$

where $\|\cdot\|$ is an appropriate complete norm on \mathbb{R}^n .

Definition 2.1 ([35, 36]). Let $z \in [0, b]$, $\Theta_z = [z, b] \times [0, c]$ and $u = (u_1, u_2)$ ($u_1, u_2 > 0$). For $\Upsilon \in L^1(\Theta_z, \mathbb{R}^n)$, its mixed Riemann-Liouville fractional integral is defined by

$${}_{(z+,0+)}\mathcal{I}_{(s,t)}^u \Upsilon(s, t) = \int_z^s \int_0^t \frac{(s-\epsilon)^{u_1-1}}{\Gamma(u_1)} \frac{(t-\tau)^{u_2-1}}{\Gamma(u_2)} \Upsilon(\epsilon, \tau) d\tau d\epsilon.$$

Definition 2.2 [29]. Let $e = (e_1, e_2)$ ($e_1, e_2 \in (0, 1]$) and $\Upsilon : \Theta_z \rightarrow \mathbb{R}^n$ satisfy $\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \in L^1(\Theta_z, \mathbb{R}^n)$. the Caputo fractional derivative of Υ with order e is defined by

$${}_{(z+,0+)}^C \mathcal{D}_{(s,t)}^e \Upsilon(s, t) = {}_{(z+,0+)}\mathcal{I}_{(s,t)}^{1-e} \left(\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \right).$$

Lemma 2.3 [32]. Let the function $g : \Theta_z \rightarrow \mathbb{R}^n$ satisfy $\frac{\partial^2}{\partial s \partial t} g(s, t) \in C(\Theta_z, \mathbb{R}^n)$. If $\Upsilon(s, t) \in C(\Theta_z, \mathbb{R}^n)$, then $\Upsilon(s, t)$ satisfies

$${}_{(z+,0+)}^C \mathcal{D}_{(s,t)}^e \Upsilon(s, t) = g(s, t), \quad (s, t) \in \Theta_z,$$

iff $\Upsilon(s, t)$ satisfies

$$\begin{aligned} \Upsilon(s, t) &= \Upsilon(s, 0) + \Upsilon(z+, t) - \Upsilon(z+, 0) + {}_{(z+,0+)}\mathcal{I}_{(s,t)}^e g(s, t) \\ &= \Upsilon(s, 0) + \Upsilon(z+, t) - \Upsilon(z+, 0) + \int_z^s \int_0^t \frac{(s-\epsilon)^{e_1-1} (t-\tau)^{e_2-1} g(\epsilon, \tau)}{\Gamma(e_1) \Gamma(e_2)} d\tau d\epsilon, \\ &\text{for } (s, t) \in \Theta_z. \end{aligned}$$

Next, we use *Definitions 2.1-2.2* to consider the properties of piecewise function. Let

$$\begin{aligned} f(s, t) &= \begin{cases} f_0(s, t), & (s, t) \in \Theta_0, \\ f_1(s, t), & (s, t) \in \Theta_1, \\ \vdots \\ f_J(s, t), & (s, t) \in \Theta_J, \end{cases} \\ &= \begin{cases} f_0(s, t), & (s, t) \in \Theta_0, \\ 0, & (s, t) \in \cup_{k=1}^J \Theta_k, \end{cases} + \begin{cases} 0, & (s, t) \in \Theta_0, \\ f_1(s, t), & (s, t) \in \Theta_1, \\ 0, & (s, t) \in \cup_{k=2}^J \Theta_k, \end{cases} \\ &\quad + \dots + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-2} \Theta_k, \\ f_{J-1}(s, t), & (s, t) \in \Theta_{J-1}, \\ 0, & (s, t) \in \Theta_J, \end{cases} + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-1} \Theta_k, \\ f_J(s, t), & (s, t) \in \Theta_J, \end{cases} \end{aligned} \tag{2.1}$$

Property 2.4. Let $e = (e_1, e_2)$ (here $e_1, e_2 \in (0, 1]$) and $f_k : \bar{\Theta}_k \rightarrow \mathbb{R}^n$ satisfy $\frac{\partial^2}{\partial s \partial t} f_k(s, t) \in C(\bar{\Theta}_k, \mathbb{R}^n)$ (here $k = 0, 1, \dots, J$), then the fractional derivative of (2.1) can be computed by

$$\begin{aligned}
{}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e f(s, t) = & \begin{cases} {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e f_0(s, t), & (s, t) \in \Theta_0, \\ \int_0^{s_1} \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 f_0(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon, & (s, t) \in \cup_{k=1}^J \Theta_k, \end{cases} \\
& + \begin{cases} 0, & (s, t) \in \Theta_0, \\ {}^C_{(s_1^+,0+)}\mathcal{D}_{(s,t)}^e f_1(s, t), & (s, t) \in \Theta_1, \\ \int_{s_1}^{s_2} \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 f_1(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon, & (s, t) \in \cup_{k=2}^J \Theta_k, \end{cases} \\
& + \dots + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-2} \Theta_k, \\ {}^C_{(s_{J-1}^+,0+)}\mathcal{D}_{(s,t)}^e f_{J-1}(s, t), & (s, t) \in \Theta_{J-1}, \\ \int_{s_{J-1}}^{s_J} \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 f_{J-1}(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon, & (s, t) \in \Theta_J, \end{cases} \\
& + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-1} \Theta_k, \\ {}^C_{(s_J^+,0+)}\mathcal{D}_{(s,t)}^e f_J(s, t), & (s, t) \in \Theta_J. \end{cases}
\end{aligned} \tag{2.2}$$

Property 2.5. Let $e = (e_1, e_2)$ (here $e_1, e_2 \in (0, 1]$) and $f_k : \bar{\Theta}_k \rightarrow \mathbb{R}^n$ satisfy $f_k(s, t) \in C(\bar{\Theta}_k, \mathbb{R}^n)$ (here $k = 0, 1, \dots, J$), then the fractional integral of (2.1) can be computed by

$$\begin{aligned}
{}_{(0+,0+)}\mathcal{I}_{(s,t)}^e f(s, t) = & \begin{cases} {}_{(0+,0+)}\mathcal{I}_{(s,t)}^e f_0(s, t), & (s, t) \in \Theta_0, \\ \int_0^{s_1} \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1} f_0(\epsilon, \tau)}{\Gamma(e_1)\Gamma(e_2)} d\tau d\epsilon, & (s, t) \in \cup_{k=1}^J \Theta_k, \end{cases} \\
& + \begin{cases} 0, & (s, t) \in \Theta_0, \\ {}_{(s_1^+,0+)}\mathcal{I}_{(s,t)}^e f_1(s, t), & (s, t) \in \Theta_1, \\ \int_{s_1}^{s_2} \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1} f_1(\epsilon, \tau)}{\Gamma(e_1)\Gamma(e_2)} d\tau d\epsilon, & (s, t) \in \cup_{k=2}^J \Theta_k, \end{cases} \\
& + \dots + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-2} \Theta_k, \\ {}_{(s_{J-1}^+,0+)}\mathcal{I}_{(s,t)}^e f_{J-1}(s, t), & (s, t) \in \Theta_{J-1}, \\ \int_{s_{J-1}}^{s_J} \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1} f_{J-1}(\epsilon, \tau)}{\Gamma(e_1)\Gamma(e_2)} d\tau d\epsilon, & (s, t) \in \Theta_J, \end{cases} \\
& + \begin{cases} 0, & (s, t) \in \cup_{k=0}^{J-1} \Theta_k, \\ {}_{(s_J^+,0+)}\mathcal{I}_{(s,t)}^e f_J(s, t), & (s, t) \in \Theta_J. \end{cases}
\end{aligned} \tag{2.3}$$

3 The integral solution of (1.1)

We first present some limit properties of (1.1) and next illustrate that there are two piecewise functions satisfying the conditions in (1.1) to expound that the above integral solutions of (1.1) given in existing papers are inappropriate, and finally use the basic limit properties and *Properties 2.4-2.5* to re-explore the correct integral solution of (1.1) in this section.

For simplifying some formulas, let

$$\begin{aligned}\Phi_i(s, t) &= \int_{s_i}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad i = 0, 1, \dots, J; \\ \Psi_j(s, t) &= \int_0^{s_j} \int_0^t \frac{(s_j-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon + \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ &\quad - \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad j = 1, 2, \dots, J.\end{aligned}\tag{3.1}$$

3.1 The basic limit properties of (1.1)

Consider two limit cases of (1.1):

$$\begin{aligned}&\lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0 \text{ for } \forall j \in \{1, 2, \dots, J\}} \{\text{system (1.1)}\} \\ &= \begin{cases} {}^C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta, \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c], \end{cases} \\ &\Leftrightarrow \Upsilon(s, t) = \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad (s, t) \in \Theta;\end{aligned}\tag{3.2}$$

$$\begin{aligned}&\lim_{s_j \rightarrow s_l \text{ for } \forall j \in \{1, 2, \dots, J\} \text{ and } l \in \{1, 2, \dots, J\}} \{\text{system (1.1)}\} \\ &= \begin{cases} {}^C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta, s \neq s_l, \\ \Upsilon(s_l^+, t) - \Upsilon(s_l^-, t) = \sum_{j=1}^J U_j(\Upsilon(s_l^-, t)), & t \in [0, c], \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c]. \end{cases}\end{aligned}\tag{3.3}$$

3.2 Two piecewise functions satisfy the conditions in (1.1)

We will expound that there are two piecewise functions satisfying the conditions in (1.1).

Because $\Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon$ satisfies ${}^C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t))$ on whole interval Θ , it is sure that the piecewise function (1.4) satisfies the conditions of (1.1).

Next we will discuss whether (1.2) meets the fractional derivative of (1.1). Use *Definition 2.1* to calculate the fractional derivative of (1.2):

$$\begin{aligned}&{}^C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) \Big|_{(s, t) \in \Theta_0} = F(s, t, \Upsilon(s, t)), \quad (s, t) \in \Theta_0, \\ &{}^C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) \Big|_{(s, t) \in \Theta_j} = \int_0^{s_1} \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \Phi_0(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon \\ &\quad + \int_{s_1}^{s_2} \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \Phi_1(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon + \dots \\ &\quad + \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \Phi_j(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon, \quad (s, t) \in \Theta_j \\ &\quad \neq \int_0^s \int_0^t \frac{(s-\epsilon)^{-e_1}(t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \Phi_0(\epsilon, \tau)}{\partial \epsilon \partial \tau} d\tau d\epsilon, \quad (s, t) \in \Theta_j \\ &= F(s, t, \Upsilon(s, t)), \quad (s, t) \in \Theta_j \text{ (here } j \in \{1, 2, \dots, J\}),\end{aligned}\tag{3.4}$$

which means that (1.2) doesn't satisfy the fractional derivative of (1.1).

However, we can use (1.2) to construct another piecewise function satisfying the conditions in (1.1) as follows

$$\begin{aligned} \Upsilon(s, t) = & \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s_j^+, t) + \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_j, j = 1, 2, \dots, J, \end{cases} \\ & - \sum_{i=0}^{J-1} \begin{cases} 0, & (s, t) \in [s_0, s_{i+1}] \times [0, c], \\ \int_{s_{i+1}}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \left[\int_{s_i}^{s_{i+1}} \int_0^\tau \frac{(\epsilon-u)^{-e_1}(\tau-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \right. \\ \quad \left. \times \frac{\partial^2[\Phi_i(u, v)]}{\partial u \partial v} dv du \right] d\tau d\epsilon, & (s, t) \in (s_{i+1}, b] \times [0, c]. \end{cases} \end{aligned} \quad (3.5)$$

We use (2.2) to obtain the fractional derivative of (3.5)

$$\begin{aligned} & {}^C_{(0+,0+)} \mathcal{D}_{(s,t)}^e \Upsilon(s, t) \\ &= \begin{cases} F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta_0, \\ \int_{s_0}^{s_1} \int_0^t \frac{(s-u)^{-e_1}(t-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\Phi_0(u, v)]}{\partial u \partial v} dv du, & (s, t) \in (s_1, b] \times [0, c], \end{cases} \\ &+ \sum_{j=1}^{J-1} \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta_j, \\ \int_{s_j}^{s_{j+1}} \int_0^t \frac{(s-u)^{-e_1}(t-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\Phi_j(u, v)]}{\partial u \partial v} dv du, & (s, t) \in (s_{j+1}, b] \times [0, c], \end{cases} \\ &+ \begin{cases} 0, & (s, t) \in [0, s_J] \times [0, c], \\ F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta_J, \end{cases} \\ &- \sum_{i=0}^{J-1} \begin{cases} 0, & (s, t) \in [0, s_{i+1}] \times [0, c], \\ \int_{s_i}^{s_{i+1}} \int_0^t \frac{(s-u)^{-e_1}(t-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\Phi_i(u, v)]}{\partial u \partial v} dv du, & (s, t) \in (s_{i+1}, b] \times [0, c], \end{cases} \\ &= F(s, t, \Upsilon(s, t)), \quad (s, t) \in \cup_{i=0}^J \Theta_i. \end{aligned} \quad (3.6)$$

Therefore, it can be verified that both (1.4) and (3.5) satisfy these conditions of the initial value, the impulses and the fractional derivative in (1.1).

Additionally, by the above discussion, (1.4) is only a *special solution* of (1.1) because it doesn't contain the key part $\Phi_j(s, t)$ ($j = 1, 2, \dots, J$), but (3.5) is not the solution of (1.1) because it doesn't satisfy (3.2).

3.3 The integral solution of (1.1)

We will use *Properties 2.4-2.5*, two limit properties ((3.2) and (3.3)) and the thoughts of (1.4) and (3.5) to re-explore the integral solution of (1.1). Define the space of functions

$$\begin{aligned}
IC_P(\Theta, \mathbb{R}^n) := & \left\{ \Upsilon(s, t) : \cup_{i=0}^J \Theta_i \rightarrow \mathbb{R}^n \text{ and } \frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \in C(\cup_{i=0}^J \Theta_i, \mathbb{R}^n), \right. \\
& \Upsilon(s_j^-, t) = \lim_{s \rightarrow s_j^-} \Upsilon(s, t) = \Upsilon(s_j, t) < \infty, \quad \Upsilon(s_j^+, t) = \lim_{s \rightarrow s_j^+} \Upsilon(s, t) < \infty, \\
& \left. \lim_{s \rightarrow s_j^-} \left[\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \right] < \infty, \lim_{s \rightarrow s_j^+} \left[\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \right] < \infty, \text{ here } j = 1, 2, \dots, J \right\}.
\end{aligned}$$

Lemma 3.1. Let $F : \Theta \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfy $\frac{\partial^2}{\partial s \partial t} F(s, t, \Upsilon(s, t)) \in C(\bar{\Theta}_i \times \mathbb{R}^n, \mathbb{R}^n)$ for $\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \in C(\bar{\Theta}_i, \mathbb{R}^n)$ (here $i = 0, 1, \dots, J$), and let $\rho(t)$ be an arbitrary continuously differentiable function of t on $[0, c]$, then

$$\begin{aligned}
C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e & \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \rho(t) \Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1 - 1} (t - \tau)^{e_2 - 1}}{\Gamma(e_1) \Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon - u)^{-e_1} (\tau - v)^{-e_2}}{\Gamma(1 - e_1) \Gamma(1 - e_2)} \frac{\partial^2 [\rho(v) \Phi_0(u, v)]}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ (s, t) \in (s_j, b] \times [0, c], \text{ here } j = 1, 2, \dots, J; \end{cases} = 0. \quad (3.7)
\end{aligned}$$

The proof will be given in the *Appendix*.

Theorem 3.2. Let $F : \Theta \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfy $\frac{\partial^2}{\partial s \partial t} F(s, t, \Upsilon(s, t)) \in C(\bar{\Theta}_i \times \mathbb{R}^n, \mathbb{R}^n)$ for $\frac{\partial^2}{\partial s \partial t} \Upsilon(s, t) \in C(\bar{\Theta}_i, \mathbb{R}^n)$ (here $i = 0, 1, \dots, J$), and let $\lambda(t)$ be an arbitrary continuously differentiable function of t on $[0, c]$.

If $\Upsilon(s, t) \in IC_P(\Theta, \mathbb{R}^n)$, then $\Upsilon(s, t)$ satisfies (1.1) iff $\Upsilon(s, t)$ satisfies

$$\begin{aligned}
\Upsilon(s, t) = & \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s - \epsilon)^{e_1 - 1} (t - \tau)^{e_2 - 1}}{\Gamma(e_1) \Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s, t) + \int_0^s \int_0^t \frac{(s - \epsilon)^{e_1 - 1} (t - \tau)^{e_2 - 1}}{\Gamma(e_1) \Gamma(e_2)} F d\tau d\epsilon \\ \quad + \sum_{k=1}^j [U_k(\Upsilon(s_k^-, t)) - U_k(\Upsilon(s_k^-, 0))], & (s, t) \in \Theta_j, j = 1, 2, \dots, J, \end{cases} \\
& + \sum_{j=1}^J \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \lambda(t) U_j(\Upsilon(s_j^-, t)) \Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1 - 1} (t - \tau)^{e_2 - 1}}{\Gamma(e_1) \Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon - u)^{-e_1} (\tau - v)^{-e_2}}{\Gamma(1 - e_1) \Gamma(1 - e_2)} \frac{\partial^2 [\lambda(v) U_j(\Upsilon(s_j^-, v)) \Phi_0(u, v)]}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ (s, t) \in (s_j, b] \times [0, c], \end{cases} \quad (3.8)
\end{aligned}$$

Proof. 'Necessity'. It will be proved that the integral solution of (1.1) is (3.8). Let us consider (1.1) with an impulse:

$$\begin{cases} C_{(0+, 0+)} \mathcal{D}_{(s, t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta, s \neq s_j, \\ \Upsilon(s_j^+, t) - \Upsilon(s_j^-, t) = U_j(\Upsilon(s_j^-, t)), & t \in [0, c], \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c], \end{cases} \quad (3.9)$$

By Lemma 2.3 and the limit property (3.2), the integral solution of (3.9) satisfies

$$\Upsilon(s, t) = \Lambda(s, t) + \int_0^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad (s, t) \in [0, s_j] \times [0, c]; \quad (3.10)$$

and

$$\lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0} \Upsilon(s, t) = \Lambda(s, t) + \int_0^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad (s, t) \in \Theta. \quad (3.11)$$

Next, by using (3.10), it is similar with (3.5) to construct a piecewise function $\tilde{\Upsilon}(s, t)$

$$\begin{aligned} & \tilde{\Upsilon}(s, t) \\ &= \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in [0, s_j] \times [0, c], \\ \Lambda(s, t) + U_j(\Upsilon(s_j^-, t)) - U_j(\Upsilon(s_j^-, 0)) + \int_0^{s_j} \int_0^t \frac{(s_j - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ \quad + \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in (s_j, b] \times [0, c], \end{cases} \\ & - \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon - u)^{-e_1} (\tau - v)^{-e_2}}{\Gamma(1 - e_1)\Gamma(1 - e_2)} \frac{\partial^2 \Phi_0(u, v)}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ & (s, t) \in (s_j, b] \times [0, c], \end{cases} \end{aligned} \quad (3.12)$$

Remark 3.3. (3.12) satisfies these conditions of the initial value, the impulses and fractional derivative in (3.9), but (3.12) doesn't satisfy (3.11) to be only as an approximate solution with $e(s, t) = \Upsilon(s, t) - \tilde{\Upsilon}(s, t)$ for $(s, t) \in \Theta$.

Thus, by (3.11) and (3.12), we have

$$\begin{aligned} & \lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0} e(s, t) = \lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0} \left\{ \Upsilon(s, t) - \tilde{\Upsilon}(s, t) \right\} \quad \text{for } (s, t) \in \Theta, \\ &= \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon - u)^{-e_1} (\tau - v)^{-e_2}}{\Gamma(1 - e_1)\Gamma(1 - e_2)} \frac{\partial^2 \Phi_0(u, v)}{\partial u \partial v} dv du \right] d\tau d\epsilon \\ \quad - \Psi_j(s, t), & (s, t) \in (s_j, b] \times [0, c], \end{cases} \end{aligned} \quad (3.13)$$

From (3.13), we think that $e(s, t)$ connects with $U_j(\Upsilon(s_j^-, t))$ and $\lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0} e(s, t)$ and assume

$$\begin{aligned} & e(s, t) = g\left(U_j(\Upsilon(s_j^-, t))\right) \lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0} e(s, t) \quad (\text{here } g(\cdot) \text{ is an undetermined function}) \\ &= - \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ g\left(U_j(\Upsilon(s_j^-, t))\right) \Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s - \epsilon)^{e_1-1} (t - \tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon - u)^{-e_1} (\tau - v)^{-e_2}}{\Gamma(1 - e_1)\Gamma(1 - e_2)} \frac{\partial^2 \left[g(U_j(\Upsilon(s_j^-, v))) \Phi_0(u, v) \right]}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ & (s, t) \in (s_j, b] \times [0, c], \end{cases} \end{aligned} \quad (3.14)$$

By (3.12) and (3.14), we get

$$\begin{aligned}
\Upsilon(s, t) &= \tilde{\Upsilon}(s, t) + e(s, t) \quad \text{for } (s, t) \in \Theta, \\
&= \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in [0, s_j] \times [0, c], \\ \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ \quad + U_j(\Upsilon(s_j^-, t)) - U_j(\Upsilon(s_j^-, 0)), & (s, t) \in (s_j, b] \times [0, c], \end{cases} \\
&\quad + \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \left[1 - g\left(U_j(\Upsilon(s_j^-, t))\right) \right] \Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon-u)^{-e_1}(\tau-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \left[(1 - g(U_j(\Upsilon(s_j^-, v)))) \Phi_0(u, v) \right]}{\partial u \partial v} dv du \right] d\tau d\epsilon \\ & (s, t) \in (s_j, b] \times [0, c], \end{cases} \quad (3.15)
\end{aligned}$$

Furthermore, by (3.2) and (3.3), it shows that the impulsive effects have the linear additivity. Therefore, by combining with the special solution (1.4), we obtain the solution of (1.1) as

$$\begin{aligned}
\Upsilon(s, t) &= \begin{cases} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, & (s, t) \in \Theta_0, \\ \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ \quad + \sum_{i=1}^j [U_i(\Upsilon(s_i^-, t)) - U_i(\Upsilon(s_i^-, 0))], & (s, t) \in \Theta_j, j = 1, 2, \dots, J, \end{cases} \\
&\quad + \sum_{j=1}^J \begin{cases} 0, & (s, t) \in [0, s_j] \times [0, c], \\ \left[1 - g\left(U_j(\Upsilon(s_j^-, t))\right) \right] \Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon-u)^{-e_1}(\tau-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 \left[(1 - g(U_j(\Upsilon(s_j^-, v)))) \Phi_0(u, v) \right]}{\partial u \partial v} dv du \right] d\tau d\epsilon \\ & (s, t) \in (s_j, b] \times [0, c], \end{cases} \quad (3.16)
\end{aligned}$$

Moreover, (3.16) satisfies (3.3) such that

$$\begin{aligned}
1 - g[U_i(\Upsilon(s_i^-, t)) + U_j(\Upsilon(s_j^-, t))] &= 1 - g(U_i(\Upsilon(s_i^-, t))) + 1 - g(U_j(\Upsilon(s_j^-, t))) \\
&\quad \text{for } \forall U_i(\Upsilon(s_i^-, t)), U_j(\Upsilon(s_j^-, t)) \in \mathbb{R}^n. \quad (3.17)
\end{aligned}$$

Then

$$1 - g(U_j(\Upsilon(s_j^-, t))) = \lambda(t) U_j(\Upsilon(s_j^-, t)) \quad (3.18)$$

here $\lambda(t)$ is an arbitrary continuously differentiable function, and then (3.16) is (3.8).

'Sufficiency'. By *Lemma 3.1*, we obtain the fractional derivative of a part of (3.8):

$${}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e \left\{ \begin{array}{l} 0, \quad (s,t) \in [0, s_j] \times [0, c], \\ \lambda(t)U_j(\Upsilon(s_j^-, t))\Psi_j(s, t) - \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \\ \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon-u)^{-e_1}(\tau-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2 [\lambda(v)U_j(\Upsilon(s_j^-, v))\Phi_0(u, v)]}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ (s,t) \in (s_j, b] \times [0, c], \text{ here } j = 1, 2, \dots, J, \end{array} \right. \quad (3.19)$$

$$= 0.$$

Then the fractional derivative of (3.8) satisfies

$$\begin{aligned} & {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e \Upsilon(s, t) \\ &= {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e \left\{ \begin{array}{l} \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad (s, t) \in \Theta_0, \\ \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \\ + \sum_{i=1}^j [U_i(\Upsilon(s_i^-, t)) - U_i(\Upsilon(s_i^-, 0))], \quad (s, t) \in \Theta_j, j = 1, 2, \dots, J, \end{array} \right. \quad (3.20) \\ &= F(s, t, \Upsilon(s, t)) \text{ for } (s, t) \in \cup_{i=0}^J \Theta_i. \end{aligned}$$

For each s_j ($j = 1, 2, \dots, J$) in (3.8), we have

$$\begin{aligned} \Upsilon(s_j^+, t) - \Upsilon(s_j^-, t) &= \Lambda(s_j^+, t) - \Lambda(s_j^-, t) + U_j(\Upsilon(s_j^-, t)) - U_j(\Upsilon(s_j^-, 0)) \\ &= \eta(s_j^+) - \eta(s_j^-) + U_j(\Upsilon(s_j^-, t)) - U_j(\Upsilon(s_j^-, 0)) \\ &= U_j(\Upsilon(s_j^-, t)). \end{aligned} \quad (3.21)$$

Finally, letting $U_j(\Upsilon(s_j^-, t)) \rightarrow 0$ for $\forall j \in \{1, 2, \dots, J\}$ in (3.8), we obtain

$$\begin{aligned} & \lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0 \text{ for all } j \in \{1, 2, \dots, J\}} \{\text{equation (3.8)}\} \\ & \Leftrightarrow \Upsilon(s, t) = \Lambda(s, t) + \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon, \quad (s, t) \in \Theta, \\ & \Leftrightarrow \begin{cases} {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e \Upsilon(s, t) = F(s, t, \Upsilon(s, t)), & (s, t) \in \Theta, \\ \Upsilon(s, 0) = \eta(s), \Upsilon(0, t) = \xi(t), & s \in [0, b], t \in [0, c], \end{cases} \quad (3.22) \\ & = \lim_{U_j(\Upsilon(s_j^-, t)) \rightarrow 0 \text{ for all } j \in \{1, 2, \dots, J\}} \{\text{system (1.1)}\}. \end{aligned}$$

Moreover, it is obvious that (3.8) satisfies the property (3.3). Thus the proof is now accomplished. \square

4 Applications

We will offer an example to show the computation of solution of the IFrPDOS and illustrate that there are many solutions for the IFrPDOS in this section.

Example 4.1. We consider a IFRPDOS

$$\begin{cases} {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e w(s,t) = st, & (s,t) \in [0,3] \times [0,3], s \neq 1, \\ w(1^+,t) - w(1^-,t) = t, & t \in [0,3], \\ w(s,0) = w(0,t) \equiv 0, & s \in [0,3], t \in [0,3], \end{cases} \quad (4.1)$$

where $e = (e_1, e_2) = (\frac{1}{2}, \frac{1}{2})$.

Using *Theorem 3.2*, the solution of (4.1) is

$$\begin{aligned} w(s,t) = & \begin{cases} \frac{16}{9\pi} s^{\frac{3}{2}} t^{\frac{3}{2}}, & (s,t) \in [0,1] \times [0,3], \\ \frac{16}{9\pi} s^{\frac{3}{2}} t^{\frac{3}{2}} + t, & (s,t) \in (1,3] \times [0,3], \end{cases} \\ & + \begin{cases} 0, & (s,t) \in [0,1] \times [0,3], \\ \frac{8\lambda(t)}{9\pi} \left[2t^{\frac{5}{2}} + (s-1)^{\frac{1}{2}}(2s+1)t^{\frac{5}{2}} - 2s^{\frac{3}{2}}t^{\frac{5}{2}} \right] - \int_1^s \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \\ \times \left[\int_0^1 \int_0^\tau \frac{(\epsilon-u)^{-\frac{1}{2}}(\tau-v)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \frac{\partial^2}{\partial u \partial v} \left(\lambda(v) \frac{16}{9\pi} u^{\frac{3}{2}} v^{\frac{5}{2}} \right) dv du \right] d\tau d\epsilon, \\ & (s,t) \in (1,3] \times [0,3], \end{cases} \end{aligned} \quad (4.2)$$

where $\lambda(t)$ is an arbitrary continuously differentiable function of t on $[0,3]$.

Because of

$$\begin{aligned} \Phi_0(s,t) &= \int_0^s \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \epsilon \tau d\tau d\epsilon = \frac{16}{9\pi} s^{\frac{3}{2}} t^{\frac{3}{2}}; \\ \Psi_1(s,t) &= \int_0^1 \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \epsilon \tau d\tau d\epsilon + \int_1^s \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \epsilon \tau d\tau d\epsilon \\ &\quad - \int_0^s \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \epsilon \tau d\tau d\epsilon \\ &= \frac{16}{9\pi} t^{\frac{3}{2}} + \frac{8}{9\pi} (s-1)^{\frac{1}{2}}(2s+1)t^{\frac{3}{2}} - \frac{16}{9\pi} s^{\frac{3}{2}} t^{\frac{3}{2}}, \end{aligned} \quad (4.3)$$

by *Lemma 3.1*, we have

$$\begin{aligned} {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e & \begin{cases} 0, & (s,t) \in [0,1] \times [0,3], \\ \frac{8\lambda(t)[2t^{\frac{5}{2}} + (s-1)^{\frac{1}{2}}(2s+1)t^{\frac{5}{2}} - 2s^{\frac{3}{2}}t^{\frac{5}{2}}]}{9\pi} - \int_1^s \int_0^t \frac{(s-\epsilon)^{-\frac{1}{2}}(t-\tau)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \\ \times \left[\int_0^1 \int_0^\tau \frac{(\epsilon-u)^{-\frac{1}{2}}(\tau-v)^{-\frac{1}{2}}}{\Gamma(\frac{1}{2})\Gamma(\frac{1}{2})} \frac{\partial^2}{\partial u \partial v} \left(\lambda(v) \frac{16}{9\pi} u^{\frac{3}{2}} v^{\frac{5}{2}} \right) dv du \right] d\tau d\epsilon, \\ & (s,t) \in (1,3] \times [0,3], \end{cases} \\ &= 0 \end{aligned} \quad (4.4)$$

Thus, we come to that (4.2) satisfies ${}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e w(s,t) = st$ for $(s,t) \in ([0,3] \times [0,3])$ and $s \neq 1$ in (4.1). Moreover, (4.2) also satisfies $w(1^+,t) - w(1^-,t) = t$ for $t \in [0,3]$ and $w(s,0) = w(0,t) \equiv 0$ for $s \in [0,3], t \in [0,3]$ in (4.1). Because of the arbitrariness of $\lambda(t)$, (4.1) has many solutions.

5 Conclusion

It is found that there are two piecewise functions to satisfy the conditions in (1.1). And then, by constructing piecewise function and combining with the limit properties of (1.1), the integral solution of (1.1) is found that is an integral equality with an arbitrary continuously differentiable function $\lambda(t)$ to reveal the non-uniqueness for solution of (1.1) due to the arbitrariness of $\lambda(t)$.

6 Acknowledgements

The work described in this paper is financially supported by the National Natural Science Foundation of China (Grant No. 21636004, 22078030) and the Natural Science Foundation of Chongqing (No. CSTB2022NSCQ-MSX1133).

7 Appendix A

We will give the proof of *Lemma 3.1*.

Proof. Using *Definitions 2.1-2.2* and *Property 2.4*, we rewrite (3.7) into

$$\begin{aligned}
 {}^C_{(0+,0+)}\mathcal{D}_{(s,t)}^e & \begin{cases} 0, & (s,t) \in [0, s_j] \times [0, c], \\ \rho(t)\Psi_j(s,t) - \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} \\ \quad \times \left[\int_0^{s_j} \int_0^\tau \frac{(\epsilon-u)^{-e_1}(\tau-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\rho(v)\Phi_0(u,v)]}{\partial u \partial v} dv du \right] d\tau d\epsilon, \\ (s,t) \in (s_j, b] \times [0, c], \end{cases} \\
 & = \begin{cases} 0, & (s,t) \in [0, s_j] \times [0, c], \\ \int_{s_j}^s \int_0^t \frac{(s-u)^{-e_1}(t-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\rho(v)\Phi_j(u,v) - \rho(v)\Phi_0(u,v)]}{\partial u \partial v} dv du \\ \quad - \int_0^{s_j} \int_0^t \frac{(s-u)^{-e_1}(t-v)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \frac{\partial^2[\rho(v)\Phi_0(u,v)]}{\partial u \partial v} dv du, \\ (s,t) \in (s_j, b] \times [0, c], \end{cases} \tag{A.1} \\
 & = \begin{cases} 0, & (s,t) \in [0, s_j] \times [0, c], \\ {}_{(s_j^+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2[\rho(t)\Phi_j(s,t)]}{\partial s \partial t} - {}_{(0+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2[\rho(t)\Phi_0(s,t)]}{\partial s \partial t}, & (s,t) \in (s_j, b] \times [0, c]. \end{cases}
 \end{aligned}$$

To calculate ${}_{(s_j^+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2[\rho(t)\Phi_j(s,t)]}{\partial s \partial t} = {}_{(s_j^+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2}{\partial s \partial t} \left[\rho(t) \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \right]$, we transform $\int_{s_j}^s \int_0^t (s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1} F d\tau d\epsilon$ into

$$\begin{aligned}
 \int_{s_j}^s \int_0^t (s-\epsilon)^{e_1-1}(t-\tau)^{e_2-1} F d\tau d\epsilon & = \frac{1}{e_1 e_2} \int_{s_j}^s \left[\int_0^t F d(t-\tau)^{e_2} \right] d(s-\epsilon)^{e_1} \\
 & = \frac{1}{e_1 e_2} \left\{ (s-s_j)^{e_1} t^{e_2} F(s_j, 0, \Upsilon(s_j, 0)) + (s-s_j)^{e_1} \int_0^t (t-\tau)^{e_2} \frac{\partial F(s_j, \tau, \Upsilon(s_j, \tau))}{\partial \tau} d\tau \right. \\
 & \quad \left. + t^{e_2} \int_{s_j}^s (s-\epsilon)^{e_1} \frac{\partial F(\epsilon, 0, \Upsilon(\epsilon, 0))}{\partial \epsilon} d\epsilon + \int_{s_j}^s (s-\epsilon)^{e_1} \left[\int_0^t (t-\tau)^{e_2} \frac{\partial^2 F(\epsilon, \tau, \Upsilon(\epsilon, \tau))}{\partial \epsilon \partial \tau} d\tau \right] d\epsilon \right\}.
 \end{aligned}$$

Then

$$\begin{aligned}
& (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2}{\partial s \partial t} \left[\rho(t) \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1} (t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \right] \\
&= \frac{1}{\Gamma(e_1)\Gamma(e_2)} (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \left\{ (s-s_j)^{e_1-1} \left[t^{e_2-1} \rho(t) + \frac{1}{e_2} t^{e_2} \rho'(t) \right] F(s_j, 0, \Upsilon(s_j, 0)) \right. \\
&\quad + (s-s_j)^{e_1-1} \int_0^t \left[(t-\tau)^{e_2-1} \rho(t) + \frac{1}{e_2} (t-\tau)^{e_2} \rho'(t) \right] \frac{\partial F(s_j, \tau, \Upsilon(s_j, \tau))}{\partial \tau} d\tau \\
&\quad + \left[t^{e_2-1} \rho(t) + \frac{1}{e_2} t^{e_2} \rho'(t) \right] \int_{s_j}^s (s-\epsilon)^{e_1-1} \frac{\partial F(\epsilon, 0, \Upsilon(\epsilon, 0))}{\partial \epsilon} d\epsilon \\
&\quad \left. + \int_{s_j}^s (s-\epsilon)^{e_1-1} \int_0^t \left[(t-\tau)^{e_2-1} \rho(t) + \frac{1}{e_2} (t-\tau)^{e_2} \rho'(t) \right] \frac{\partial^2 F(\epsilon, \tau, \Upsilon(\epsilon, \tau))}{\partial \epsilon \partial \tau} d\tau d\epsilon \right\}. \tag{A.2}
\end{aligned}$$

We divide the right side of (A.2) into four parts (I)-(IV) as follows

$$\begin{aligned}
\text{(I)} \quad & \frac{1}{\Gamma(e_1)\Gamma(e_2)} (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \left\{ (s-s_j)^{e_1-1} \left[t^{e_2-1} \rho(t) + \frac{1}{e_2} t^{e_2} \rho'(t) \right] F(s_j, 0, \Upsilon(s_j, 0)) \right\} \\
&= \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{1-e_1-1} (\epsilon-s_j)^{e_1-1} (t-\tau)^{1-e_2-1}}{\Gamma(e_1)\Gamma(1-e_1)\Gamma(e_2)\Gamma(1-e_2)} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau d\epsilon \\
&\quad \times F(s_j, 0, \Upsilon(s_j, 0)) \quad (\text{let } \epsilon - s_j = \phi) \\
&= \int_0^{s-s_j} \frac{(s-s_j-\phi)^{1-e_1-1} \phi^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)} d\phi \int_0^t \frac{(t-\tau)^{1-e_2-1}}{\Gamma(e_2)\Gamma(1-e_2)} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau \\
&\quad \times F(s_j, 0, \Upsilon(s_j, 0)) \\
&= F(s_j, 0, \Upsilon(s_j, 0)) \int_0^t \frac{(t-\tau)^{1-e_2-1}}{\Gamma(e_2)\Gamma(1-e_2)} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau; \\
\text{(II)} \quad & (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \left\{ \frac{(s-s_j)^{e_1-1}}{\Gamma(e_1)\Gamma(e_2)} \int_0^t \left[(t-v)^{e_2-1} \rho(t) + \frac{(t-v)^{e_2}}{e_2} \rho'(t) \right] \frac{\partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} dv \right\} \\
&= \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{-e_1} (t-\tau)^{-e_2}}{\Gamma(1-e_1)\Gamma(1-e_2)} \left\{ \frac{(\epsilon-s_j)^{e_1-1}}{\Gamma(e_1)\Gamma(e_2)} \int_0^\tau \left[(\tau-v)^{e_2-1} \rho(\tau) + \frac{1}{e_2} (\tau-v)^{e_2} \rho'(\tau) \right] \right. \\
&\quad \times \left. \frac{\partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} dv \right\} d\tau d\epsilon \quad (\text{let } \epsilon - s_j = \phi) \\
&= \int_0^{s-s_j} \frac{(s-s_j-\phi)^{1-e_1-1} \phi^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)} d\phi \int_0^t \frac{(t-\tau)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \\
&\quad \times \left(\int_0^\tau \left[(\tau-v)^{e_2-1} \rho(\tau) + \frac{1}{e_2} (\tau-v)^{e_2} \rho'(\tau) \right] \frac{\partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} dv \right) d\tau \\
&= \int_0^t \int_v^t \frac{(t-\tau)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \left[(\tau-v)^{e_2-1} \rho(\tau) + \frac{1}{e_2} (\tau-v)^{e_2} \rho'(\tau) \right] \frac{\partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} d\tau dv \\
&\quad (\text{let } \tau - v = \varphi) \\
&= \int_0^t \int_0^{t-v} \frac{(t-v-\varphi)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \left[\varphi^{e_2-1} \rho(\varphi+v) + \frac{1}{e_2} \varphi^{e_2} \rho'(\varphi+v) \right] \frac{\partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} d\varphi dv;
\end{aligned}$$

$$\begin{aligned}
\text{(III)} \quad & (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \left\{ \left[t^{e_2-1} \rho(t) + \frac{1}{e_2} t^{e_2} \rho'(t) \right] \int_{s_j}^s \frac{(s-u)^{e_1-1}}{\Gamma(e_1)\Gamma(e_2)} \frac{\partial F(u, 0, \Upsilon(u, 0))}{\partial u} du \right\} \\
&= \int_{s_j}^s \frac{\partial F(u, 0, \Upsilon(u, 0))}{\partial u} \int_u^s \frac{(s-\epsilon)^{-e_1} (\epsilon-u)^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)\Gamma(e_2)\Gamma(1-e_2)} d\epsilon du \\
&\quad \times \int_0^t (t-\tau)^{-e_2} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau \quad (\text{let } \epsilon - u = \phi) \\
&= \int_{s_j}^s \frac{\partial F(u, 0, \Upsilon(u, 0))}{\partial u} \left[\int_0^{s-u} \frac{(s-u-\phi)^{-e_1} \phi^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)} d\phi \right] du \\
&\quad \times \int_0^t \frac{(t-\tau)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau \\
&= \frac{F(s, 0, \Upsilon(s, 0)) - F(s_j, 0, \Upsilon(s_j, 0))}{\Gamma(e_2)\Gamma(1-e_2)} \int_0^t (t-\tau)^{-e_2} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau; \\
\text{(IV)} \quad & (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \left\{ \int_{s_j}^s \frac{(s-u)^{e_1-1}}{\Gamma(e_1)\Gamma(e_2)} \int_0^t \left[(t-v)^{e_2-1} \rho(t) + \frac{(t-v)^{e_2}}{e_2} \rho'(t) \right] \frac{\partial^2 F(u, v, \Upsilon(u, v))}{\partial u \partial v} dv du \right\} \\
&= \int_{s_j}^s \frac{(s-\epsilon)^{-e_1}}{\Gamma(e_1)\Gamma(1-e_1)} \int_{s_j}^\epsilon (\epsilon-u)^{e_1-1} \left\{ \int_0^t \frac{\partial^2 F(u, v, \Upsilon(u, v))}{\partial u \partial v} \int_v^t \frac{(t-\tau)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \right. \\
&\quad \times \left. \left[(\tau-v)^{e_2-1} \rho(\tau) + \frac{1}{e_2} (\tau-v)^{e_2} \rho'(\tau) \right] d\tau dv \right\} d\epsilon du \quad (\text{let } \tau - v = \varphi) \\
&= \int_{s_j}^s \int_u^s \frac{(s-\epsilon)^{-e_1} (\epsilon-u)^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)} \left\{ \int_0^t \frac{\partial^2 F(u, v, \Upsilon(u, v))}{\partial u \partial v} \int_0^{t-v} \frac{(t-v-\varphi)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \right. \\
&\quad \times \left. \left[\varphi^{e_2-1} \rho(\varphi+v) + \frac{1}{e_2} \varphi^{e_2} \rho'(\varphi+v) \right] d\varphi dv \right\} d\epsilon du \quad (\text{let } \epsilon - u = \phi) \\
&= \int_{s_j}^s \int_0^{s-u} \frac{(s-u-\phi)^{-e_1} \phi^{e_1-1}}{\Gamma(e_1)\Gamma(1-e_1)} \left\{ \int_0^t \frac{\partial^2 F(u, v, \Upsilon(u, v))}{\partial u \partial v} \int_0^{t-v} \frac{(t-v-\varphi)^{-e_2}}{\Gamma(e_2)\Gamma(1-e_2)} \right. \\
&\quad \times \left. \left[\varphi^{e_2-1} \rho(\varphi+v) + \frac{1}{e_2} \varphi^{e_2} \rho'(\varphi+v) \right] d\varphi dv \right\} d\phi du \\
&= \int_0^t \int_0^{t-v} \frac{(t-v-\varphi)^{-e_2} \left[\varphi^{e_2-1} \rho(\varphi+v) + \frac{1}{e_2} \varphi^{e_2} \rho'(\varphi+v) \right]}{\Gamma(e_2)\Gamma(1-e_2)} \\
&\quad \times \frac{\partial F(s, v, \Upsilon(s, v)) - \partial F(s_j, v, \Upsilon(s_j, v))}{\partial v} d\varphi dv.
\end{aligned}$$

Substituting (I)-(IV) into (A.2), we have

$$\begin{aligned}
& (s_j^+, 0+) \mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2}{\partial s \partial t} \left[\rho(t) \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1} (t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \right] \\
&= \int_0^t \int_0^{t-v} \frac{(t-v-\varphi)^{-e_2} \left[\varphi^{e_2-1} \rho(\varphi+v) + \frac{1}{e_2} \varphi^{e_2} \rho'(\varphi+v) \right]}{\Gamma(e_2)\Gamma(1-e_2)} \frac{\partial F(s, v, \Upsilon(s, v))}{\partial v} d\varphi dv \\
&\quad + \frac{F(s, 0, \Upsilon(s, 0))}{\Gamma(e_2)\Gamma(1-e_2)} \int_0^t (t-\tau)^{-e_2} \left[\tau^{e_2-1} \rho(\tau) + \frac{1}{e_2} \tau^{e_2} \rho'(\tau) \right] d\tau.
\end{aligned} \tag{A.3}$$

In addition, letting $s_j = 0$ in (A.3), we obtain

$$\begin{aligned} & {}_{(0+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2}{\partial s \partial t} \left[\rho(t) \int_0^s \int_0^t \frac{(s-\epsilon)^{e_1-1} (t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \right] \\ &= {}_{(s_j^+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2}{\partial s \partial t} \left[\rho(t) \int_{s_j}^s \int_0^t \frac{(s-\epsilon)^{e_1-1} (t-\tau)^{e_2-1}}{\Gamma(e_1)\Gamma(e_2)} F d\tau d\epsilon \right] \end{aligned} \quad (\text{A.4})$$

then

$${}_{(s_j^+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2 [\rho(t)\Phi_j(s,t)]}{\partial s \partial t} - {}_{(0+,0+)}\mathcal{I}_{(s,t)}^{1-e} \frac{\partial^2 [\rho(t)\Phi_0(s,t)]}{\partial s \partial t} = 0. \quad (\text{A.5})$$

This proof is completed. \square

References

- [1] J.W. Zhou, Y.M. Deng, Y.N. Wang, Variational approach to p-Laplacian fractional differential equations with instantaneous and non-instantaneous impulses, *Applied Mathematics Letters*, 104, 106251 (2020)
- [2] J.Y. Cao, L.Z. Chen, Z.Q. Wang, A block-by-block method for the impulsive fractional ordinary differential equations, *Journal of Applied analysis and Computation*, 10, 3, 853-874 (2020)
- [3] H.D. Gou, Y.X. Li, A study on impulsive Hilfer fractional evolution equations with nonlocal conditions, *International Journal of Nonlinear Sciences and Numerical Simulation*, 21, 2, 205-218 (2020)
- [4] H.D. Gou, Y.X. Li, The method of lower and upper solutions for impulsive fractional evolution equations in Banach spaces, *Journal of the Korean Mathematical Society*, 57, 1, 61-88 (2020)
- [5] K.D. Kucche, J.P. Kharade, J.V.D.C. Sousa, On the nonlinear impulsive Ψ -Hilfer fractional differential equations, *Mathematical Modelling and Analysis*, 25, 4, 642-660 (2020)
- [6] S. Heidarkhani, A Salari, Nontrivial solutions for impulsive fractional differential systems through variational methods, *Mathematical Methods in the Applied Sciences*, 43, 10, 6529-6541 (2020)
- [7] H.D. Gou, Y.X. Li, The method of lower and upper solutions for impulsive fractional evolution equations, *Annals of Functional Analysis*, 11, 2, 350-369 (2020)
- [8] J. You, S.R. Sun, On impulsive coupled hybrid fractional differential systems in Banach algebras, *Journal of applied Mathematics and Computing*, 62, 1-2, 189-205 (2020)
- [9] H.D. Gou, Y.X. Li, A study on impulsive fractional hybrid evolution equations using sequence method, *Computational and Applied Mathematics*, 39, 3, 225 (2020)
- [10] C. Ravichandran, K. Logeswari, S.K. Panda, K.S. Nisar, On new approach of fractional derivative by Mittag-Leffler kernel to neutral integro-differential systems with impulsive conditions, *Chaos Solitons and Fractals*, 139, 110012 (2020)

- [11] D.D. Min, F.Q. Chen, Existence of solutions for a fractional advection-dispersion equation with impulsive effects via variational approach, *Journal of Applied Analysis and Computation*, 10, 3, 1005-1023 (2020)
- [12] L.M. Feng, Z.L. Han, Oscillation behavior of solution of impulsive fractional differential equation, *Journal of Applied analysis and Computation*, 10, 1, 223-233 (2020)
- [13] L.M. Feng, Y.B. Sun, Z.L. Han, Philos-type oscillation criteria for impulsive fractional differential equations, *Journal of applied Mathematics and Computing*, 62, 1-2, 361-376 (2020)
- [14] T.W. Zhang, L.L. Xiong, Periodic motion for impulsive fractional functional differential equations with piecewise Caputo derivative, *Applied Mathematics Letters*, 2020, 101 (2020)
- [15] M.R. Xu, S.R. Sun, Z.L. Han, Solvability for impulsive fractional Langevin equation, *Journal of Applied analysis and Computation*, 10, 2, 486-494 (2020)
- [16] L.J. Cheng, L.Y. Hu, Y. Ren, Perturbed impulsive neutral stochastic functional differential equations, *Qualitative Theory of Dynamical Systems*, 20, 2, 27 (2021)
- [17] J.K. Liu, W. Xu, An averaging result for impulsive fractional neutral stochastic differential equations, *Applied Mathematics Letters*, 114, 106892 (2021)
- [18] K.D. Kucche, J.P. Kharade, Analysis of impulsive ϕ -Hilfer fractional differential equations, *Mediterranean Journal of Mathematics*, 17, 5, 163 (2020)
- [19] R. Agarwal, S. Hristova, D. O'Regan, A survey of Lyapunov functions, stability and impulsive Caputo fractional differential equations, *Fractional Calculus and Applied Analysis*, 19, 290-318 (2016)
- [20] G. Wang, B. Ahmad, L. Zhang, J.J. Nieto, Comments on the concept of existence of solution for impulsive fractional differential equations, *Communications in Nonlinear Science and Numerical Simulation*, 19, 3, 401-403 (2014)
- [21] M. Feckan, Y. Zhou, J.R. Wang, On the concept and existence of solution for impulsive fractional differential equations, *Communications in Nonlinear Science and Numerical Simulation*, 17, 7, 3050-3060 (2012)
- [22] J.R. Wang, X. Li, W. Wei, On the natural solution of an impulsive fractional differential equation of order $q \in (1, 2)$, *Communications in Nonlinear Science and Numerical Simulation*, 17, 11, 4384-4394 (2012)
- [23] M. Feckan, Y. Zhou, J.R. Wang, Response to "Comments on the concept of existence of solution for impulsive fractional differential equations [Commun Nonlinear Sci Numer Simul 2014;19:401-3.]", *Communications in Nonlinear Science and Numerical Simulation*, 19,12, 4213-4215 (2014)
- [24] J.R. Wang, M. Fekan, Y. Zhou, A survey on impulsive fractional differential equations, *Fractional Calculus and Applied Analysis*, 19, 806-831 (2016)

- [25] Y. Liu, On piecewise continuous solutions of higher order impulsive fractional differential equations and applications, *Applied Mathematics and Computation*, 287-288, 38-49 (2016)
- [26] Y. Liu, Survey and new results on boundary-value problems of singular fractional differential equations with impulse effects, *Electronic Journal of Differential Equations*, 2016, 296, 1-177 (2016)
- [27] Y. Liu, A new method for converting boundary value problems for impulsive fractional differential equations to integral equations and its applications, *Advances in Nonlinear Analysis*, 8, 1, 386-454 (2019)
- [28] X. Zhang, Z. Liu, S. Yang, Z. Peng, Y. He, L. Wei, The right equivalent integral equation of impulsive Caputo fractional-order system of order $\epsilon \in (1, 2)$, *Fractal Fract.*, 2023, 7, 37 (2023)
- [29] S. Abbas, M. Benchohra, Upper and lower solutions method for impulsive partial hyperbolic differential equations with fractional order, *Nonlinear Analysis: Hybrid Systems*, 4, 3, 406-413 (2010)
- [30] S. Abbas, M. Benchohra, Impulsive partial hyperbolic functional differential equations of fractional order with state-dependent delay, *Fractional Calculus and Applied Analysis*, 13, 225-242 (2010)
- [31] S. Abbas, M. Benchohra, Lech Gorniewicz, Existence theory for impulsive partial hyperbolic functional differential equations involving the Caputo fractional derivative, *Scientiae Mathematicae Japonicae*, e-2010, 271-282 (2010)
- [32] S. Abbas, Ravi P. Agarwal, M. Benchohra, Darboux problem for impulsive partial hyperbolic differential equations of fractional order with variable times and infinite delay, *Nonlinear Analysis: Hybrid Systems*, 4, 818-829 (2010)
- [33] T.L. Guo, K.J. Zhang, Impulsive fractional partial differential equations, *Applied Mathematics and Computation*, 257, 581-590 (2015)
- [34] X.M. Zhang, Non-uniqueness of solution for initial value problem of impulsive fractional partial differential equations, *International Journal of Dynamical Systems and Differential Equations*, 12, 4, 316-338 (2022)
- [35] S.G. Samko, A.A. Kilbas and O.N. Marichev, Integrals and derivatives of fractional order and their applications [in Russian], *Tekhnika*, Minsk (1987)
- [36] A.N. Vityuk, A.V. Golushkov, Existence of solutions of systems of partial differential equations of fractional order, *Nonlinear Oscillations*, 7, 3, 318-325 (2004)