

Peng et al.(2014)
Wegner et al.(2013)
Wegner et al.(2013)
Peng et al.(2014)
Wang et al.(2014)
Sugiyama et al. (2015)
Rodrigues et al.(2017)
Chen et al. (2019)
Lin et al.(2019)
Park et al.(2012)
Lin et al.(2019)
Antunes et al.(2017)
Sugiyama et al. (2015)
Northam et al.(2005)
Sugiyama et al. (2015)
Northam et al.(2005)
Sugiyama et al. (2015)
Northam et al.(2005)
Park et al.(2012)
Wang et al.(2014)
Rodrigues et al.(2017)
Wang et al.(2014)
Wang et al.(2014)
Peng et al.(2014)
Lin et al.(2019)
Yu et al.(2016)

Sex (M/F)
Age (18-75) [years]

Weight

Co-administration with inducer (Y/N)

Co-administration with VPA (Y/N)

Co-administration with LEV or LTG (Y/N)

Dose (75-1350) [mg/d]

eGFR (20-120) [ml/min]

ALT (50-149) [U/L]

Children (16-40) [kg]

Both children and adults (16-100) [kg]

Adults (40-100) [kg]

0 0.5 0.8 1 1.25 1.5 2
Covariate effect on CL