

Identification of novel arsenic resistance genes in yeast

Esin Isik¹, Çiğdem Balkan¹, Vivien Karl², Hüseyin Çağlar Karakaya¹, Sansan Hua², Sebastien Rauch³, Markus J. Tamás^{2*} and Ahmet Koc^{1,4*}

¹Department of Molecular Biology and Genetics, Izmir Institute of Technology, Izmir, Turkey

²Department of Chemistry and Molecular Biology, University of Gothenburg, SE-405 30 Gothenburg, Sweden

³Water Environment Technology, Department of Architecture and Civil Engineering, Chalmers University of Technology, Gothenburg, Sweden

⁴Department of Genetics, School of Medicine, Inonu University, 44280 Malatya, Turkey

*Corresponding authors:

Ahmet Koc, Department of Genetics, School of Medicine, Inonu University, 44280 Malatya, Turkey. Email: ahmet.koc@inonu.edu.tr

Markus J. Tamás, Department of Chemistry and Molecular Biology, University of Gothenburg, SE-405 30 Gothenburg, Sweden. Email: markus.tamas@cmb.gu.se

ABSTRACT

Arsenic is a toxic metalloid that affects human health by causing numerous diseases and by being used in the treatment of acute promyelocytic leukemia. *Saccharomyces cerevisiae* (budding yeast) has been extensively utilized to elucidate the molecular mechanisms underlying arsenic toxicity and resistance in eukaryotes. In this study, we applied a genomic DNA overexpression strategy to identify yeast genes that provide arsenic resistance in wild-type and arsenic-sensitive *S. cerevisiae* cells. In addition to known arsenic-related genes, our genetic screen revealed novel genes, including *PHO86*, *VBA3*, *UGP1*, and *TUL1*, whose overexpression conferred resistance. To gain insights into possible resistance mechanisms, we addressed the contribution of these genes to cell growth, intracellular arsenic, and protein aggregation during arsenate exposure. Overexpression of *PHO86* resulted in higher cellular arsenic levels but no additional effect on protein aggregation, indicating that these cells efficiently protect their intracellular environment. *VBA3* overexpression caused resistance despite higher intracellular arsenic and protein aggregation levels. Overexpression of *UGP1* led to lower intracellular arsenic and protein aggregation levels whilst *TUL1* overexpression had no impact on intracellular arsenic or protein aggregation levels. Thus, the identified genes appear to confer arsenic resistance through distinct mechanisms but the molecular details remain to be elucidated.

Keywords: arsenate, arsenite, yeast, Pho86, Vba3, Tul1, Ugp1, arsenic transport

INTRODUCTION

The metalloid arsenic is abundantly present in the environment mainly as pentavalent arsenate [As(V)] and trivalent arsenite [As(III)]. Both forms of arsenic are toxic and chronic exposure causes cancers of the skin, lung, bladder, kidney, and liver, and is associated with various neurodegenerative disorders. Exposure occurs mainly via toxic waste or through the consumption of contaminated drinking water and food. Chronic arsenic exposure is a significant public health concern with hundreds of millions of persons worldwide estimated to be affected (Clemens & Ma, 2016; Naujokas et al, 2013).

On the cellular level, arsenic's toxic and carcinogenic effects may be caused by the inactivation of specific enzymes, induction of oxidative stress, inhibition of DNA repair systems, and deregulation of cell proliferation. As(V) resembles phosphate and may replace phosphate in some reactions and interfere with ATP synthesis. As(III) has a high affinity for sulfhydryl groups and can bind to cysteine residues in proteins, thereby altering their folding and activity. Additionally, As(III) may cause oxidative damage to proteins, lipids, and DNA (Hughes et al, 2011; Shen et al, 2013; Tamás et al, 2014). Despite its toxicity, arsenic has a long history of usage as a therapeutic agent and is currently used to treat acute promyelocytic leukemia. The disease-causing and therapeutic effects of arsenicals have spurred research into the molecular mechanisms that underlie their toxicity and the mechanisms that cells use to develop resistance. Most of these studies have focused on model organisms such as bacteria and the eukaryote model *Saccharomyces cerevisiae* (budding yeast) (Garbinski et al, 2019; Rosen & Tamás, 2010; Slyemi & Bonnefoy, 2012; Wysocki & Tamás, 2010).

Arsenic exerts its toxic effects inside cells. Therefore, transport proteins that allow the influx of arsenic or mediate its export are central players of toxicity and resistance. In budding yeast, like in most cells, As(V) is taken up by phosphate transporters (Pho84 in *S. cerevisiae*) (Bun-ya et al, 1996; Yompakdee et al, 1996), while As(III) is taken up by aquaglyceroporins (Fps1) (Wysocki et al, 2001) and hexose transporters (Liu et al, 2004). Intracellular As(V) is reduced to As(III) by the arsenate reductase Acr2 (also called Arr2) (Mukhopadhyay & Rosen, 1998; Mukhopadhyay et al, 2000) followed by As(III) export from the cytosol through the plasma membrane arsenite efflux protein Acr3 (also called Arr3) (Wysocki et al, 1997) or sequestration into the vacuole in form of the glutathione conjugate As(GS)₃ catalyzed by the ABC (ATP-binding cassette) transporter Ycf1 (Ghosh et al, 1999). These transport proteins are regulated at the transcriptional and post-translational levels during As(III) stress. For instance, expression of *ACR2* (*ARR2*) and *ACR3* (*ARR3*) is regulated by the arsenic-sensing transcription

factor Yap8 (also called Acr1 or Arr1) (Kumar et al, 2016; Menezes et al, 2004; Wysocki et al, 2004), *YCF1* expression is regulated by Yap1 whilst expression of glutathione biosynthesis-related genes involves the transcription factors Yap1 and Met4 (Thorsen et al, 2007; Wysocki et al, 2004). The activity of the yeast aquaglyceroporin Fps1 is regulated by the kinases Hog1 and Slt2 (Ahmadpour et al, 2016; Lee & Levin, 2018; Thorsen et al, 2006) as well as the Rgc1 and Rgc2 proteins (Beese et al, 2009) whereas hexose transporters are degraded in response to As(III), probably as a way to gain resistance (Jochem et al, 2019). Another cellular defense strategy employed by most cells is arsenic chelation. *S. cerevisiae* increases the production of the tri-peptide glutathione in response to As(III) exposure for intracellular and extracellular chelation and detoxification (Talemi et al, 2014; Thorsen et al, 2012; Thorsen et al, 2007). Since arsenic induces widespread misfolding and aggregation of newly synthesized proteins (Ibstedt et al, 2014; Jacobson et al, 2012), decreased protein synthesis and proteasomal degradation of misfolded/aggregated proteins represent additional mechanisms used by cells to evade arsenic toxicity (Andersson et al, 2021; Guerra-Moreno et al, 2015; Jacobson et al, 2012). A number of other genes and pathways with a putative role in resistance or toxicity have been identified in comprehensive genome-wide phenotypic screens using the yeast genome-wide deletion collection (Haugen et al, 2004; Jin et al, 2008; Jo et al, 2009; Johnson et al, 2016; Pan et al, 2010; Thorsen et al, 2009; Zhou et al, 2009) or via transcriptomic or proteomic approaches (da Silva et al, 2018; Haugen et al, 2004; Hosiner et al, 2009; Jin et al, 2008; Jochem et al, 2019; Matia-Gonzalez & Rodriguez-Gabriel, 2011; Thorsen et al, 2009). In this current study, we performed a high-copy genomic DNA (gDNA) library screening in yeast and identified genes whose overexpression confers arsenic resistance.

MATERIALS AND METHODS

Yeast strains, growth conditions, plasmids, and gDNA library screening

The *S. cerevisiae* strains used in this study are listed in Table A1. Deletion mutants were purchased from EUROSCARF and the *HSP104*-GFP strain was purchased from Invitrogen. Haploid and diploid deletion mutants are respectively in BY4741 and BY4743 backgrounds lacking the corresponding gene but carrying the kanamycin resistance gene instead. YPAD (1% yeast extract, 2% peptone, 0.04 % adenine, 2% glucose) was used as a rich medium and SD (synthetic dextrose with 0.67% YNB (yeast nitrogen base) without amino acids, 2% glucose

and supplemented with auxotrophic requirements) was used as a selective medium for regular cultivation of yeast cells. Sodium arsenate, Na_2HAsO_4 (Cat. No: A6756), and sodium arsenite, NaAsO_2 (Cat. No: S7400), were purchased from Sigma-Aldrich.

High copy yeast genomic DNA library (AB320 genomic DNA library in YEp13 plasmid in *Escherichia coli* obtained from the American Type Culture Collection ATCC (No: 37323)) was introduced into yeast cells by transformation for subsequent genomic library screening. The transformation was carried out by the standard LiAc method.

The set-up of the screen and the arsenic concentrations used are illustrated in Fig. 1a. The plasmids present in the resistant colonies were isolated by a commercial miniprep kit (GeneJET Plasmid Miniprep Kit, Thermo-Molecular Biology) after treating the yeast cells with lyticase (5u/ml) for 30 min. The isolated plasmids were amplified in *JM109 E. coli* cells and sequenced with vector-specific sequencing primers using DNA sequencer ABI3130xl with ABI PRISM sequencing analysis v5.1 program at IzTech Biotechnology Center (Izmir, Turkey).

Cloning of selected candidate genes

Selected candidate genes were cloned from the yeast Open Reading Frame (ORF) collection (in *E.coli*, YSC3868) into the pAG425GPD-ccdB plasmid by the Gateway Cloning System. Since the *PHO86* gene was not available in the ORF collection, it was amplified by PCR using a pair of Gateway-compatible primers and then cloned into pAG425GPD-ccdB. The correct cloning of the candidate genes was confirmed by restriction enzyme analysis and sequencing with vector-specific primers.

Growth assay on gradient plates

A gradient plate was prepared with two different layers of agar media to produce a continuous concentration gradient of the metalloid in a Petri dish, as previously reported (Szybalski & Bryson, 1952). Initially, 50 ml of agar medium without metalloid was poured as a lower layer onto a square Petri dish (120 mm x 120 mm) and then tilted at an angle to cover the complete bottom surface of the plate. After solidification, the plate was positioned horizontally and 50 ml of medium with the corresponding arsenical at the desired concentration was poured on top of the solid medium. The arsenicals in the upper layer diffused towards the lower layer in proportion to the heights of the layers and, because of this diffusion, a uniform concentration

gradient was established increasing from 0 mM on one side to the highest arsenic concentration on the other side. An overnight culture of yeast cells in the corresponding growth medium was diluted and grown to the logarithmic phase. Then, 5 μl of the cells with an OD_{600} value of 0.02 were spotted onto the gradient plates starting from one end of the plate to the other end. The plates were incubated at 30°C for 3-5 days and the growth of the cells was recorded.

Arsenic measurements

Plasmids overexpressing the indicated genes were introduced into yeast cells (BY4741). The transformants were grown to exponential phase in a YNB medium containing 2% glucose and subsequently, cells were either left untreated or exposed to 1 mM As(V) for 1 h. Samples were collected before and after As(V) exposure and washed twice in ice-cold water. The cell pellet was then resuspended in water, boiled for 10 min, and centrifuged to collect the supernatant. To measure arsenic efflux, cells were first incubated with 1 mM As(V) for 1 h to allow arsenic accumulation, then the cells were washed with As(V)-free medium and samples taken for intracellular arsenic determination. The arsenic content of each sample was determined by inductively coupled plasma-mass spectrometry (ICP-MS) following a 20-fold dilution with ultrapure water from a GenPure water purification system (Thermo Scientific Barnstead, resistivity 18.2 M Ω cm) and acidification to 1% volume HNO₃ (Sharlau, HNO₃, 65% for trace analysis) before the analysis. The analysis was performed on an ICAP Q ICP-MS (Thermo Fisher Scientific) with an SC-FAST automated sample introduction system (Elemental Scientific). Arsenic was measured at $m/z=75$ and the instrument was operated in the Kinetic Energy Discrimination (KED) mode with He as the collision gas to remove potential interference from ArCl⁺ at $m/z=75$. Calibration was performed using a set of arsenic standards with concentrations up to 500 $\mu\text{g L}^{-1}$. A solution of 1 $\mu\text{g L}^{-1}$ indium was continuously injected for internal standardization. The detection limit is estimated to be 0.1 $\mu\text{g As L}^{-1}$.

Protein aggregation assay

Plasmids were introduced into yeast cells (BY4741) harboring a genomic copy of the Hsp104-GFP fusion protein (Huh et al, 2003). The transformants were grown to mid-log phase in YNB medium and exposed to 2 mM As(V). Cells were collected at the indicated time points, fixed with formaldehyde for 30 min at room temperature, and washed with phosphate-buffered saline (PBS). The GFP signals were observed using a Zeiss Axiovert 200M (Carl Zeiss MicroImaging)

fluorescence microscope equipped with Plan-Apochromat 1.40 objectives and appropriate fluorescence light filter sets. Images were taken with a digital camera (AxioCamMR3) and processed with Zeiss Zen software. To quantify protein aggregation, the fraction of cells with aggregates/Hsp104–GFP foci was determined by visual inspection of 110–230 cells per strain and time point using Image J software.

Localization of Pho84-GFP and Pho86-GFP

The plasmids *pPHO84-GFP* (EB0666) and *pPHO86-GFP* (EB0667) (Lau et al, 2000) were introduced into BY4741 cells. Overnight cultures of the transformants were diluted in fresh liquid media and then treated with 0.5 mM As(V) for 1 h. Cells were transferred to microscope slides and analyzed with a fluorescent microscope.

RESULTS

High-copy genomic library screening for genes conferring resistance to arsenicals

To identify novel arsenic resistance genes, we screened a high-copy gDNA library in two different yeast backgrounds: W303-1A (wild-type) and the arsenic sensitive GK4 (Δ *acr1* Δ *acr2* Δ *acr3*) strains (Fig. 1a; Appendix Table A1). Initially, we performed growth assays on plates to assess minimum inhibitory concentration (MIC) values of As(III) and As(V). 1.5 mM and 0.5 mM of As(V) completely inhibited growth of W303-1A and GK4, respectively, while the MIC of As(III) was determined to be 1.0 mM and 0.1 mM for W303-1A and GK4 cells, respectively (Appendix Figures. A1, A2). After determining the MICs, the yeast genome library was inserted into the W303-1A and GK4 strains, and the transformants were screened on plates containing As(V) or As(III) at lethal concentrations (>MIC) (Fig. 1a). Surviving colonies were selected for further experiments.

We first confirmed the arsenic resistance of the surviving colonies from the library screen by growth assay on gradient plates. Subsequently, plasmids present in the resistant colonies were isolated and amplified in *E. coli*. The amplified plasmids were inserted back into the corresponding yeast strain. Growth assays were carried out to confirm that the resistance observed in the original transformants was due to the presence of the plasmids. Consequently,

ten colonies for As(V) (9 for W303-1A and 1 for GK4) and four colonies for As(III) (GK4) resistance were selected (Figs. 1b, 1c, A3). The plasmids from these colonies were sequenced with vector-specific primers. Subsequent bioinformatics analysis revealed the genomic regions present in these plasmids (Table 1). The genomic library used in this study was constructed by ligation of the fragments generated by *Sau3AI* partial digestion (Nasmyth & Reed, 1980); thus, most of the cassettes contained multiple genes, except for one colony (AsIII.3, Table 1).

Genes whose overexpression confers arsenate resistance

Ten different As(V) resistant colonies were obtained: eight of those contained genes from the same genomic fragment encompassing the genes *ASF1*, *NCA3*, *PHO86*, *YJL118W*, *RPE1*, *ALB1*, *MTC1*, *LSMI*, whilst two of the colonies contained distinct expression cassettes with the genes *ARR1*, *ARR2*, *ARR3*, *SGE1*, *YPR196W* and *TUL1*, *UGP1*, *AIM26*, respectively (Table 1). The presence of the known arsenic resistance gene cluster, consisting of *ARR1/ACR1/YAP8*, *ARR2/ACR2*, and *ARR3/ACR3* (Bobrowicz et al, 1997), in one of the colonies (AsV.3) validates the gDNA library screening approach. The other genes in that cassette (*SGE1*, *YPR196W*) have no known role in arsenic resistance. Likewise, none of the genes in cassette AsV.4 (*AIM26*, *UGP1*, *TUL1*) have previously been associated with arsenic resistance. *AIM26* encodes a protein of unknown function, *UGP1* encodes a UDP-glucose pyrophosphorylase, and *TUL1* encodes a Golgi-localized RING-finger ubiquitin ligase. The remaining eight cassettes comprised overlapping gene fragments from Chromosome X with *PHO86* and *YJL118W* genes present in all cassettes. Pho86 is an endoplasmic reticulum (ER)-resident protein that is required for ER exit of the inorganic phosphate transporter Pho84 (Lau et al, 2000). Both *PHO86* and *PHO84* have been implicated in As(V) resistance; Pho84 is responsible for the internalization of arsenate into yeast cells (Shen et al, 2012), and deletion of either *PHO84* or *PHO86* increases As(V) resistance (Bun-ya et al, 1996). *YJL118W* encodes a protein of unknown function with no known role in arsenic resistance. Deletions of some of the other genes in these cassettes were found in genome-wide screens to cause As(III) sensitivity, including in *ASF1*, involved in nucleosome assembly and DNA repair (Pan et al, 2010), and *RPE1* encoding a D-ribulose-5-phosphate 3-epimerase (Haugen et al, 2004; Thorsen et al, 2009). The genes *NCA3*, *ALB1*, *MTC1*, and *LSMI* have not been implicated in arsenic resistance previously.

Although we performed the gDNA library screening in both W303-1A and GK4 cells, we obtained As(III) resistant colonies only in the GK4 background. Out of the four colonies, one contained the known arsenic resistance genes *ARR1/ACR1/YAP8*, *ARR2/ACR2*, and

ARR3/ACR3 (Bobrowicz et al, 1997) validating our approach. Two colonies had identical cassettes containing the *GIS4*, *YAPI*, *ERG6*, and *MRPL39* genes. Among these, deletion of *YAPI* and *ERG6* sensitizes cells to As(III) (Haugen et al, 2004; Pan et al, 2010; Thorsen et al, 2009; Wysocki et al, 2004), and *YAPI* overexpression improves arsenic resistance in cells lacking the vacuolar ABC transporter Ycf1 (Bouganim et al, 2001). *GIS4* and *MRPL39* have not been implicated in arsenic resistance previously. One colony contained a single gene in the cassette, *VBA3*, encoding a vacuolar basic amino acid transporter with no previous function in arsenic resistance.

Cloning and overexpression of selected genes

We next cloned and expressed selected genes from a high-copy expression plasmid (pAG425GPD) in W303-1A and GK4 cells. The growth of yeast cells overexpressing these candidate genes was then tested in the presence of As(V) and As(III). The different cassettes with *PHO86* and *YJL118W* contained eight genes in total. Overexpression of the *PHO86* gene provided resistance to As(V) in both wild-type W303-1A and arsenic-sensitive GK4 cells (Figs. 2a, 2b) but did not improve As(III) resistance, indicating an As(V)-specific role for Pho86. In contrast, individual overexpression of *YJL118W*, *NCA3*, *RPE1*, *ALB1*, or *LSM1* did not confer resistance to As(III) or As(V) (Fig. A4). Cells overexpressing *MTC1* or *ASF1* grew poorly both in the absence and presence of As(V) (Fig. A4), suggesting that these genes have a negative impact on growth and resistance. Of the three genes in cassette AsV.4, individual overexpression of *TUL1* and *UGP1* improved As(V) resistance of W303-1A cells but not of arsenic-sensitive GK4 cells whilst overexpression of *AIM26* did not improve growth in either strain background (Figs 2a, 2b). Thus, of the three genes in colony AsV.4, both *TUL1* and *UGP1* contributed to the observed resistance whereas *AIM26* did not. Overexpression of the tested As(III) resistance genes (*VBA3*, *YAPI*, *ERG6*) improved As(III) resistance of GK4 cells whilst in W303-1A cells, only *VBA3* overexpression conferred As(III) resistance (Figs. 2c, 2d). *VBA3* overexpression also resulted in As(V) resistance in the GK4 background. For our follow-up experiments, we decided to focus on selected genes and their roles in As(V) resistance in a strain background (BY4741) that could be used for all further experiments. We confirmed the As(V) resistant phenotype of wild-type cells of the BY4741 background overexpressing *PHO86*, *VBA3*, *TUL1*, and *UGP1* (Fig. 2e). Although all four genes conferred As(V) resistance upon overexpression, we noted that their potency was different in the BY4741 background compared to W303-1A. W303-1A is generally more sensitive to various stress conditions than

BY4741, and this is also the case for As(III) and As(V) (see Appendix Figures A1, A2). Thus, the degree of the benefit provided by overexpression of individual genes appears to depend on the strain background.

Intracellular arsenic levels

Since arsenic exerts its toxic effects through intracellular targets, we reasoned that some of the identified genes might affect arsenic uptake, efflux, or intracellular sequestration pathways. To address this, we exposed BY4741 cells to 1 mM As(V) for 1 h and determined intracellular arsenic levels (Fig. 3a). There was a significant increase in intracellular arsenic in cells overexpressing *PHO86* ($p=0.01$) and *VBA3* ($p=0.001$) compared to cells carrying the empty plasmid (p -only). Thus, *PHO86* and *VBA3* overexpressing cells are resistant despite elevated intracellular arsenic levels. Intracellular arsenic levels were unchanged in cells overexpressing *TUL1* ($p=0.97$) and slightly lower in cells overexpressing *UGPI* (borderline of statistical significance with $p=0.08$). Thus, enhanced resistance of *UGPI* overexpressing cells might be related to less intracellular arsenic.

Chelation and vacuolar sequestration are major arsenic detoxification pathways in yeast (Rosen & Tamás, 2010; Wysocki & Tamás, 2010) and we have previously shown that cells with efficient vacuolar sequestration capacity retain more intracellular arsenic upon removal of extracellular arsenic (Talemi et al, 2014). We therefore asked whether intracellular arsenic in *PHO86* and *VBA3* overexpressing cells is retained or whether it can leave cells upon removal of extracellular As(V). For this, we first incubated cells with 1 mM As(V) for 1 h, then washed the cells with As(V)-free medium and measured how fast the cells exported the intracellular arsenic. Cells overexpressing *VBA3* exported arsenic more efficiently than wild-type cells carrying the empty plasmid whilst the difference between *PHO86* overexpressing and control cells was largely non-significant (Fig. 3b). These data argue against a model in which *PHO86* and *VBA3* overexpressing cells have a more efficient vacuolar sequestration capacity than wild-type cells.

Protein aggregation analyses

Arsenic has been shown to induce protein misfolding and aggregation (Jacobson et al, 2012; Tamás et al, 2014) and the protein aggregation levels can be used as a measure of intracellular damage caused by arsenic (Andersson et al, 2021; Jacobson et al, 2012; Talemi et al, 2014). To

assess protein aggregation *in vivo*, we monitored and quantified the subcellular distribution of GFP-tagged Hsp104, a molecular chaperone that associates with misfolded and aggregated proteins during As(III) stress (Jacobson et al, 2012). Plasmids overexpressing *PHO86*, *TUL1*, *VBA3*, or *UGP1* were introduced into BY4741 cells harboring *HSP104-GFP* integrated into the genome, the transformants exposed to 2 mM As(V) and the fraction of cells containing Hsp104-GFP foci/protein aggregates were quantified. As shown in Fig. 4, 60-70% of the cells had protein aggregates/Hsp104-GFP foci after 1 h of As(V) exposure. The fraction of cells with aggregates/Hsp104-GFP foci declined during the time-course of the experiment. Interestingly, *UGP1* overexpressing cells had significantly less aggregates after 3 h ($p=0.0002$) and 5 h ($p=0.02$) of As(V) exposure than cells carrying the empty vector. This is in line with the observation above that *UGP1* overexpressing cells accumulate less intracellular arsenic (Fig. 3). *PHO86* and *TUL1* overexpressing cells had similar levels of aggregated proteins as control cells carrying the empty plasmid (Fig. 4). Thus, the increased intracellular arsenic measured in *PHO86* overexpressing cells does not seem to cause enhanced protein aggregation. Cells overexpressing *VBA3* contained more protein aggregates at the 3 h ($p=0.004$) and 5 h time-points ($p=0.0002$) than control cells (Fig. 4). Thus, the increased intracellular arsenic levels in *VBA3* overexpressing cells (Fig. 3) appear to enhance protein misfolding/damage.

***PHO86* overexpression does not further improve As(V) resistance in mutants lacking specific phosphate transporters**

The high-affinity phosphate transporter Pho84 is known to play a role in As(V) import; its overexpression causes over-accumulation of As(V) in the cell (Shen et al, 2012), and *PHO84* deletion results in increased As(V) resistance (Bun-ya et al, 1996). Pho86 is required for the exit of Pho84 from the ER to the plasma membrane. In a *PHO86* deletion mutant, Pho84 is unable to localize to the plasma membrane (Lau et al, 2000) and *pho86Δ* cells are As(V) resistant (Bun-ya et al, 1996). To address whether Pho84 localization is affected by *PHO86* overexpression, we visualized the GFP-fused Pho84 protein in the absence and presence of As(V). We also visualized the GFP-fused Pho86 protein. As reported earlier (Lau et al, 2000), Pho84-GFP is localized to the plasma membrane whilst Pho86-GFP is localized to the ER (Fig. 5a). The presence of As(V) seemed to influence the localization of neither Pho84-GFP nor Pho86-GFP. In *PHO86* overexpressing cells, the overall signal intensity of Pho84-GFP was increased and Pho84-GFP was clearly visible in internal structures, likely the ER, and on the cell surface (Fig. 5a). Thus, more Pho84 appears to be present on the plasma membrane when

PHO86 is overexpressed. This is consistent with the observation that *PHO86* overexpressing cells have enhanced intracellular arsenic levels (Fig. 3). Because of the close functional relationship between Pho86 and Pho84, we asked whether Pho84 is needed for *PHO86* overexpression to confer As(V) resistance. In contrast to the effect in wild-type cells, overexpression of *PHO86* in the *pho84*Δ mutant did not result in an additional gain of As(V) resistance (Fig. 5b). The *S. cerevisiae* genome encodes five phosphate transporters and we overexpressed *PHO86* in mutants lacking the high-affinity transporter *PHO89* and the low-affinity transporters *PHO87*, *PHO90*, and *PHO91* (Wykoff & O'Shea, 2001) and scored growth of the transformants. Cells lacking *PHO87* and *PHO89* were more As(V) resistant than wild-type or *pho84*Δ cells carrying an empty plasmid (Fig. 5b). Thus, in addition to Pho84, Pho87 and Pho89 might be physiologically relevant As(V) import routes in yeast. The growth of *pho90*Δ and *pho91*Δ was similar to wild-type cells in the presence of As(V). *PHO86* overexpression enhanced As(V) resistance of the *pho90*Δ and *pho91*Δ mutants but not of the arsenate-resistant *pho87*Δ and *pho89*Δ mutants (Fig. 5b). We conclude that *PHO86* overexpression does not further improve As(V) resistance in mutants lacking specific phosphate transporters.

DISCUSSION

The yeast genome-wide deletion collection has provided important insights into arsenic toxicity and resistance mechanisms (Haugen et al, 2004; Jin et al, 2008; Jo et al, 2009; Johnson et al, 2016; Pan et al, 2010; Thorsen et al, 2009; Zhou et al, 2009). In contrast, genome-wide overexpression to identify arsenic resistance genes has, to our knowledge, only been performed once; that screen identified the arsenic resistance locus containing *ARR1/ACR1/YAP8*, *ARR2/ACR2*, and *ARR3/ACR3* (Bobrowicz et al, 1997) showing the power of the overexpression approach. In this current study, we screened a yeast genomic library and identified novel genes, namely *PHO86*, *TUL1*, *VBA3*, *UGP1*, and *ERG6* that provide resistance to arsenicals in yeast cells when overexpressed. Our screen also identified known arsenic resistance-related genes including *ARR1/ACR1/YAP8*, *ARR2/ACR2*, *ARR3/ACR3*, and *YAP1*. Some of the genes in the cassettes did not enhance resistance when overexpressed individually (*YJL118W*, *AIM26*, *MTC1*, *ASF1*, *NCA3*, *RPE1*, *ALB1*, *LSMI*). Individual overexpression of some genes (*SGE1*, *YPR196W*, *GIS4*, *MRPL39*) was not tested.

PHO86 overexpression conferred resistance to As(V) in wild-type and arsenic-sensitive GK4 cells but did not result in As(III) resistance, suggesting an As(V)-specific function of

Pho86 in resistance. Pho86 is an ER-resident protein that is necessary for the exit of Pho84 from the ER into COPII vesicles (Lau et al, 2000). Pho84 is a phosphate transporter that also mediates arsenate import into cells (Shen et al, 2012). Pho84 is not properly delivered to the plasma membrane in *pho86Δ* cells (Lau et al, 2000) leading to As(V) resistance (Bun-ya et al, 1996). Using a Pho84-GFP construct, we found that *PHO86* overexpression resulted in increased levels of Pho84 at the plasma membrane. Additionally, *PHO86* overexpressing cells had elevated levels of intracellular arsenic, consistent with a higher As(V) uptake capacity, likely mediated by Pho84. Indeed, overexpression of *PHO84* has previously been shown to result in higher intracellular accumulation of arsenic in the cell (Shen et al, 2012). Despite this, *PHO86* overexpressing cells did not have elevated intracellular damage, as determined by protein aggregation assays. These observations suggest that *PHO86* overexpressing cells efficiently protect the proteome from intracellular arsenic preventing its toxic effects on protein folding and activity. In addition to *pho84Δ*, we found that *pho87Δ* and *pho89Δ* cells were As(V) resistant suggesting that Pho87 and Pho89 are physiologically relevant As(V) import routes. *PHO86* overexpression did not further enhance resistance in the arsenate-resistant *pho84Δ*, *pho87Δ*, and *pho89Δ* mutants, suggesting that *PHO86*-mediated resistance is particularly relevant when cells accumulate high intracellular As(V). Indeed, we observed a substantial growth improvement upon *PHO86* overexpression of the As(V) sensitive strain GK4 that lacks arsenate reductase activity and capacity to convert intracellular As(V) to As(III). How does *PHO86* overexpression confer As(V) resistance despite higher intracellular arsenic levels? The yeast vacuole is a major storage compartment for phosphate and the vacuolar membrane contains proteins that mediate phosphate and As(V) transport (Booth & Guidotti, 1997). Our data indicate that *PHO86* overexpressing cells may not sequester large amounts of As(V) in the vacuoles since these cells exported arsenic as fast as control cells upon removal of extracellular As(V). Instead, we hypothesize that higher levels of intracellular phosphate may counteract the toxic effects of As(V) as increased amounts of Pho84 at the plasma membrane caused by *PHO86* overexpression not only results in increased As(V) uptake but also in increased phosphate uptake (Rothstein, 1963). Nevertheless, we cannot rule out other possible mechanisms. For example, similar to our observations, *PHO84* overexpression was reported to confer resistance to Mg^{2+} , Cu^{2+} , and Co^{2+} despite higher levels of intracellular metal accumulation. The same study showed that ectopic *PHO84* overexpression activated the unfolded protein response (UPR) pathway (Ofiteru et al, 2012). Thus, increased levels of Pho84, as observed in *PHO86* overexpressing cells, might result in UPR activation that in turn mitigates metal toxicity. Alternatively, As(V) can induce iron deficiency by triggering the

degradation of Fet3, a protein required for high-affinity iron uptake in yeast (Batista-Nascimento et al, 2013). Interestingly, Pho86 was shown to interact with Fet3 in a large-scale screen of membrane protein interactors (Miller et al, 2005) raising the possibility that *PHO86* overexpression alleviates As(V) toxicity by enhancing iron uptake. Nevertheless, the exact mechanism by which *PHO86* overexpression results in improved As(V) resistance remains to be elucidated.

VBA3 overexpression conferred resistance to both As(V) and As(III) and the effect was particularly pronounced in GK4 cells lacking the *ARR/ACR* gene cluster. *VBA3* encodes a basic amino acid transporter that mediates the influx of lysine and histidine into vacuoles (Shimazu et al, 2005). *VBA3* overexpressing cells had elevated intracellular arsenic content and protein misfolding/damage, yet these cells were As(V) resistant. One way this could be achieved is by Vba3-mediated transport of arsenic into the vacuole. However, *VBA3* overexpressing cells were more efficient in exporting arsenic than control cells once extracellular As(V) was removed, arguing against a function of Vba3 in vacuolar arsenic sequestration. We have previously shown that cells increase their protein degradation capacity during arsenic exposure to mitigate toxicity (Jacobson et al, 2012). It is reasonable to assume that there is a concomitant increase in amino acid uptake into vacuoles to maintain cytosolic concentrations of amino acids at appropriate levels during arsenic stress since high intracellular amino acid levels are toxic to cells (Hughes & Gottschling, 2012; Hughes et al, 2020; Ruiz et al, 2020). Whether *VBA3* overexpression confers As(V) resistance through an increased capacity to transport amino acids into the vacuole or through another mechanism remains to be established.

UGPI overexpression conferred As(V) resistance. Ugp1 catalyzes the reversible formation of UDP-glucose from glucose 1-phosphate and UTP, and the protein is involved in a variety of metabolic pathways. *UGPI* overexpression resulted in a modest decrease in intracellular arsenic and less damage as measured by protein aggregation. Elevated resistance in *UGPI* overexpressing cells might thus be caused by this reduction in intracellular arsenic. The mechanisms by which Ugp1 affects arsenic uptake or efflux remain unknown.

TUL1 overexpression conferred As(V) resistance without affecting intracellular arsenic or protein aggregation levels. Tul1 is a ubiquitin ligase required to sort membrane proteins into multivesicular bodies for further delivery to the vacuole (Reggiori & Pelham, 2002). Interestingly, *TUL1* overexpression has been shown to induce the degradation of several vacuolar membrane proteins, including Ycf1 (Li et al, 2015). Whether Tul1 contributes to

As(V) resistance through ubiquitination and degradation of these proteins or other mechanisms remains to be unveiled.

To sum up, this current study took advantage of a yeast genome library to identify genes that contribute to arsenic resistance upon overexpression. Some of the overexpressed genes had an impact on intracellular arsenic and protein aggregation levels giving clues on how these genes confer resistance. However, dedicated follow-up experiments will be required to provide insights into the underlying mechanisms.

Acknowledgments

The *PHO86*-GFP and *PHO84*-GFP plasmids were kindly provided by Erin K. O'Shea (Howard Hughes Medical Institute) and the GK4 strain was kindly provided by Robert Wysocki (University of Wroclaw). We also thank Stefanie Andersson for her expert technical assistance. This work was financially supported by the Swedish Research Council – Swedish Research Links program (VR: grant number 348–2014–4296) and the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (Formas: grant number 942–2015–376) to M.J.T.

Author contributions

Esin Isik: Investigation; Formal Analysis; Writing - Original Draft Preparation. **Çiğdem Balkan:** Investigation; Formal Analysis. **Vivien Karl:** Investigation; Formal Analysis. **Hüseyin Çağlar Karakaya:** Investigation; Formal Analysis. **Sansan Hua:** Investigation; Formal Analysis. **Sebastien Rauch:** Investigation; Formal Analysis. **Markus J. Tamás:** Conceptualization; Formal Analysis; Funding Acquisition; Writing - Original Draft Preparation; Writing – Review & Editing. **Ahmet Koc:** Conceptualization; Formal Analysis; Writing - Original Draft Preparation; Writing – Review & Editing.

Conflict of interest. None declared.

Ethics statement. None required.

Data availability. All data generated or analyzed during this study are included in this published article.

REFERENCES

- Ahmadpour, D., Maciaszczyk-Dziubinska, E., Babazadeh, R., Dahal, S., Migocka, M., Andersson, M., Wysocki, R., Tamás, M. J. & Hohmann, S. (2016) The mitogen-activated protein kinase Slr2 modulates arsenite transport through the aquaglyceroporin Fps1. *FEBS Lett*, 590(20), 3649-3659.
- Andersson, S., Romero, A., Rodrigues, J. I., Hua, S., Hao, X., Jacobson, T., Karl, V., Becker, N., Ashouri, A., Rauch, S., Nyström, T., Liu, B. & Tamás, M. J. (2021) Genome-wide imaging screen uncovers molecular determinants of arsenite-induced protein aggregation and toxicity. *J Cell Sci*, 134(11).
- Batista-Nascimento, L., Toledano, M. B., Thiele, D. J. & Rodrigues-Pousada, C. (2013) Yeast protective response to arsenate involves the repression of the high affinity iron uptake system. *Biochim Biophys Acta*, 1833(5), 997-1005.
- Beese, S. E., Negishi, T. & Levin, D. E. (2009) Identification of positive regulators of the yeast Fps1 glycerol channel. *PLoS Genet*, 5(11), e1000738.
- Bobrowicz, P., Wysocki, R., Owsianik, G., Goffeau, A. & Ulaszewski, S. (1997) Isolation of three contiguous genes, *ACR1*, *ACR2* and *ACR3*, involved in resistance to arsenic compounds in the yeast *Saccharomyces cerevisiae*. *Yeast*, 13(9), 819-828.
- Booth, J. W. & Guidotti, G. (1997) Phosphate transport in yeast vacuoles. *J Biol Chem*, 272(33), 20408-13.
- Bouganim, N., David, J., Wysocki, R. & Ramotar, D. (2001) Yap1 overproduction restores arsenite resistance to the ABC-transporter deficient mutant *ycf1* by activating *ACR3* expression. *Biochem Cell Biol*, 79, 441-448.
- Bun-ya, M., Shikata, K., Nakade, S., Yompakdee, C., Harashima, S. & Oshima, Y. (1996) Two new genes, *PHO86* and *PHO87*, involved in inorganic phosphate uptake in *Saccharomyces cerevisiae*. *Curr Genet*, 29(4), 344-351.
- Clemens, S. & Ma, J. F. (2016) Toxic Heavy Metal and Metalloid Accumulation in Crop Plants and Foods. *Annu Rev Plant Biol*, 67, 489-512.
- da Silva, S. M., Batista-Nascimento, L., Gaspar-Cordeiro, A., Vernis, L., Pimentel, C. & Rodrigues-Pousada, C. (2018) Transcriptional regulation of FeS biogenesis genes: A possible shield against arsenate toxicity activated by Yap1. *Biochim Biophys Acta Gen Subj*, 1862(10), 2152-2161.
- Garbinski, L. D., Rosen, B. P. & Chen, J. (2019) Pathways of arsenic uptake and efflux. *Environment International*, 126, 585-597.
- Ghosh, M., Shen, J. & Rosen, B. P. (1999) Pathways of As(III) detoxification in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA*, 96(9), 5001-5006.
- Guerra-Moreno, A., Isasa, M., Bhanu, M. K., Waterman, D. P., Eapen, V. V., Gygi, S. P. & Hanna, J. (2015) Proteomic Analysis Identifies Ribosome Reduction as an Effective Proteotoxic Stress Response. *J Biol Chem*, 290(50), 29695-706.
- Haugen, A. C., Kelley, R., Collins, J. B., Tucker, C. J., Deng, C., Afshari, C. A., Brown, J. M., Ideker, T. & Van Houten, B. (2004) Integrating phenotypic and expression profiles to map arsenic-response networks. *Genome Biol*, 5(12), R95.
- Hosiner, D., Lempiainen, H., Reiter, W., Urban, J., Loewith, R., Ammerer, G., Schweyen, R., Shore, D. & Schuller, C. (2009) Arsenic toxicity to *Saccharomyces cerevisiae* is a consequence of inhibition of the TORC1 kinase combined with a chronic stress response. *Mol Biol Cell*, 20(3), 1048-57.
- Hughes, A. L. & Gottschling, D. E. (2012) An early age increase in vacuolar pH limits mitochondrial function and lifespan in yeast. *Nature*, 492(7428), 261-5.

- Hughes, C. E., Coody, T. K., Jeong, M. Y., Berg, J. A., Winge, D. R. & Hughes, A. L. (2020) Cysteine Toxicity Drives Age-Related Mitochondrial Decline by Altering Iron Homeostasis. *Cell*, 180(2), 296-310 e18.
- Hughes, M. F., Beck, B. D., Chen, Y., Lewis, A. S. & Thomas, D. J. (2011) Arsenic exposure and toxicology: a historical perspective. *Toxicol Sci*, 123(2), 305-32.
- Huh, W. K., Falvo, J. V., Gerke, L. C., Carroll, A. S., Howson, R. W., Weissman, J. S. & O'Shea, E. K. (2003) Global analysis of protein localization in budding yeast. *Nature*, 425(6959), 686-91.
- Ibstedt, S., Sideri, T. C., Grant, C. M. & Tamás, M. J. (2014) Global analysis of protein aggregation in yeast during physiological conditions and arsenite stress. *Biol Open*, 3(10), 913-23.
- Jacobson, T., Navarrete, C., Sharma, S. K., Sideri, T. C., Ibstedt, S., Priya, S., Grant, C. M., Christen, P., Goloubinoff, P. & Tamás, M. J. (2012) Arsenite interferes with protein folding and triggers formation of protein aggregates in yeast. *J Cell Sci*, 125(Pt 21), 5073-83.
- Jin, Y. H., Dunlap, P. E., McBride, S. J., Al-Refai, H., Bushel, P. R. & Freedman, J. H. (2008) Global transcriptome and deletome profiles of yeast exposed to transition metals. *PLoS Genet*, 4(4), e1000053.
- Jo, W. J., Loguinov, A., Wintz, H., Chang, M., Smith, A. H., Kalman, D., Zhang, L., Smith, M. T. & Vulpe, C. D. (2009) Comparative functional genomic analysis identifies distinct and overlapping sets of genes required for resistance to monomethylarsonous acid (MMAIII) and arsenite (AsIII) in yeast. *Toxicol Sci*, 111(2), 424-36.
- Jochem, M., Ende, L., Isasa, M., Ang, J., Schnell, H., Guerra-Moreno, A., Micoogullari, Y., Bhanu, M., Gygi, S. P. & Hanna, J. (2019) Targeted Degradation of Glucose Transporters Protects against Arsenic Toxicity. *Mol Cell Biol*, 39(10).
- Johnson, A. J., Veljanoski, F., O'Doherty, P. J., Zaman, M. S., Petersingham, G., Bailey, T. D., Munch, G., Kersaitis, C. & Wu, M. J. (2016) Molecular insight into arsenic toxicity via the genome-wide deletion mutant screening of *Saccharomyces cerevisiae*. *Metallomics*, 8(2), 228-35.
- Kumar, N. V., Yang, J., Pillai, J. K., Rawat, S., Solano, C., Kumar, A., Grotli, M., Stemmler, T. L., Rosen, B. P. & Tamás, M. J. (2016) Arsenic directly binds to and activates the yeast AP-1-like transcription factor Yap8. *Mol Cell Biol*, 36(6), 913-22.
- Lau, W. T., Howson, R. W., Malkus, P., Schekman, R. & O'Shea, E. K. (2000) Pho86p, an endoplasmic reticulum (ER) resident protein in *Saccharomyces cerevisiae*, is required for ER exit of the high-affinity phosphate transporter Pho84p. *Proc Natl Acad Sci USA*, 97(3), 1107-1112.
- Lee, J. & Levin, D. E. (2018) Intracellular mechanism by which arsenite activates the yeast stress MAPK Hog1. *Mol Biol Cell*, 29(15), 1904-1915.
- Li, M., Koshi, T. & Emr, S. D. (2015) Membrane-anchored ubiquitin ligase complex is required for the turnover of lysosomal membrane proteins. *J Cell Biol*, 211(3), 639-52.
- Liu, Z., Boles, E. & Rosen, B. P. (2004) Arsenic trioxide uptake by hexose permeases in *Saccharomyces cerevisiae*. *J Biol Chem*, 279(17), 17312-8.
- Matia-Gonzalez, A. M. & Rodriguez-Gabriel, M. A. (2011) Slt2 MAPK pathway is essential for cell integrity in the presence of arsenate. *Yeast*, 28(1), 9-17.
- Menezes, R. A., Amaral, C., Delaunay, A., Toledano, M. & Rodrigues-Pousada, C. (2004) Yap8p activation in *Saccharomyces cerevisiae* under arsenic conditions. *FEBS Lett*, 566(1-3), 141-6.
- Miller, J. P., Lo, R. S., Ben-Hur, A., Desmarais, C., Stagljar, I., Noble, W. S. & Fields, S. (2005) Large-scale identification of yeast integral membrane protein interactions. *Proc Natl Acad Sci USA*, 102(34), 12123-8.

- Mukhopadhyay, R. & Rosen, B. P. (1998) *Saccharomyces cerevisiae* ACR2 gene encodes an arsenate reductase. *FEMS Microbiol Lett*, 168(1), 127-136.
- Mukhopadhyay, R., Shi, J. & Rosen, B. P. (2000) Purification and characterization of Acr2p, the *Saccharomyces cerevisiae* arsenate reductase. *J Biol Chem*, 275(28), 21149-21157.
- Nasmyth, K. A. & Reed, S. I. (1980) Isolation of genes by complementation in yeast: molecular cloning of a cell-cycle gene. *Proc Natl Acad Sci U S A*, 77(4), 2119-23.
- Naujokas, M. F., Anderson, B., Ahsan, H., Aposhian, H. V., Graziano, J. H., Thompson, C. & Suk, W. A. (2013) The broad scope of health effects from chronic arsenic exposure: update on a worldwide public health problem. *Environ Health Perspect*, 121(3), 295-302.
- Ofiteru, A. M., Ruta, L. L., Rotaru, C., Dumitru, I., Ene, C. D., Neagoe, A. & Farcasanu, I. C. (2012) Overexpression of the *PHO84* gene causes heavy metal accumulation and induces Ire1p-dependent unfolded protein response in *Saccharomyces cerevisiae* cells. *Appl Microbiol Biotechnol*, 94(2), 425-35.
- Pan, X., Reissman, S., Douglas, N. R., Huang, Z., Yuan, D. S., Wang, X., McCaffery, J. M., Frydman, J. & Boeke, J. D. (2010) Trivalent arsenic inhibits the functions of chaperonin complex. *Genetics*, 186(2), 725-34.
- Reggiori, F. & Pelham, H. R. (2002) A transmembrane ubiquitin ligase required to sort membrane proteins into multivesicular bodies. *Nat Cell Biol*, 4(2), 117-23.
- Rosen, B. P. & Tamás, M. J. (2010) Arsenic transport in prokaryotes and eukaryotic microbes. *Adv Exp Med Biol*, 679, 47-55.
- Rothstein, A. (1963) Interactions of arsenate with the phosphate-transporting system of yeast. *J Gen Physiol*, 46, 1075-85.
- Ruiz, S. J., van 't Klooster, J. S., Bianchi, F. & Poolman, B. (2020) Growth Inhibition by Amino Acids in *Saccharomyces cerevisiae*. *Microorganisms*, 9(1).
- Shen, M. W., Shah, D., Chen, W. & Da Silva, N. (2012) Enhanced arsenate uptake in *Saccharomyces cerevisiae* overexpressing the Pho84 phosphate transporter. *Biotechnol Prog*, 28(3), 654-61.
- Shen, S., Li, X. F., Cullen, W. R., Weinfeld, M. & Le, X. C. (2013) Arsenic Binding to Proteins. *Chem Rev*.
- Shimazu, M., Sekito, T., Akiyama, K., Ohsumi, Y. & Kakinuma, Y. (2005) A family of basic amino acid transporters of the vacuolar membrane from *Saccharomyces cerevisiae*. *J Biol Chem*, 280(6), 4851-7.
- Slyemi, D. & Bonnefoy, V. (2012) How prokaryotes deal with arsenic. *Environ Microbiol Rep*, 4(6), 571-86.
- Szybalski, W. & Bryson, V. (1952) Genetic studies on microbial cross resistance to toxic agents. I. Cross resistance of *Escherichia coli* to fifteen antibiotics. *J Bacteriol*, 64(4), 489-99.
- Talemi, S. R., Jacobson, T., Garla, V., Navarrete, C., Wagner, A., Tamás, M. J. & Schaber, J. (2014) Mathematical modelling of arsenic transport, distribution and detoxification processes in yeast. *Mol Microbiol*, 92(6), 1343-56.
- Tamás, M. J., Sharma, K. S., Ibstedt, S., Jacobson, T. & Christen, P. (2014) Heavy metals and metalloids as a cause for protein misfolding and aggregation. *Biomolecules*, 4, 252-267.
- Thorsen, M., Di, Y., Tangemo, C., Morillas, M., Ahmadpour, D., Van der Does, C., Wagner, A., Johansson, E., Boman, J., Posas, F., Wysocki, R. & Tamás, M. J. (2006) The MAPK Hog1p modulates Fps1p-dependent arsenite uptake and tolerance in yeast. *Mol Biol Cell*, 17(10), 4400-4410.
- Thorsen, M., Jacobson, T., Vooijs, R., Navarrete, C., Bliet, T., Schat, H. & Tamás, M. J. (2012) Glutathione serves an extracellular defence function to decrease arsenite accumulation and toxicity in yeast. *Mol Microbiol*, 84(6), 1177-88.

- Thorsen, M., Lagniel, G., Kristiansson, E., Junot, C., Nerman, O., Labarre, J. & Tamás, M. J. (2007) Quantitative transcriptome, proteome, and sulfur metabolite profiling of the *Saccharomyces cerevisiae* response to arsenite. *Physiol Genomics*, 30(1), 35-43.
- Thorsen, M., Perrone, G. G., Kristiansson, E., Traini, M., Ye, T., Dawes, I. W., Nerman, O. & Tamás, M. J. (2009) Genetic basis of arsenite and cadmium tolerance in *Saccharomyces cerevisiae*. *BMC Genomics*, 10, 105.
- Wykoff, D. D. & O'Shea, E. K. (2001) Phosphate transport and sensing in *Saccharomyces cerevisiae*. *Genetics*, 159(4), 1491-9.
- Wysocki, R., Bobrowicz, P. & Ulaszewski, S. (1997) The *Saccharomyces cerevisiae* ACR3 gene encodes a putative membrane protein involved in arsenite transport. *J Biol Chem*, 272(48), 30061-30066.
- Wysocki, R., Chéry, C. C., Wawrzycka, D., Van Hulle, M., Cornelis, R., Thevelein, J. M. & Tamás, M. J. (2001) The glycerol channel Fps1p mediates the uptake of arsenite and antimonite in *Saccharomyces cerevisiae*. *Mol Microbiol*, 40, 1391-1401.
- Wysocki, R., Fortier, P. K., Maciaszczyk, E., Thorsen, M., Leduc, A., Odhagen, A., Owsianik, G., Ulaszewski, S., Ramotar, D. & Tamás, M. J. (2004) Transcriptional activation of metalloids tolerance genes in *Saccharomyces cerevisiae* requires the AP-1-like proteins Yap1p and Yap8p. *Mol Biol Cell*, 15(5), 2049-60.
- Wysocki, R. & Tamás, M. J. (2010) How *Saccharomyces cerevisiae* copes with toxic metals and metalloids. *FEMS Microbiol Rev*, 34(6), 925-51.
- Yompakdee, C., Bun-ya, M., Shikata, K., Ogawa, N., Harashima, S. & Oshima, Y. (1996) A putative new membrane protein, Pho86p, in the inorganic phosphate uptake system of *Saccharomyces cerevisiae*. *Gene*, 171(1), 41-47.
- Zhou, X., Arita, A., Ellen, T. P., Liu, X., Bai, J., Rooney, J. P., Kurtz, A. D., Klein, C. B., Dai, W., Begley, T. J. & Costa, M. (2009) A genome-wide screen in *Saccharomyces cerevisiae* reveals pathways affected by arsenic toxicity. *Genomics*, 94(5), 294-307.

TABLES

Table 1: The genomic regions of the plasmids from arsenate-/arsenite-resistant colonies.

Colony #	Chromosome	Region	Candidate Genes
AsV.1	X	188721 bp to 195519 bp	NCA3, PHO86, YJL118W, RPE1, ALB1, MTC1
AsV.2	X	191189 bp to 196530 bp	ASF1, NCA3, PHO86, YJL118W
AsV.3	XVI	929988 bp to 941100 bp	ARR1, ARR2, ARR3, SGE1, YPR196W
AsV.4	X	368844 bp to 373815 bp	TUL1, UGP1, AIM26
AsV.5	X	188958 bp to 196530 bp	ASF1, NCA3, PHO86, YJL118W, RPE1, ALB1, MTC1
AsV.6	X	191207 bp to 196530 bp	NCA3, PHO86, YJL118W
AsV.7	X	189834 bp to 204440 bp	ASF1, NCA3, PHO86, YJL118W, RPE1, ALB1
AsV.8	X	190300 bp to 196690 bp	ASF1, NCA3, PHO86, YJL118W, RPE1
AsV.9	X	188958 bp to 194280 bp	NCA3, PHO86, YJL118W, RPE1, ALB1
AsV.10	X	186335 bp to 193712 bp	PHO86, YJL118W, RPE1, ALB1, MTC1, LSM1
AsIII.1	XVI	932323 bp to 943489 bp	ARR1, ARR2, ARR3, SGE1
AsIII.2	XIII	250986 bp to 256885 bp	GIS4, YAP1, ERG6, MRPL39
AsIII.3	III	9676 bp to 14296 bp	VBA3
AsIII.4	XIII	250986 bp to 256885 bp	GIS4, YAP1, ERG6, MRPL39

FIGURE LEGENDS

Fig. 1 Screening of a genomic library for arsenic resistance in yeast and identified hits. (a) Cartoon of the screening set-up and arsenic concentrations used. Growth of yeast gDNA transformants on gradient plates containing (b) arsenate [As(V)] and (c) arsenite [As(III)]. Growth was scored after incubating the cells for 3-5 days at 30°C. P-only represents the empty vector containing cells.

Fig. 2 Growth of yeast cells (wild-type W303-1A and arsenic sensitive GK4) overexpressing individual candidate genes in the presence of arsenate [As(V)] (a, b) and arsenite [As(III)] (c, d). Growth was scored on gradient plates after incubating the cells for 3-5 days at 30°C. Growth of BY4741 cells overexpressing individual candidate genes in the presence of As(V) (e). Growth on plates containing the given As(V) concentrations was scored after incubating the cells for 2-3 days at 30°C. P-only represents the empty vector containing cells.

Fig. 3 Intracellular arsenic concentrations of BY4741 cells overexpressing the *PHO86*, *VBA3*, *TUL1*, and *UGP1* genes or the empty vector control (p-only). (a) Arsenic accumulation. Intracellular arsenic levels were measured in untreated cells and cells exposed to 1 mM As(V) for 1 h. Values represent averages from six independent biological replicates (n=6) and the error bars represent the standard deviation. (b) Arsenic efflux. Cells were incubated with 1 mM As(V) for 1 h, then washed with As(V)-free medium, and samples were taken at the indicated time points for intracellular arsenic determination. The intracellular arsenic levels measured before the wash were set to 1 to allow comparisons between cells overexpressing different genes. The values represent averages from three independent biological replicates (n=3) and the error bars represent the standard deviation. Statistical analyses were performed by Student's t-test, and significant differences are indicated; * $p < 0.05$ and ** $p < 0.005$.

Fig. 4 Protein aggregation levels of cells overexpressing the *PHO86*, *VBA3*, *TUL1*, and *UGP1* genes. P-only represents the empty vector containing cells. Hsp104-GFP localization was monitored in BY4741 cells by fluorescence microscopy before and after exposure to 2 mM As(V). The fractions of cells containing aggregates/Hsp104-GFP foci were determined by visual inspection of 144-282 cells per strain and time point. Values represent averages from three independent biological replicates (n=3) and the error bars represent the standard deviation. Statistical analyses were performed by Student's t-test, and significant differences are indicated; * $p < 0.05$ and ** $p < 0.005$.

Fig. 5 *PHO86* overexpression and phosphate transporters. (a) Localization of Pho84-GFP in BY4741 cells overexpressing *PHO86* and exposed to As(V). (b) Growth of BY4741 cells (wild-type and mutants lacking individual phosphate transporters) overexpressing *PHO86* in the presence of As(V). Growth was scored after incubating the cells for 2-3 days at 30°C. P-only represents the empty vector containing cells.

Appendix

Table A1: *S. cerevisiae* strains used in this study

Strain	Genotype	Source
BY4741	MATa , his3Δ1 leu2Δ0 met15Δ0 ura3Δ0	EUROSCARF
BY4743	MATa /α his3Δ1/his3Δ1 leu2Δ0/leu2Δ0 LYS2/lys2Δ0 met15Δ0/MET15 ura3Δ0/ura3Δ0	EUROSCARF
(n) Δgene	BY4741 ΔDeleted gene:KAN ^R	EUROSCARF
(2n) Δgene	BY4743 ΔDeleted gene:KAN ^R	EUROSCARF
W303-1A	MATa ura3-1 leu2-3/112 trp1 -1 his3-11/15 ade2- 1 can1-100 GAL SUC2 mal0	Thomas and Rothstein (1989) (b)
GK4(a)	W303-1A acr1Δ acr2Δ acr3Δ::loxP-kanMX-loxP	Robert Wysocki
HSP104-GFP	BY4741 HSP104-GFP- HIS3-MX6	Invitrogen

(a)*ACR1* is also called *ARR1/YAP8*; *ACR2* is also called *ARR2*; *ACR3* is also called *ARR3*

(b)Thomas, B. J. & Rothstein, R. (1989). Elevated recombination rates in transcriptionally active DNA. *Cell* 56, 619-630.

Figure A1: Determining the minimum inhibitory concentration (MIC) for arsenate [As(V)] in the indicated yeast strains.

Figure A2: Determining the minimum inhibitory concentration (MIC) for arsenite [As(III)] in the indicated yeast strains.

Figure A3: Growth of yeast gDNA transformants on plates containing given concentrations of (a) arsenate [As(V)] and (b) arsenite [As(III)]. Growth was scored after incubating the cells for 5 days at 30°C. P-only represents the empty vector.

Figure A4: Growth of yeast (W303-1A) cells expressing individual candidate genes in the absence and presence of given concentrations of As(V). Growth was scored after incubating the cells for 3-5 days at 30°C. P-only represents the empty vector.




Figure 1





Figure 2

(a)



(b)




Figure 3




Figure 4




Figure 5




Figure A1





Figure A2

(a)



(b)







Figure A3

**Figure A4**