

STABILITY OF STEADY-STATE SOLUTIONS OF A CLASS OF KELLER-SEGEL MODELS WITH MIXED BOUNDARY CONDITIONS

ZEFU FENG, JING JIA, AND SHOUMING ZHOU

ABSTRACT. In this paper, we investigate the the existence and stability of non-trivial steady state solutions of a class of chemotaxis models with zero-flux boundary conditions and Dirichlet boundary conditions on one-dimensional bounded interval. By using upper-lower solution and the monotone iteration scheme method, we get the existence of the steady-state solution of the chemotaxis model. Moreover, by adopting the “inverse derivative” technique and the weighted energy method to obtain the stability of the steady-state solution of this chemotaxis model.

1. INTRODUCTION

Chemotaxis phenomenon is an essential and basic property of the living cells which responses to the environments changes, it describes a directional movement of the cells to the gradient of chemical concentrations, such as aggregation of bacteria, slime mould formation, fish pigmentation, tumor angiogenesis, blood vessel formation, wound healing/inflammation, and cancer metastasis[2, 3, 33]. The chemotaxis model was first proposed by Patlak [18] when studying the random motion of particles, and later Keller and Segel [19] proposed a chemotaxis model when studying amoeba aggregation effects. The general mathematical Keller-Segel model reads as:

$$\begin{cases} u_t = \Delta u - \nabla \cdot (u \nabla \phi(w)), & x \in \Omega, \\ w_t = D \Delta w + g(u, w), & x \in \Omega, \end{cases} \quad (1.1)$$

where u and w represent the bacterial density and oxygen concentration at position x and time t , respectively. The constant $D > 0$ represent the chemical diffusion coefficient. The functional $\phi(w)$ is called the chemotaxis sensitivity function accounting for the signal response mechanism and the reaction term $g(u, w)$ is the chemical kinetics (growth and degradation). The chemotactic sensitivity function $\phi(w)$ usually has two prototypes: $\phi(w) = \ln w$ (logarithmic sensitivity) and $\phi(w) = w$ (linear sensitivity). Logarithmic sensitivity was originally used in [19], based on Weber-Fechner’s law (the sensory response to stimuli is logarithmic), and it has various biological applications (cf. [9, 16, 21]). The reaction function usually can be chosen as $g(u, w) = u^\gamma w^m$ with nonnegative constant parameters $\gamma \geq 1$ and $m \geq 0$, when $\gamma = 1$ and for any $0 \leq m \leq 1$, the existence of the traveling wave solution of (1.1) for logarithmic sensitivity was reported in [17, 30, 34]. The stability of the travel wavefront ($m = 1$) was investigated in [6, 7, 23, 24, 25, 29]. The instability of pulsation waves ($m = 0$) was studied in [8, 26] and the stability of boundary layers (see [14, 15]). We also refer to [10, 31, 32] for the global existence and large-time behavior of solution to (1.1) with $\gamma = m = 1$ and linear sensitivity under the Neumann boundary conditions $\partial_\nu u|_{\partial\Omega} = \partial_\nu w|_{\partial\Omega} = 0$. When the physical boundary conditions, namely, the zero-flux boundary condition and Dirichlet boundary condition are imposed to cell density u and chemical concentration w

2020 *Mathematics Subject Classification.* 35B40, 35B45, 35Q92.

Key words and phrases. Chemotaxis, steady state, existence, stability.

(see also the experiment in [1]):

$$\partial_\nu u - u\partial_\nu w = 0, w = w_*, x \in \partial\Omega, \quad (1.2)$$

where ν is the outward unit normal vector of $\partial\Omega$ and $w_* > 0$ represents the boundary data of w on the $\partial\Omega$. Until very recently, significant progress has been made for (1.1)-(1.2), the authors of [5] showed that under the boundary conditions (1.2) and the parametric restrictions $\gamma = 1$, $m \geq 0$ and $\chi > |m - 1|$, system (1.1)-(1.2) with $\phi(w) = \ln w$ generates the so-called boundary spike layer (BSL) solutions that exist on the half-line: $[0, \infty)$ under smallness assumptions on the initial perturbations. When $\gamma = 1$, $m = 1$ and $\phi(w) = w$, Braukhoff and Lankeit in [4] proved the stationary solution under the no-flux boundary conditions for u and the physically meaningful condition for w in bounded domain $\Omega \subset \mathbb{R}^n, n \geq 1$. Later, Lee et.al [20] proved the existence and uniqueness of the steady-state solution of equations (1.1)-(1.2) in space $\Omega \subset \mathbb{R}^n (n \geq 1)$; in [35], Winkler consider the existence of globally weak solutions in one-dimensional bounded interval; later, Hong and Wang [13] improved the result of [20] to obtain the asymptotic stability of the steady-state solution in the bounded domain $\Omega = (0, 1)$. When $\gamma > 1, m = 0$, we refer the readers to [36, 12] where the global existence and large time behavior of the one dimensional Cauchy problem solution of equation and dynamic boundary condition problem were proved, respectively. In [11], the author also extended the result [5] to the case $1 < \gamma < 2$ on the half line. However, the stability of steady-state solution to (1.1)-(1.2) with linear sensitivity function is an open problem, which has more biological significance. Motivated by the ideas of [20] and [13], we are devoted to investigating the following system of reaction-diffusion-advection equations:

$$\begin{cases} u_t = u_{xx} - \chi(uw_x)_x, & \text{in } \mathcal{I}, \\ w_t = Dw_{xx} - u^\gamma, & \text{in } \mathcal{I}, \\ (u, w)(x, 0) = (u_0, w_0)(x), & \text{in } \mathcal{I}, \end{cases} \quad (1.3)$$

with physical mixed boundary condition

$$(u_x - \chi uw_x)|_{x=0,1} = 0, w(0, t) = w(1, t) = w_*. \quad (1.4)$$

where $(x, t) \in \mathcal{I} \times \mathbb{R}^+$ and $1 < \gamma \leq 2$.

With the zero-flux boundary condition (1.4) on u , by integrating the first equation (1.3), we immediately find that the cell mass is preserved in time, namely :

$$\int_{\mathcal{I}} u(x, t) dx = \int_{\mathcal{I}} u_0(x) dx := M,$$

where $M > 0$ denotes the initial cell mass. Then the stationary solution (\bar{u}, \bar{w}) of (1.3) subject to boundary condition satisfies

$$\begin{cases} \bar{u}_{xx} - \chi(\bar{u}\bar{w}_x)_x = 0, & x \in \mathcal{I}, \\ D\bar{w}_{xx} - \bar{u}^\gamma = 0, & x \in \mathcal{I}, \\ \int_{\mathcal{I}} \bar{u}(x, t) dx = M, \\ (\bar{u}_x - \chi\bar{u}\bar{w}_x)|_{x=0,1} = 0, \bar{w}|_{x=0,1} = w_*. \end{cases} \quad (1.5)$$

The goal of this paper is twofold. First, we can use a similar method as in [20] to prove the existence of the classical solution to equation (1.5). The second goal of this paper is to show the nonlinear stability of the stationary solution (\bar{u}, \bar{w}) , we can show that the system (1.3) with (1.4) admits a unique solution (u, w) satisfying:

$$\|(u, w) - (\bar{u}, \bar{w})\|_{L^\infty} \rightarrow 0 \text{ as } t \rightarrow \infty.$$

The rationale underlining the studies of such a model is that monomials are the building blocks of genuinely nonlinear functions. Hence, the study of the model will help shed light on how to advance fundamental research of chemotaxis models with nonlinear rates of chemical production/consumption. Our intention of this paper is to continue exploring the mathematical properties of the model by investigating the existence and stability of steady state solutions under the same boundary conditions as specified in (1.4).

The rest of this paper is organized as follows: In Sec. 2, we state our main results on the existence and stability of stationary solution to the (1.3)–(1.4). Then we prove the existence results and stability results in Sec. 3.

Notations. Throughout the paper, we denote by L^∞, L^2, H_0^1 and H^k the standard function spaces $L^\infty(\mathcal{I}), L^2(\mathcal{I}), H_0^1(\mathcal{I})$ and $H^k(\mathcal{I})$, respectively. We denote $\bar{\mathcal{I}}$ by the closure of \mathcal{I} and by C a generic time-independent constant which may take different values in different places. In the sequel, we often omit \mathcal{I} without ambiguity.

2. MAIN RESULTS

In this section, we introduce the existence of the stationary solution of the equation (1.5) and state our main results on the stability of stationary solutions. The first results on the existence of steady state solutions are stated below.

Theorem 2.1. *For any $M \in (0, \infty)$, the system (1.5) admits a unique classical non-constant solution $(\bar{u}, \bar{w}) \in C^1(\bar{\mathcal{I}}) \cap C^\infty(\mathcal{I})$ such that*

$$\bar{u} = \frac{M}{\int_{\mathcal{I}} e^{x\bar{w}} dx} e^{x\bar{w}}, \bar{u} > 0, 0 < \bar{w} \leq w_* \text{ for any } x \in \bar{\mathcal{I}}. \quad (2.1)$$

The second result is the nonlinear local stability of stationary solutions obtained in Theorem 2.1 for the initial-boundary value problem (1.3)-(1.4) as time goes to infinity.

Theorem 2.2. *Suppose that $u_0 \in H^1$ and $w_0 \in H^2$ with $u_0 \geq 0, w_0 \geq 0$ such that $\int_{\mathcal{I}} u_0 dx = M$. Let (\bar{u}, \bar{w}) be the stationary solution given in Theorem 2.1 with $\int_{\mathcal{I}} \bar{u} dx = M$ and define*

$$\varphi_0(x) = \int_0^x (u_0(y) - \bar{u}(y)) dy.$$

Then there exists a constant $\delta_0 > 0$ such that if

$$\|\varphi_0\|_{H^2}^2 + \|w_0 - \bar{w}\|_{H^2}^2 \leq \delta_0,$$

then the initial-boundary value problem (1.3)–(1.4) admits a unique global solution (u, w) satisfying

$$u \in C([0, \infty); H^1) \cap L^2([0, \infty); H^2), w \in C([0, \infty); H^2) \cap L^2([0, \infty); H^3),$$

and the following large time behavior:

$$\|(u - \bar{u}, w - \bar{w})(\cdot, t)\|_{L^\infty} \rightarrow 0. \quad (2.2)$$

3. EXISTENCE AND STABILITY FOR THE CASE $D > 0$

3.1. Existence of stationary solution. In this section, we first prove the existence of steady state of (1.5). We start the proof by considering the following auxiliary problem

$$\begin{cases} D\bar{w}_{\kappa,xx} = (\kappa M e^{\chi \bar{w}_{\kappa}})^{\gamma}, & x \in \mathcal{I}, \\ \bar{w}_{\kappa} = w_*, & x = 0, 1, \end{cases} \quad (3.1)$$

where κ is an arbitrary positive constant. Since $w_* > 0$, by maximal principle we have

$$0 \leq \bar{w}_{\kappa} \leq w_*, x \in \mathcal{I}.$$

Then it is not difficult to see that $\bar{w}_{\kappa} = w_*$ is a super-solution of (3.1), while $\bar{w}_{\kappa} = 0$ provides a sub-solution. Therefore, following the standard monotone iteration scheme and the fact that $f(x) = e^x$ is increasing function for x positive, we immediately know that (3.1) has a unique classical solution \bar{w}_{κ} satisfying

$$0 \leq \bar{w}_{\kappa} \leq w_*.$$

Now we claim that there exists κ_m such that

$$\kappa_m \int_{\mathcal{I}} e^{\chi \bar{w}_{\kappa_m}} dx = 1. \quad (3.2)$$

We postpone the proof of (3.2) in lemma 3.1. Using this claim we can easily see that $\bar{w} = \bar{w}_{\kappa_m}$ is a solution of (1.5).

In order to prove the claim (3.2), we give the following lemma.

Lemma 3.1. *Let $\kappa_1 \geq \kappa_2 > 0$ and $\bar{w}_{\kappa_i}, i = 1, 2$ be the solutions of (3.1) with $\kappa = \kappa_i, i = 1, 2$ respectively. Then*

$$0 \leq \bar{w}_{\kappa_2} - \bar{w}_{\kappa_1} \leq \left(\left(\frac{\kappa_1}{\kappa_2} \right)^{\gamma} - 1 \right) \frac{e^{\gamma \chi w_*}}{\gamma \chi}. \quad (3.3)$$

Moreover, there exists a constant κ_m such that

$$\kappa_m \int_{\mathcal{I}} e^{\chi \bar{w}_{\kappa_m}} dx = 1. \quad (3.4)$$

Proof. The left-hand side inequality follows from the standard comparison theorem directly. We only prove the inequality for the right-hand side. Due to the fact $\kappa_1 \geq \kappa_2 > 0$ and $\bar{w}_{\kappa_1} > 0$, it's easy to see that

$$\begin{aligned} D(\bar{w}_{\kappa_1,xx} - \bar{w}_{\kappa_2,xx}) &= (\kappa_1 M e^{\chi \bar{w}_{\kappa_1}})^{\gamma} - (\kappa_2 M e^{\chi \bar{w}_{\kappa_2}})^{\gamma} \\ &= (\kappa_2 M)^{\gamma} (e^{\gamma \chi \bar{w}_{\kappa_1}} - e^{\gamma \chi \bar{w}_{\kappa_2}}) + ((\kappa_1 M)^{\gamma} - (\kappa_2 M)^{\gamma}) e^{\gamma \chi \bar{w}_{\kappa_1}} \\ &\leq (\kappa_2 M)^{\gamma} F(\gamma \chi \bar{w}_{\kappa_1}, \gamma \chi \bar{w}_{\kappa_2}) (\gamma \chi \bar{w}_{\kappa_1} - \gamma \chi \bar{w}_{\kappa_2}) \\ &\quad + ((\kappa_1 M)^{\gamma} - (\kappa_2 M)^{\gamma}) e^{\gamma \chi w_*} \end{aligned} \quad (3.5)$$

where

$$F(a, b) = \begin{cases} \frac{e^a - e^b}{a - b}, & \text{if } a \neq b, \\ e^a, & \text{if } a = b. \end{cases}$$

From the fact

$$0 < \bar{w}_{\kappa_1} \leq \bar{w}_{\kappa_2} \leq w_*, \quad (3.6)$$

we have

$$1 < F(\gamma \chi \bar{w}_{\kappa_1}, \gamma \chi \bar{w}_{\kappa_2}) \leq e^{\gamma \chi w_*}. \quad (3.7)$$

By (3.5) and (3.7), we have

$$\begin{aligned}\bar{w}_{\kappa_2} - \bar{w}_{\kappa_1} &\leq \frac{((\kappa_1 M)^\gamma - (\kappa_2 M)^\gamma) e^{\gamma \chi \bar{w}_{\kappa_1}}}{(\kappa_2 M)^\gamma F(\gamma \chi \bar{w}_{\kappa_1}, \gamma \chi \bar{w}_{\kappa_2}) \gamma \chi} \\ &\leq \left(\left(\frac{\kappa_1}{\kappa_2} \right)^\gamma - 1 \right) \frac{e^{\gamma \chi w_*}}{\gamma \chi}.\end{aligned}$$

Thus, we prove the right-hand side of (3.3). It implies that the continuity of \bar{w}_κ with respect to κ . On the other hand, we have

$$\lim_{\kappa \rightarrow 0^+} \kappa \int_{\mathcal{I}} e^{\chi \bar{w}_\kappa} dx = 0 \text{ and } \lim_{\kappa \rightarrow \infty} \kappa \int_{\mathcal{I}} e^{\chi \bar{w}_\kappa} dx = \infty.$$

Then we can find κ_m such that $\kappa_m \int_{\mathcal{I}} e^{\chi \bar{w}_{\kappa_m}} dx = 1$ and it completes the proof of lemma 3.1. \square

3.2. Nonlinear asymptotic stability. In this section, we shall focus on attention to investigate the global well-posedness and stability of the steady state of (1.3)-(1.4) for $D > 0$ by the method of energy estimates when the initial data (u_0, w_0) is a small perturbation of (\bar{u}, \bar{w}) . Before proceeding, we present an well-known inequality that will be frequently used in the sequel.

Lemma 3.2. (cf.[27]). *For any $f \in H^1(\mathcal{I})$, there exists a constant $c_1 > 0$ such that*

$$\|f\|_{L^\infty} \leq c_1 (\|f\|_{L^2}^{\frac{1}{2}} \|f_x\|_{L^2}^{\frac{1}{2}} + \|f\|_{L^2}). \quad (3.8)$$

Furthermore, if $f \in H_0^1(\mathcal{I})$, then it holds that

$$\|f\|_{L^\infty} \leq c_2 \|f\|_{L^2}^{\frac{1}{2}} \|f_x\|_{L^2}^{\frac{1}{2}} \text{ and } \|f\|_{L^\infty} \leq c_3 \|f_x\|_{L^2} \quad (3.9)$$

for some constants $c_2, c_3 > 0$.

3.2.1. Reformulation of the problem. Now let us use energy methods and anti-derivatives technique to prove Theorem 2.2. Integrating the equation (1.5)₁ over $(0, x)$, we obtain that the stationary solution (\bar{u}, \bar{w}) satisfies

$$\begin{cases} \bar{u}_x - \chi(\bar{u}\bar{w})_x = 0, \\ D\bar{w}_{xx} - \bar{u}^\gamma = 0, \\ \bar{w}(0) = \bar{w}(1) = w_*. \end{cases} \quad (3.10)$$

With the zero-flux boundary condition of u in (1.4), which combined with the facts $\int_{\mathcal{I}} \bar{u} dx = \int_{\mathcal{I}} u_0 dx = M$ implies that

$$\int_{\mathcal{I}} (u(x, t) - \bar{u}(x)) dx = 0$$

for any $t \geq 0$. This enables us to define anti-derivatives of the perturbed functions and more importantly, carry out stability analysis in the framework of Sobolev spaces. Define

$$\varphi(x, t) = \int_0^x (u(y, t) - \bar{u}(y)) dy, \psi = w - \bar{w},$$

which implies

$$u = \varphi_x + \bar{u}, w = \psi + \bar{w}. \quad (3.11)$$

Substituting (3.11) into (1.3), by integrating the first resulting equation with the spatial variable from 0 to x , then using (3.10), we can show that:

$$\begin{cases} \varphi_t = \varphi_{xx} - \chi(\varphi_x \psi_x + \bar{u} \psi_x + \varphi_x \bar{w}_x), \\ \psi_t = D\psi_{xx} + \bar{u}^\gamma - (\varphi_x + \bar{u})^\gamma, \end{cases} \quad (3.12)$$

with the initial data

$$(\varphi, \psi)(x, 0) = (\varphi_0, \psi_0) = \left(\int_0^x (u_0(y) - \bar{u}(y)) dy, w_0 - \bar{w} \right) \quad (3.13)$$

and the boundary conditions

$$(\varphi, \psi)(0, t) = (\varphi, \psi)(1, t) = 0. \quad (3.14)$$

By standard approaches, such as iteration scheme and fixed point theorems (cf. [22, 28]), one can prove the local existence of solutions to the initial-boundary value problem (3.12)-(3.14). To simplify the presentation, we only record the result here without producing the proof. Precisely, for any $T > 0$, if we define

$$\begin{aligned} X(0, T) := & \{(\varphi, \psi) | \varphi \in \mathcal{C}([0, T]; H_0^1 \cap H^2) \cap L^2([0, T]; H^3), \\ & \psi \in \mathcal{C}([0, T]; H_0^1 \cap H^2) \cap L^2([0, T]; H^3)\} \end{aligned}$$

and denote

$$N(T) := \sup_{0 \leq t \leq T} (\|\varphi\|_{H^2}^2 + \|\psi\|_{H^2}^2),$$

then we have the following local existence result.

Proposition 3.3. (*Local existence*). *Let $\varphi_0 \in H_0^1 \cap H^2$ and $\psi_0 \in H_0^1 \cap H^2$ such that*

$$\varphi_{0,x} + \bar{u} \geq 0, \psi_0 + \bar{w} \geq 0$$

for any $x \in \mathcal{I}$. Then there exists a positive constant T_0 depending on N_0, \bar{u} and \bar{w} such that the initial-boundary value problem (3.12)-(3.14) admits a unique solution $(\varphi, \psi) \in X(0, T_0)$ satisfying $N(T_0) \leq 2N(0)$ and

$$\varphi_x + \bar{u} \geq 0, \psi + \bar{w} \geq 0$$

for any $(x, t) \in \mathcal{I} \times [0, T_0]$.

Next, we shall derive the *a priori* estimates of the local solution, in order to extend it to a global one and to study the stability of stationary solutions to the initial-boundary value problem (1.3)-(1.4).

Proposition 3.4. *Assume the conditions of Proposition 3.3 hold. Then there exists a positive constant δ_1 , such that if $\|\varphi_0\|_{H^2}^2 + \|\psi_0\|_{H^2}^2 \leq \delta_1$, then the problem (3.12)-(3.14) admits a unique global solution $(\varphi, \psi) \in X(0, \infty)$ which satisfies that for all $t \geq 0$,*

$$\|\varphi(\cdot, t)\|_{H^2}^2 + \|\psi(\cdot, t)\|_{H^2}^2 + \int_0^t (\|\varphi\|_{H^3}^2 + \|\psi\|_{H^3}^2 + \|\varphi_\tau\|_{H^1}^2 + \|\psi_\tau\|_{H^1}^2) d\tau \leq CN^2(0). \quad (3.15)$$

for some constants $C > 0$ independent of $t \in (0, \infty)$.

By the local existence result and the standard continuation argument, to prove the Proposition 3.4, we only need to prove the requisite *a priori* estimates below.

Proposition 3.5. (*A priori estimates*). *For any $T > 0$ and any solution $(\varphi, \psi) \in X(0, T)$ to the problem (3.12)-(3.14) with $(\varphi_0, \psi_0) \in H_0^1 \cap H^2$, there exists a suitably small $\delta_2 > 0$ independent of T such that if $\|\varphi\|_{H^2}^2 + \|\psi\|_{H^2}^2 \leq \delta_2$, then (φ, ψ) satisfies (3.15) for any $t \in [0, T]$.*

Before proceeding to the estimate of (φ, ψ) , we collect some technical lemmas, which will often be used in the proof of Proposition 3.5.

Lemma 3.6. *Let (\bar{u}, \bar{w}) be the stationary solution of (1.3), (1.4) stated in Proposition 2.1. Then it holds that*

$$0 < C_1^{-1} \leq \bar{u}, \bar{w} \leq C_1 \quad (3.16)$$

for some constant $C_1 > 0$, and that

$$D\bar{w}_x^2 \leq \frac{2}{\gamma\chi} \bar{u}^\gamma. \quad (3.17)$$

Proof. According to Proposition 3.3, it is trivial to obtain (3.16) by comparison principle, so we only need to prove (3.17). For any $x \in \tilde{\mathcal{I}}$, when $0 < \bar{w}(x) \leq w_*$ there exists an $x_0 \in (0, 1)$ such that

$$0 < \bar{w}(x_0) = \min_{x \in \tilde{\mathcal{I}}} \bar{w}(x) \text{ and } \bar{w}_x(x_0) = 0.$$

Multiplying the second equation in (1.5) by \bar{w}_x , and integrating the resulting equation from x_0 to x , we get

$$\frac{D}{2} \bar{w}_x^2 = \int_{x_0}^x \bar{u}^\gamma \bar{w}_y dy = \int_{\bar{w}(x_0)}^{\bar{w}(x)} \bar{u}^\gamma d\bar{w} = \lambda^\gamma \int_{\bar{w}(x_0)}^{\bar{w}(x)} e^{\gamma\chi s} ds \leq \lambda^\gamma \int_0^{\bar{w}(x)} e^{\gamma\chi s} ds$$

with $\lambda = \frac{M}{\int_{\mathcal{I}} e^{\chi\bar{w}} dx}$, where we have used the following identity

$$\bar{u} = \frac{M}{\int_{\mathcal{I}} e^{\chi\bar{w}} dx} e^{\chi\bar{w}} =: \lambda e^{\chi\bar{w}} \quad (3.18)$$

from (2.1). Thanks to (3.18), we can get

$$\frac{D}{2} \bar{w}_x^2 \leq \frac{1}{\gamma\chi} \lambda^\gamma (e^{\gamma\chi\bar{w}} - 1) \leq \frac{1}{\gamma\chi} \lambda^\gamma e^{\gamma\chi\bar{w}} = \frac{1}{\gamma\chi} \bar{u}^\gamma.$$

The proof is completed. \square

Lemma 3.7 ([36]). *For any $a \geq -1$ and $1 < \gamma < 2$, it holds that*

$$(a+1)^\gamma - 1 - \gamma a \leq a^2.$$

Lemma 3.8 ([36]). *For any $a \geq 0$ and $0 < \gamma < 1$, it holds that*

$$|a^\gamma - 1| \leq |a - 1|.$$

In the following part, we will consider the global existence of (3.12)-(3.14). Let us begin with the estimate of the zeroth and first order frequencies of the perturbation.

Lemma 3.9. *For any solution $(\varphi, \psi) \in X(0, T)$ to the problem (3.12)-(3.14) satisfying $N(t) \ll 1$, it holds that*

$$\int_{\mathcal{I}} \frac{\varphi^2}{\bar{u}} + \bar{u}^{1-\gamma} \psi^2 dx + \int_0^t (||\psi||_{L^2}^2 + ||\varphi_x||_{L^2}^2 + ||\psi_x||_{L^2}^2) d\tau \leq C (||\varphi_0||_{L^2}^2 + ||\psi_0||_{L^2}^2) \quad (3.19)$$

for any $t \in [0, T]$, where $C > 0$ is a constant independent of t .

Proof. We Multiply the first equation in (3.12) by $\gamma \frac{\varphi}{\bar{u}}$, then integrating by parts with respect to x , we can show that

$$\begin{aligned} & \frac{\gamma}{2} \frac{d}{dt} \int_{\mathcal{I}} \frac{\varphi^2}{\bar{u}} dx + \gamma \int_{\mathcal{I}} \frac{\varphi_x^2}{\bar{u}} dx \\ &= -\gamma \int_{\mathcal{I}} \varphi \varphi_x \left[\left(\frac{1}{\bar{u}} \right)_x + \chi \frac{\bar{w}_x}{\bar{u}} \right] dx - \gamma\chi \int_{\mathcal{I}} \frac{\varphi \varphi_x \psi_x}{\bar{u}} dx - \gamma\chi \int_{\mathcal{I}} \varphi \psi_x dx. \end{aligned} \quad (3.20)$$

By using (3.10), we can show that

$$\left(\frac{1}{\bar{u}}\right)_x + \chi \frac{\bar{w}_x}{\bar{u}} = -\frac{1}{\bar{u}^2}(\bar{u}_x - \chi \bar{u} \bar{w}_x) = 0,$$

thus we get

$$-\gamma \int_{\mathcal{I}} \varphi \varphi_x \left[\left(\frac{1}{\bar{u}}\right)_x + \chi \frac{\bar{w}_x}{\bar{u}} \right] dx = 0. \quad (3.21)$$

Using the fact that $\|\varphi\|_{L^\infty} \leq N(t)$ and the Cauchy-Schwarz inequality, we can show that

$$\gamma \chi \int_{\mathcal{I}} \frac{\varphi \varphi_x \psi_x}{\bar{u}} dx \leq C \|\varphi\|_{L^\infty} \|\psi_x\|_{L^2} \|\varphi_x\|_{L^2} \leq CN(t) (\|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2). \quad (3.22)$$

Substituting (3.21) and (3.22) into (3.20), we have

$$\frac{\gamma}{2} \frac{d}{dt} \int_{\mathcal{I}} \frac{\varphi^2}{\bar{u}} dx + \gamma \int_{\mathcal{I}} \frac{\varphi_x^2}{\bar{u}} dx \leq -\gamma \chi \int_{\mathcal{I}} \varphi \psi_x dx + CN(t) (\|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2). \quad (3.23)$$

We Multiply the first equation in (3.12) by $\chi \bar{u}^{1-\gamma} \psi$ and then integrating by parts with respect to x , we get

$$\begin{aligned} \frac{\chi}{2} \frac{d}{dt} \int_{\mathcal{I}} \bar{u}^{1-\gamma} \psi^2 dx + \chi D \int_{\mathcal{I}} \bar{u}^{1-\gamma} \psi_x^2 dx &= -\chi D \int_{\mathcal{I}} (\bar{u}^{1-\gamma})_x \psi \psi_x dx \\ &\quad - \chi \int_{\mathcal{I}} \bar{u} \psi \left[\left(\frac{\varphi_x}{\bar{u}} + 1\right)^\gamma - 1 - \gamma \frac{\varphi_x}{\bar{u}} \right] dx \\ &\quad - \gamma \chi \int_{\mathcal{I}} \psi \varphi_x dx, \end{aligned} \quad (3.24)$$

For the first term on the right hand side of (3.24), by (3.14) and then integrating by parts, we can show that

$$-\chi D \int_{\mathcal{I}} (\bar{u}^{1-\gamma})_x \psi \psi_x dx = \frac{1}{2} \chi D \int_{\mathcal{I}} (\bar{u}^{1-\gamma})_{xx} \psi^2 dx, \quad (3.25)$$

by (3.10), (3.16) and (3.17), we have

$$\begin{aligned} \frac{D\chi}{2} (\bar{u}^{1-\gamma})_{xx} &= -\frac{D\chi}{2} \gamma (1-\gamma) \bar{u}^{-\gamma-1} \bar{u}_x^2 + \frac{D\chi}{2} (1-\gamma) \bar{u}^{-\gamma} \bar{u}_{xx} \\ &= \frac{D\chi}{2} (1-\gamma)^2 \bar{u}^{1-\gamma} \chi^2 \bar{w}_x^2 + \frac{D\chi}{2} (1-\gamma) \bar{u}^{1-\gamma} \chi \bar{w}_{xx} \\ &\leq \left[\frac{(1-\gamma)^2}{\gamma} + \frac{1-\gamma}{2} \right] \chi^2 \bar{u} \end{aligned}$$

Hence, when $1 < \gamma \leq 2$, we update (3.25) as

$$-\chi D \int_{\mathcal{I}} (\bar{u}^{1-\gamma})_x \psi \psi_x dx \leq \left(\frac{(1-\gamma)^2}{\gamma} + \frac{1-\gamma}{2} \right) \chi^2 \int_{\mathcal{I}} \bar{u} \psi^2 dx := -\mathcal{K} \int_{\mathcal{I}} \bar{u} \psi^2 dx, \quad (3.26)$$

where $\mathcal{K} > 0$ when $1 < \gamma < 2$ and $\mathcal{K} = 0$ when $\gamma = 2$. For the second term on the right hand side of (3.24), using (3.9), (3.16), Lemma 3.7 and the Hölder inequality, we have

$$-\chi \int_{\mathcal{I}} \bar{u} \psi \left[\left(\frac{\varphi_x}{\bar{u}} + 1\right)^\gamma - 1 - \gamma \frac{\varphi_x}{\bar{u}} \right] dx \leq C \|\psi\|_{L^\infty} \|\varphi_x\|_{L^2}^2 \leq CN(t) \|\varphi_x\|_{L^2}^2, \quad (3.27)$$

then substituting (3.26) and (3.27) into (3.24), we get

$$\begin{aligned} & \frac{\chi}{2} \frac{d}{dt} \int_{\mathcal{I}} \bar{u}^{1-\gamma} \psi^2 dx + D\chi \int_{\mathcal{I}} \bar{u}^{1-\gamma} \psi_x^2 dx + \mathcal{K} \int_{\mathcal{I}} \bar{u} \psi^2 dx \\ & \leq -\gamma\chi \int_{\mathcal{I}} \psi \varphi_x dx + CN(t) \|\varphi_x\|_{L^2}^2. \end{aligned} \quad (3.28)$$

Adding (3.28) to (3.23), we can show that

$$\frac{1}{2} \frac{d}{dt} \int_{\mathcal{I}} \gamma \frac{\varphi^2}{\bar{u}} + \chi \bar{u}^{1-\gamma} \psi^2 dx + \int_{\mathcal{I}} \mathcal{K} \bar{u} \psi^2 + \gamma \frac{\varphi_x^2}{\bar{u}} + D\chi \bar{u}^{1-\gamma} \psi_x^2 dx \leq CN(t) (\|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2).$$

By (3.16), we obtain

$$\begin{aligned} & \frac{1}{2} \min\{\gamma, \chi\} \frac{d}{dt} \int_{\mathcal{I}} \frac{\varphi^2}{\bar{u}} + \bar{u}^{1-\gamma} \psi^2 dx + \min\{C_1 \mathcal{K}, \frac{\gamma}{C_1}, D\chi C_1^{1-\gamma}\} \int_{\mathcal{I}} \psi^2 + \varphi_x^2 + \psi_x^2 dx \\ & \leq CN(t) (\|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2), \end{aligned}$$

which implies that

$$\frac{d}{dt} \int_{\mathcal{I}} \frac{\varphi^2}{\bar{u}} + \bar{u}^{1-\gamma} \psi^2 dx + \beta (\|\psi\|_{L^2}^2 + \|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2) \leq 0 \quad (3.29)$$

with

$$CN(t) \leq \frac{\min\{\frac{\gamma}{C_1}, D\chi C_1^{1-\gamma}, C_1 \mathcal{K}\}}{2 \min\{\gamma, \chi\}} =: \frac{\beta}{2}.$$

Integrating (3.29) over $(0, t)$, we get (3.19) and hence completes the proof of the lemma. \square

We are ready to derive the estimate on φ_x and ψ_x in the next lemma.

Lemma 3.10. *For any solution $(\varphi, \psi) \in X(0, T)$ to the problem (3.12)-(3.14) satisfying $N(t) \ll 1$, it holds that for any $t \in [0, T]$ that*

$$\|\varphi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^2 + \int_0^t \|\varphi_\tau\|_{L^2}^2 + \|\psi_\tau\|_{L^2}^2 d\tau \leq C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2). \quad (3.30)$$

Proof. Multiplying the first equation of (3.12) by φ_t , then integrating by parts, we can show that

$$\frac{1}{2} \frac{d}{dt} \int_{\mathcal{I}} \varphi_x^2 dx + \int_{\mathcal{I}} \varphi_t^2 dx = -\chi \int_{\mathcal{I}} \bar{u} \varphi_t \psi_x dx - \chi \int_{\mathcal{I}} \bar{w}_x \varphi_t \varphi_x dx - \chi \int_{\mathcal{I}} \varphi_t \varphi_x \psi_x dx, \quad (3.31)$$

Using (3.16), (3.17) and the Cauchy-Schwarz inequality, we have

$$-\chi \int_{\mathcal{I}} \bar{w}_x \varphi_t \varphi_x dx \leq \chi \|\bar{w}_x\|_{L^\infty} \|\varphi_t\|_{L^2} \|\varphi_x\|_{L^2} \leq \varepsilon \|\varphi_t\|_{L^2}^2 + C_\varepsilon \|\varphi_x\|_{L^2}^2 \quad (3.32)$$

and

$$-\chi \int_{\mathcal{I}} \bar{u} \varphi_t \psi_x dx \leq \chi \|\bar{u}\|_{L^\infty} \|\varphi_t\|_{L^2} \|\psi_x\|_{L^2} \leq \varepsilon \|\varphi_t\|_{L^2}^2 + C_\varepsilon \|\psi_x\|_{L^2}^2, \quad (3.33)$$

for any $\varepsilon > 0$. Using $\|\varphi_x\|_{L^\infty} \leq N(t)$ and the Cauchy-Schwarz inequality, we can show that

$$-\chi \int_{\mathcal{I}} \varphi_t \varphi_x \psi_x dx \leq \chi \|\varphi_x\|_{L^\infty} \|\varphi_t\|_{L^2} \|\psi_x\|_{L^2} \leq CN(t) (\|\varphi_t\|_{L^2}^2 + \|\psi_x\|_{L^2}^2). \quad (3.34)$$

Then we substitute (3.32)-(3.34) into (3.31), by choosing ε suitably small and letting $N(t)$ small enough such that

$$CN(t) \leq \frac{1}{4},$$

we obtain

$$\frac{d}{dt} \int_{\mathcal{I}} \varphi_x^2 dx + \int_{\mathcal{I}} \varphi_t^2 dx \leq C \|\psi_x\|_{L^2}^2. \quad (3.35)$$

Using (3.19) and integrating by parts, we update (3.35) as

$$\begin{aligned} \|\varphi_x\|_{L^2}^2 + \int_0^t \|\varphi_\tau\|_{L^2}^2 d\tau &\leq \|\varphi_{0,x}\|_{L^2}^2 + C \int_0^t \|\psi_x\|_{L^2}^2 d\tau \\ &\leq \|\varphi_{0,x}\|_{L^2}^2 + C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{L^2}^2) \\ &\leq C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{L^2}^2). \end{aligned} \quad (3.36)$$

Multiplying the second equation of (3.12) by ψ_t and then integrating the resultant equation with respect to x , we can show that

$$\frac{1}{2} \frac{d}{dt} D \int_{\mathcal{I}} \psi_x^2 dx + \int_{\mathcal{I}} \psi_t^2 dx = - \int_{\mathcal{I}} \bar{u}^\gamma \psi_t \left[\left(\frac{\varphi_x}{\bar{u}} + 1 \right)^\gamma - 1 - \gamma \frac{\varphi_x}{\bar{u}} \right] dx - \gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} \psi_t \varphi_x dx. \quad (3.37)$$

For the two terms on the right side of (3.37), by (3.9), (3.16), Lemma 3.7 and the Cauchy-Schwarz inequality, we can show that

$$\begin{aligned} - \int_{\mathcal{I}} \bar{u}^\gamma \psi_t \left[\left(\frac{\varphi_x}{\bar{u}} + 1 \right)^\gamma - 1 - \gamma \frac{\varphi_x}{\bar{u}} \right] dx &\leq \int_{\mathcal{I}} \bar{u}^{\gamma-2} |\psi_t| \varphi_x^2 dx \\ &\leq C \|\varphi_x\|_{L^\infty} \|\psi_t\|_{L^2} \|\varphi_x\|_{L^2} \\ &\leq CN(t) (\|\psi_t\|_{L^2}^2 + \|\varphi_x\|_{L^2}^2) \end{aligned}$$

and

$$\begin{aligned} -\gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} \psi_t \varphi_x dx &\leq \gamma \|\bar{u}^{\gamma-1}\|_{L^\infty} \|\psi_t\|_{L^2} \|\varphi_x\|_{L^2} \leq C \|\psi_t\|_{L^2} \|\varphi_x\|_{L^2} \\ &\leq \varepsilon \|\psi_t\|_{L^2}^2 + C_\varepsilon \|\varphi_x\|_{L^2}^2 \end{aligned}$$

for any $\varepsilon > 0$. Choosing ε and $N(t)$ small enough, we thus update (3.37) as

$$D \frac{d}{dt} \int_{\mathcal{I}} \psi_x^2 dx + \int_{\mathcal{I}} \psi_t^2 dx \leq (CN(t) + C_\varepsilon) \|\varphi_x\|_{L^2}^2. \quad (3.38)$$

Using (3.19) and integration by parts, for any $t \in [0, T]$, it implies that

$$\begin{aligned} D \int_{\mathcal{I}} \psi_x^2 dx + \int_0^t \|\psi_\tau\|_{L^2}^2 d\tau &\leq \|\psi_{0,x}\|_{L^2}^2 + (CN(t) + C_\varepsilon) \int_0^t \|\varphi_x\|_{L^2}^2 d\tau \\ &\leq C \|\psi_{0,x}\|_{L^2}^2 + C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{L^2}^2), \end{aligned}$$

thus

$$\|\psi_x\|_{L^2}^2 + \int_0^t \|\psi_\tau\|_{L^2}^2 d\tau \leq C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2). \quad (3.39)$$

Adding (3.39) to (3.36), we complete the proof of Lemma 3.9. \square

In the following, we establish the estimates of the second order derivatives of (φ, ψ) .

Lemma 3.11. *Let $(\varphi, \psi) \in X(0, T)$ be a solution to the initial-boundary value problem (3.12)-(3.14) and assume the conditions of Lemma 3.9 hold. Then it holds for any $t \in [0, T]$ that*

$$\|\varphi_{xx}\|_{L^2}^2 + \|\psi_{xx}\|_{L^2}^2 + \int_0^t \int_{\mathcal{I}} \varphi_{xxx}^2 + \psi_{xxx}^2 dx d\tau \quad (3.40)$$

$$\leq C (\|\varphi_0\|_{H^2}^2 + \|\psi_0\|_{H^2}^2) \quad (3.41)$$

Proof. By differentiating (3.12) with respect to t to get

$$\begin{cases} \varphi_{tt} = \varphi_{xxt} - \chi(\bar{w}_x \varphi_{xt} + \bar{u} \psi_{xt} + \varphi_{xt} \psi_x + \varphi_x \psi_{xt}), \\ \psi_{tt} = D\psi_{xxt} - \gamma(\varphi_x + \bar{u})^{\gamma-1} \varphi_{xt}. \end{cases} \quad (3.42)$$

Multiplying the first equation in (3.42) by φ_t and the second one by ψ_t , then integrate by parts over \mathcal{I} , and adding the results to show that

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathcal{I}} \varphi_t^2 + \psi_t^2 dx + \int_{\mathcal{I}} \varphi_{xt}^2 + D\psi_{xt}^2 dx = \\ & -\chi \int_{\mathcal{I}} \bar{w}_x \varphi_t \varphi_{xt} dx - \chi \int_{\mathcal{I}} \bar{u} \varphi_t \psi_{xt} dx - \chi \int_{\mathcal{I}} \varphi_t \varphi_{xt} \psi_x dx - \chi \int_{\mathcal{I}} \varphi_t \varphi_x \psi_{xt} dx \\ & - \gamma \int_{\mathcal{I}} (\varphi_x + \bar{u})^{\gamma-1} \psi_t \varphi_{xt} dx, \end{aligned} \quad (3.43)$$

where we have used (3.14) to get $\varphi_t(0, t) = \varphi_t(1, t) = 0$. Now, we estimate the right hand side of (3.43). Using (3.16), (3.17) and the Cauchy-Schwarz inequality, we can show that

$$-\chi \int_{\mathcal{I}} \bar{w}_x \varphi_t \varphi_{xt} dx - \chi \int_{\mathcal{I}} \bar{u} \varphi_t \psi_{xt} dx \leq C_\varepsilon \|\varphi_t\|_{L^2}^2 + \varepsilon (\|\varphi_{xt}\|_{L^2}^2 + \|\psi_{xt}\|_{L^2}^2) \quad (3.44)$$

for any $\varepsilon > 0$. Again, by using $\|\varphi_x\|_{L^\infty}, \|\psi_x\|_{L^\infty} \leq N(t)$ and the Cauchy-Schwarz inequality, we have

$$\begin{aligned} -\chi \int_{\mathcal{I}} \varphi_t \varphi_{xt} \psi_x dx - \chi \int_{\mathcal{I}} \varphi_t \varphi_x \psi_{xt} dx & \leq \chi \|\psi_x\|_{L^\infty} \|\varphi_t\|_{L^2} \|\varphi_{xt}\|_{L^2} + \chi \|\varphi_x\|_{L^\infty} \|\varphi_t\|_{L^2} \|\psi_{xt}\|_{L^2} \\ & \leq CN(t) (\|\varphi_t\|_{L^2}^2 + \|\varphi_{xt}\|_{L^2}^2 + \|\psi_{xt}\|_{L^2}^2). \end{aligned} \quad (3.45)$$

For the last term on the right hand side of (3.43), using (3.9), Lemma 3.8 and the Cauchy-Schwarz inequality we can show that

$$\begin{aligned} -\gamma \int_{\mathcal{I}} (\varphi_x + \bar{u})^{\gamma-1} \psi_t \varphi_{xt} dx & = -\gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} \left(\frac{\varphi_x}{\bar{u}} + 1 \right)^{\gamma-1} \psi_t \varphi_{xt} dx \\ & \leq \gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} \left| \left(\frac{\varphi_x}{\bar{u}} + 1 \right)^{\gamma-1} - 1 \right| |\psi_t| |\varphi_{xt}| dx + \gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} |\psi_t| |\varphi_{xt}| dx \\ & \leq \gamma \int_{\mathcal{I}} \bar{u}^{\gamma-2} |\varphi_x| |\psi_t| |\varphi_{xt}| dx + \gamma \int_{\mathcal{I}} \bar{u}^{\gamma-1} |\psi_t| |\varphi_{xt}| dx \\ & \leq C \|\varphi_x\|_{L^\infty} \|\psi_t\|_{L^2} \|\varphi_{xt}\|_{L^2} + C_\varepsilon \|\psi_t\|_{L^2}^2 + \varepsilon \|\varphi_{xt}\|_{L^2}^2 \\ & \leq CN(t) (\|\psi_t\|_{L^2}^2 + \|\varphi_{xt}\|_{L^2}^2) + C_\varepsilon \|\psi_t\|_{L^2}^2 + \varepsilon \|\varphi_{xt}\|_{L^2}^2 \\ & \leq (C_\varepsilon + CN(t)) \|\psi_t\|_{L^2}^2 + (\varepsilon + CN(t)) \|\varphi_{xt}\|_{L^2}^2. \end{aligned} \quad (3.46)$$

Substituting (3.44)-(3.46) into (3.43), we update (3.43) as

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\mathcal{I}} \varphi_t^2 + \psi_t^2 dx + \int_{\mathcal{I}} \varphi_{xt}^2 + D\psi_{xt}^2 dx \\ & \leq (C_\varepsilon + CN(t)) (\|\varphi_t\|_{L^2}^2 + \|\psi_t\|_{L^2}^2) + (\varepsilon + CN(t)) (\|\varphi_{xt}\|_{L^2}^2 + \|\psi_{xt}\|_{L^2}^2). \end{aligned}$$

Then we choose ε and δ small enough such that

$$\varepsilon + CN(t) \leq \frac{1}{2}, \quad (3.47)$$

thus

$$\frac{d}{dt} \int_{\mathcal{I}} \varphi_t^2 + \psi_t^2 dx + \int_{\mathcal{I}} \varphi_{xt}^2 + D\psi_{xt}^2 dx \leq (C_\varepsilon + C\delta) (\|\varphi_t\|_{L^2}^2 + \|\psi_t\|_{L^2}^2). \quad (3.48)$$

Combine (3.39), (3.36) and (3.48), we can show that

$$\begin{aligned} & \int_{\mathcal{I}} \varphi_t^2 + \psi_t^2 dx + \int_0^t \int_{\mathcal{I}} \varphi_{x\tau}^2 + \psi_{x\tau}^2 dx d\tau \\ & \leq C \int_{\mathcal{I}} (\varphi_{0,xx}^2 + \varphi_{0,x}^2 + \psi_{0,xx}^2 + \psi_{0,x}^2 + \varphi_{0,x}^4 + \psi_{0,x}^4) dx \\ & \quad + (\varepsilon + CN(t)) \int_0^t \|\varphi_\tau\|_{L^2}^2 + \|\psi_\tau\|_{L^2}^2 d\tau \\ & \leq C \left(\|\varphi_{0,x}\|_{H^1}^2 + \|\psi_{0,x}\|_{H^1}^2 + (\|\varphi_{0,x}\|_{L^2}^2 + \|\psi_{0,x}\|_{L^2}^2)^2 \right) \\ & \quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2), \end{aligned} \quad (3.49)$$

where we have used the following inequality

$$\begin{aligned} \int_{\mathcal{I}} \varphi_t^2 dx|_{t=0} & \leq C \int_{\mathcal{I}} (\varphi_{0,xx}^2 + \varphi_{0,x}^2 + \psi_{0,x}^2 + \varphi_{0,x}^2 \psi_{0,x}^2) dx \\ & \leq C \int_{\mathcal{I}} (\varphi_{0,xx}^2 + \varphi_{0,x}^2 + \psi_{0,x}^2 + \varphi_{0,x}^4 + \psi_{0,x}^4) dx \end{aligned} \quad (3.50)$$

and

$$\begin{aligned} \int_{\mathcal{I}} \psi_t^2 dx|_{t=0} & \leq \int_{\mathcal{I}} (D\psi_{0,xx}^2 + \bar{u}^\gamma - (\varphi_{0,x} + \bar{u})^\gamma)^2 dx \\ & \leq C \int_{\mathcal{I}} \psi_{0,xx}^2 dx + C \int_{\mathcal{I}} ((\varphi_{0,x} + \bar{u})^\gamma - \bar{u}^\gamma)^2 dx \\ & \leq C \int_{\mathcal{I}} \psi_{0,xx}^2 dx + C \int_{\mathcal{I}} (\gamma \bar{u}^{\gamma-1} \varphi_{0,x} + o(\varphi_{0,x}))^2 dx \\ & \leq C (\|\psi_{0,xx}\|_{L^2}^2 + \|\varphi_{0,x}\|_{L^2}^2), \end{aligned} \quad (3.51)$$

in which we have also used $\varphi_t|_{t=0} = \varphi_{0,xx} - \chi(\varphi_{0,x}\psi_{0,x} + \bar{u}\psi_{0,x} + \bar{w}_x\varphi_{0,x})$ and $\psi_t|_{t=0} = D\psi_{0,xx} + \bar{u}^\gamma - (\varphi_{0,x} + \bar{u})^\gamma$ from (3.12). For the first equation in (3.12) and using the Cauchy-Schwarz inequality, we get from (3.8), (3.16) and (3.17) that

$$\begin{aligned} \int_{\mathcal{I}} \varphi_{xx}^2 dx & \leq C \int_{\mathcal{I}} (\varphi_t^2 + \varphi_x^2 + \psi_x^2) dx + \chi \|\varphi_x\|_{L^\infty}^2 \int_{\mathcal{I}} \psi_x^2 dx \\ & \leq C \int_{\mathcal{I}} (\varphi_t^2 + \varphi_x^2 + \psi_x^2) dx + C (\|\varphi_x\|_{L^2} \|\varphi_{xx}\|_{L^2} + \|\varphi_x\|_{L^2}^2) \int_{\mathcal{I}} \psi_x^2 dx \\ & \leq C \int_{\mathcal{I}} (\varphi_t^2 + \varphi_x^2 + \psi_x^2) dx + \frac{1}{2} \|\varphi_{xx}\|_{L^2}^2 + C \|\varphi_x\|_{L^2}^2 (\|\psi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^4), \end{aligned}$$

thus

$$\int_{\mathcal{I}} \varphi_{xx}^2 dx \leq C \int_{\mathcal{I}} (\varphi_t^2 + \varphi_x^2 + \psi_x^2) dx. \quad (3.52)$$

This together with (3.19), (3.36), (3.39) and (3.49) can obtain that

$$\begin{aligned}
\int_{\mathcal{I}} \varphi_{xx}^2 dx &\leq C \left(\|\varphi_{0,x}\|_{H^1}^2 + \|\psi_{0,x}\|_{H^1}^2 + (\|\varphi_{0,x}\|_{L^2}^2 + \|\psi_{0,x}\|_{L^2}^2)^2 \right) + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2) \\
&\quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{L^2}^2) \left(\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2 + (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 \right) \\
&\leq C \left(\|\varphi_{0,x}\|_{H^1}^2 + \|\psi_{0,x}\|_{H^1}^2 + (\|\varphi_{0,x}\|_{L^2}^2 + \|\psi_{0,x}\|_{L^2}^2)^2 \right) \\
&\quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2) \left(1 + \|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2 + (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 \right) \\
&\leq C \left(\|\varphi_{0,x}\|_{H^1}^2 + \|\psi_{0,x}\|_{H^1}^2 + (\|\varphi_{0,x}\|_{L^2}^2 + \|\psi_{0,x}\|_{L^2}^2)^2 \right) \\
&\quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2) (1 + \|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2
\end{aligned} \tag{3.53}$$

and

$$\begin{aligned}
\int_0^t \int_{\mathcal{I}} \varphi_{xx}^2 dx d\tau &\leq C \int_0^t \int_{\mathcal{I}} (\varphi_\tau^2 + \varphi_x^2 + \psi_x^2) dx d\tau + C \int_0^t \|\varphi_x\|_{L^2}^2 (\|\psi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^4) d\tau \\
&\leq C \int_0^t \int_{\mathcal{I}} (\varphi_\tau^2 + \varphi_x^2 + \psi_x^2) dx d\tau + C \sup_{\tau \in [0,t]} (\|\psi_x\|_{L^2}^2 + \|\psi_x\|_{L^2}^4) \int_0^t \|\varphi_x\|_{L^2}^2 d\tau \\
&\leq C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{L^2}^2) \\
&\quad + C \left(\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2 + (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 \right) (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{L^2}^2) \\
&\leq C \left(1 + \|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2 + (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 \right) (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{L^2}^2) \\
&\leq C (1 + \|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{L^2}^2).
\end{aligned} \tag{3.54}$$

By using the same method and utilizing (3.14), (3.19), (3.36), (3.39), (3.49) and the second equation in (3.12), we obtain

$$\begin{aligned}
\int_{\mathcal{I}} \psi_{xx}^2 dx &= \frac{1}{D^2} \int_{\mathcal{I}} (\psi_t - \bar{u}^\gamma + (\varphi_x + \bar{u})^\gamma)^2 dx \\
&\leq C \int_{\mathcal{I}} \psi_t^2 dx + C \int_{\mathcal{I}} ((\varphi_x + \bar{u})^\gamma - \bar{u}^\gamma)^2 dx \\
&\leq C \int_{\mathcal{I}} \psi_t^2 dx + C \int_{\mathcal{I}} (\gamma \bar{u}^{\gamma-1} \varphi_x + o(\varphi_x))^2 dx \\
&\leq C \int_{\mathcal{I}} \psi_t^2 dx + C \int_{\mathcal{I}} \varphi_x^2 dx
\end{aligned} \tag{3.55}$$

and

$$\begin{aligned}
\int_0^t \int_{\mathcal{I}} \psi_{xx}^2 dx d\tau &\leq C \int_0^t \|\psi_\tau\|_{L^2}^2 d\tau + C \int_0^t \|\varphi_x\|_{L^2}^2 d\tau \\
&\leq C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2) + C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{L^2}^2) \\
&\leq C (\|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2).
\end{aligned} \tag{3.56}$$

Combine (3.49), (3.53) and (3.55), we get

$$\begin{aligned}
& \|\varphi_{xx}\|_{L^2}^2 + \|\psi_{xx}\|_{L^2}^2 + \int_0^t \int_{\mathcal{I}} \varphi_{x\tau}^2 + \psi_{x\tau}^2 dx d\tau \\
& \leq C \left(\|\varphi_{0,x}\|_{H^1}^2 + \|\psi_{0,x}\|_{H^1}^2 + (\|\varphi_{0,x}\|_{L^2}^2 + \|\psi_{0,x}\|_{L^2}^2)^2 \right) \\
& \quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2) (1 + \|\varphi_0\|_{L^2}^2 + \|\psi_0\|_{H^1}^2)^2 \\
& \quad + C (\|\varphi_0\|_{H^1}^2 + \|\psi_0\|_{H^1}^2) \\
& \leq \|\varphi_0\|_{H^2}^2 + \|\psi_0\|_{H^2}^2.
\end{aligned} \tag{3.57}$$

Next we differentiate the first equation in (3.12) with respect to x to obtain

$$\varphi_{xxx} = \varphi_{xt} + \chi(\bar{w}_{xx}\varphi_x + \bar{w}_x\varphi_{xx} + \bar{u}_x\psi_x + \bar{u}\psi_{xx} + \varphi_{xx}\psi_x + \varphi_x\psi_{xx}),$$

then we combine (3.16), (3.17), (3.19), (3.54) and (3.57) the Sobolev inequality (3.8) to show that

$$\int_0^t \int_{\mathcal{I}} \varphi_{xxx}^2 dx d\tau \leq C (\|\varphi_0\|_{H^2}^2 + \|\psi_0\|_{H^2}^2). \tag{3.58}$$

Similarly, using (3.9), (3.16), (3.17), (3.19), (3.54) and (3.57) we have

$$\int_0^t \int_{\mathcal{I}} \psi_{xxx}^2 dx d\tau \leq C (\|\varphi_0\|_{H^2}^2 + \|\psi_0\|_{H^2}^2). \tag{3.59}$$

Combining (3.54), (3.56), (3.57), (3.58) and (3.59), we then get (3.40) and complete the proof of Lemma 3.11. \square

Utilizing the *a priori* estimates established in §3.2, the global well-posedness of the initial and boundary value problem (3.12)–(3.14) can be proved by combining Proposition 3.3 and standard continuation argument. To complete the proof of Theorem 2.2, it remains to derive the energy estimates leading to the long-time behavior of the solution.

3.3. Long time behavior. In this section, we are ready to prove (2.2).

Step 1. To prove the convergence of the zeroth order frequency of the perturbation, we note that under the conditions of Theorem 2.2, Lemma 3.9 implies

$$(\|\varphi_x(t)\|_{L^2}^2 + \|\psi(t)\|_{L^2}^2 + \|\psi_x(t)\|_{L^2}^2) \in L^1(0, \infty). \tag{3.60}$$

Applying the arguments in the proof of Lemma 3.9 to (3.29), Lemma 3.10 to (3.35) and (3.38), we can show that

$$\left| \frac{d}{dt} (\|\varphi_x(t)\|_{L^2}^2 + \|\psi(t)\|_{L^2}^2 + \|\psi_x(t)\|_{L^2}^2) \right| \leq C (\|\varphi_x\|_{H^1}^2 + \|\psi_x\|_{H^1}^2). \tag{3.61}$$

Integrating (3.61) with respect to time and applying Lemmas 3.9–3.10 yield

$$\frac{d}{dt} (\|\varphi_x(t)\|_{L^2}^2 + \|\psi_x(t)\|_{L^2}^2) \in L^1(0, \infty). \tag{3.62}$$

Since a positive function $f(t) \in W^{1,1}(0, \infty)$ converges to zero as $t \rightarrow \infty$, (3.60) and (3.62) imply

$$(\|\varphi_x(t)\|_{L^2}^2 + \|\psi(t)\|_{L^2}^2 + \|\psi_x(t)\|_{L^2}^2) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \tag{3.63}$$

Since $\psi = w - \bar{w}$ and $(\varphi_x, \psi_x) = (u - \bar{u}, w_x - \bar{w}_x)$, it follows from (3.63) that

$$(\|(u - \bar{u})(t)\|_{L^2}^2 + \|(w - \bar{w})(t)\|_{H^1}^2) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \tag{3.64}$$

Step 2. To show the convergence of the first order spatial derivative of the perturbation, we first note that Lemma 3.10 imply that

$$(\|\varphi_t\|_{L^2}^2 + \|\psi_t\|_{L^2}^2) \in L^1(0, \infty). \quad (3.65)$$

Moreover, applying the arguments in the proof of Lemma 3.11 to (3.48), we can show that

$$\begin{aligned} & \left| \frac{d}{dt} (\|\varphi_t(t)\|_{L^2}^2 + \|\psi_t(t)\|_{L^2}^2) \right| \\ & \leq C(\|\varphi_t\|_{H^1}^2 + \|\psi_t\|_{H^1}^2). \end{aligned} \quad (3.66)$$

Integrating (3.66) with respect to time and applying Lemmas 3.10–3.11, we obtain

$$\frac{d}{dt} (\|\varphi_t(t)\|_{L^2}^2 + \|\psi_t(t)\|_{L^2}^2) \in L^1(0, \infty). \quad (3.67)$$

It follows from (3.65) and (3.67) that

$$(\|\varphi_t(t)\|_{L^2}^2 + \|\psi_t(t)\|_{L^2}^2) \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (3.68)$$

As a direct consequence of (3.52), (3.63) and (3.68), we have

$$\|\varphi_{xx}(t)\|_{L^2}^2 \rightarrow 0 \quad \text{as } t \rightarrow \infty, \quad (3.69)$$

which implies

$$\|(u_x - \bar{u}_x)(t)\|_{L^2}^2 \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (3.70)$$

which, together with (3.48), (3.63) and (3.68), implies

$$\|\psi_{xx}(t)\|_{L^2}^2 \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (3.71)$$

It then follows that

$$\|(w_{xx} - \bar{w}_{xx})(t)\|_{L^2}^2 \rightarrow 0 \quad \text{as } t \rightarrow \infty. \quad (3.72)$$

This finishes the proof of Theorem 2.2.

ACKNOWLEDGEMENTS

Support for this work came in part from National Natural Science Foundation of China Awards 12101095 (ZF) and 11971082 (SZ), Natural Science Foundation of Chongqing Awards cstc2021jcyj-msxmX0224 (ZF) and csts2020jcyj-jqX0022 (SZ), Science and Technology Research Program of Chongqing Municipal Education Commission Awards KJQN202100517 (ZF) and KJZD-M201900501(SZ), a research project of Chongqing Education Commission CXQT21014 (ZF), a grant of Chongqing Young Experts' Workshop (ZF), and from Chongqing Youth Talent Support Program Award cstc2021ycjh-bgzxm0130 (SZ).

REFERENCES

- [1] J. Adler, *Chemotaxis in bacteria*, Science, 153 (1966), pp. 708-716.
- [2] N. Bellomo, A. Bellouquid, Y. Tao, and M. Winkler, *Toward a mathematical theory of Keller-Segel models of pattern formation in biological tissues*, Math. Models Methods Appl. Sci., 25 (2015), pp. 1663-1763.
- [3] J. Bonner, *The Cellular Slime Molds*, 2nd ed, Princeton University Press, 1967.
- [4] M. Braukhoff and J. Lankeit, *Stationary solutions to a chemotaxis-consumption model with realistic boundary conditions for the oxygen*, Math. Models Methods Appl. Sci., 29 (2019), pp. 2033-2062.
- [5] J. Carrillo, J. Li, and Z.-A. Wang, *Boundary spike-layer solutions of the singular keller-segel system: existence and stability*, Proc. Lond. Math. Soc., (Doi:10.1112/plms.12319, 2020).

- [6] M. Chae and K. Choi, *Nonlinear stability of planar traveling waves in a chemotaxis model of tumor angiogenesis with chemical diffusion*, J. Differential Equations, 268 (2020), pp. 3449-3496.
- [7] K. Choi, M.-J. Kang, and A. F. Vasseur, *Global well-posedness of large perturbations of traveling waves in a hyperbolic-parabolic system arising from a chemotaxis model*, J. Math. Pures Appl. (9), 142 (2020), pp. 266-297.
- [8] P. Davis, P. van Heijster, and R. Marangell, *Absolute instabilities of travelling wave solutions in a keller-segel model*, Nonlinearity, 30 (2017), p. 4029.
- [9] S. Dehaene, *The neural basis of the Weber-Fechner law: a logarithmic mental number line*, Trends Cogn. Sci., 7 (2003), pp. 145-147.
- [10] L. Fan, H.-Y. Jin, *Global existence and asymptotic behavior to a chemotaxis system with consumption of chemoattractant in higher dimensions*, J. Math. Phys. 58 (2017), 011503.
- [11] Z.-F. Feng, K. Zhao, and S.-M. Zhou, *Existence and stability of boundary spike layer solutions of an attractive chemotaxis model with singular sensitivity and nonlinear consumption rate of chemical stimuli*, Preprint.
- [12] Z.-F. Feng, J. Xu, L. Xue, and K. Zhao, *Initial and boundary value problem for a system of balance laws from chemotaxis: global dynamics and diffusivity limit*, Ann. Appl. Math. 37 (2021), pp. 61-110.
- [13] G.-Y. Hong and Z.-A. Wang, *Asymptotic stability of exogenous chemotaxis systems with physical boundary conditions*, Quart. Appl. Math. 79 (2021), pp. 717-743.
- [14] Q. Hou, C.-J. Liu, Y.-G. Wang, and Z. Wang, *Stability of boundary layers for a viscous hyperbolic system arising from chemotaxis: one-dimensional case*, SIAM J. Math. Anal., 50 (2018), pp. 3058-3091.
- [15] Q. Hou and Z. Wang, *Convergence of boundary layers for the Keller-Segel system with singular sensitivity in the half-plane*, J. Math. Pures Appl. (9), 130 (2019), pp. 251-287.
- [16] Y. Kalinin, L. Jiang, Y. Tu, and M. Wu, *Logarithmic sensing in escherichia coli bacterial chemotaxis*, Biophys. J., 96 (2009), pp. 2439-2448.
- [17] E. Keller and G. Odell, *Necessary and sufficient conditions for chemotactic bands*, Math. Biosci., 27 (1975), pp. 309-317.
- [18] E. F. Keller and L. A. Segel, *Initiation of slime mold aggregation viewed as an instability*, J. Theor. Biol., 26 (1970), pp. 399-415.
- [19] E. F. Keller and L. A. Segel, *Traveling bands of chemotactic bacteria: a theoretical analysis*, J. Theor. Biol., 30 (1971), pp. 235-248.
- [20] C.-C. Lee, Z. Wang, and W. Yang, *Boundary-layer profile of a singularly perturbed nonlocal semi-linear problem arising in chemotaxis*, Nonlinearity, 33 (2020), pp. 5111-5141.
- [21] H. A. Levine, B. D. Sleeman, and M. Nilsen-Hamilton, *A mathematical model for the roles of pericytes and macrophages in the initiation of angiogenesis. I. The role of protease inhibitors in preventing angiogenesis*, Math. Biosci., 168 (2000), pp. 77-115.
- [22] H. Li and K. Zhao, *Initial-boundary value problems for a system of hyperbolic balance laws arising from chemotaxis*, J. Differential Equations, 258 (2015), pp. 302-338.
- [23] J. Li, T. Li, and Z. Wang, *Stability of traveling waves of the Keller-Segel system with logarithmic sensitivity*, Math. Models Methods Appl. Sci., 24 (2014), pp. 2819-2849.
- [24] T. Li and Z. Wang, *Nonlinear stability of traveling waves to a hyperbolic-parabolic system modeling chemotaxis*, SIAM J. Appl. Math., 70 (2009), pp. 1522-1541.
- [25] T. Li and Z. Wang, *Asymptotic nonlinear stability of traveling waves to conservation laws arising from chemotaxis*, J. Differential Equations, 250 (2011), pp. 1310-1333.
- [26] T. Nagai and T. Ikeda, *Traveling waves in a chemotactic model*, J. Math. Biol., 30 (1991), pp. 169-184.
- [27] L. Nirenberg, *An extended interpolation inequality*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3), 20 (1966), pp. 733-737.
- [28] T. Nishida, *Nonlinear hyperbolic equations and related topics in fluid dynamics*, Département de Mathématique, Université de Paris-Sud, Orsay, 1978. Publications Mathématiques d'Orsay, No. 78-02.
- [29] H. Peng and Z.-A. Wang, *Nonlinear stability of strong traveling waves for the singular keller-segel system with large perturbations*, J. Differential Equations, 265 (2018), pp. 2577-2613.
- [30] H. R. Schwetlick, *Travelling fronts for multidimensional nonlinear transport equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 17 (2000), pp. 523-550.
- [31] Y. Tao, *Boundedness in a chemotaxis model with oxygen consumption by bacteria*, J. Math. Anal. Appl. 381 (2011), pp. 521C9.

- [32] Y. Tao and M. Winkler, *Eventual smoothness and stabilization of large-data solutions in a three dimensional chemotaxis system with consumption of chemoattractant*, J. Differential Equations, 252 (2012), 2520C43.
- [33] Y. Wang, C.-L. Chen, and M. Iijima, *Signaling mechanisms for chemotaxis*, Development, Growth & Differentiation, 53 (2011), pp. 495-502.
- [34] Z.-A. Wang, *Mathematics of traveling waves in chemotaxis-review paper*, Discrete Contin. Dyn. Syst. Ser. B, 18 (2013), pp. 601-641.
- [35] M. Winkler, *Singular structure formation in a degenerate haptotaxis model involving myopic diffusion*, J. Math. Pures Appl. (9), 112 (2018), pp. 118-169.
- [36] N. Zhu, Z. Liu, V. R. Martinez, and K. Zhao, *Global Cauchy problem of a system of parabolic conservation laws arising from a Keller-Segel type chemotaxis model*, SIAM J. Math. Anal. 50 (2018), pp. 5380-5425.

(Z. Feng) SCHOOL OF MATHEMATICAL SCIENCES, CHONGQING NORMAL UNIVERSITY, CHONGQING 400047, P. R. CHINA

E-mail address: zefufeng@mails.cnu.edu.cn

(J. Jia) SCHOOL OF MATHEMATICAL SCIENCES, CHONGQING NORMAL UNIVERSITY, CHONGQING 400047, P. R. CHINA

E-mail address: jj9702112022@163.com

(S. Zhou) SCHOOL OF MATHEMATICAL SCIENCES, CHONGQING NORMAL UNIVERSITY, CHONGQING 400047, P. R. CHINA

E-mail address: zhoushouming76@163.com